



Perspective

Safe hospital preparedness in the era of COVID-19: The Swiss cheese model



Ji Yun Noh, Joon Young Song*, Jin Gu Yoon, Hye Seong, Hee Jin Cheong, Woo Joo Kim

Division of Infectious Diseases, Department of Internal Medicine, Korea University College of Medicine, Seoul, Republic of Korea

ARTICLE INFO

Article history:

Received 22 May 2020

Received in revised form 26 June 2020

Accepted 26 June 2020

Keywords:

COVID-19

SARS-CoV-2

Infection control

Health personnel

Hospital planning

ABSTRACT

Since it first emerged in December 2019, coronavirus disease 2019 (COVID-19) has spread rapidly worldwide. During the pandemic of an emerging infectious disease, it is very important to prevent nosocomial outbreaks and operate hospitals safely to maintain their functions. In this article, we present the strategies for safe hospital operations based on the experiences of the Republic of Korea early in the COVID-19 pandemic. Each hospital should maintain multiple layers of defenses to prevent even small cracks in the hospital's quarantine system.

© 2020 The Author(s). Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

During the 2015 Middle East respiratory syndrome (MERS) outbreak in South Korea, most cases were hospital-acquired infections, and 21% of them occurred among healthcare personnel (HCP) (Choi et al., 2016). Based on the experience gained during the 2015 MERS outbreak, the health authorities and individual hospitals in South Korea implemented enhanced responses to the coronavirus disease 2019 (COVID-19) pandemic. With the implementation of a reinforced hospital action plan, nosocomial transmission of COVID-19 was quite limited (Supplementary Material Figure S1) (Korea Ministry of Health and Welfare and Centers for Disease Control and Prevention, 2020). Only 1.4% of COVID-19 cases occurred in acute care hospitals, whereas more than 6% of cases occurred in long-term care facilities and psychiatric long-term care hospitals (Korea Ministry of Health and Welfare and Centers for Disease Control and Prevention, 2020). It is necessary to support and strengthen the infection control systems of long-term care facilities, which may be places of unexpected outbreaks with higher case fatality rates (McMichael et al., 2020).

According to a report from the International Council of Nurses, at least 90 000 HCP worldwide are believed to have been infected with COVID-19 (Nebhay, 2020). HCP comprised about 6% of the 3.5 million cases of COVID-19 in the world, as of May 6, 2020. In South Korea, HCP accounted for 2.4% of the total patients with

COVID-19, significantly lower than the rates in other countries (9.1–29.0%) (Supplementary Material Figure S2) (Korea Ministry of Health and Welfare and Centers for Disease Control and Prevention, 2020; COVID-19 Response Team of Centers for Disease Control and Prevention, 2020; Wang et al., 2020). It was possible to lower the infection rate in HCP by reducing the workload through the efficient distribution of medical resources and providing sufficient personal protective equipment (PPE).

The outbreak of COVID-19 in hospitals will not only lead to missed opportunities of treat chronic diseases in patients with high-risk conditions, but will also lead to high mortality rates if these patients become infected. Even patients with mild symptoms can spread severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) to many people in the enclosed spaces of hospitals. Thus, each hospital should bear in mind the Swiss cheese theory to provide multiple layers of defenses to prevent even small cracks in the hospital's quarantine system (Table 1).

In addition to providing enhanced education and training, most hospitals made it mandatory for hospital staff to wear masks, and only visitors wearing a mask were allowed to enter the hospital. In several hospitals in South Korea, with both the medical staff and patients wearing masks, secondary transmission did not occur even when infected patients stayed in the hospital for a long time (Lee and Jeong, 2020). Both the wearing of face masks and strict hand hygiene are essential for preventing the hospital spread of SARS-CoV-2.

The government has introduced a number of sequential strategies to prevent COVID-19 inflow into hospitals and operate hospitals safely (Figure 1). First, as a quarantine measure for patients suspected of having COVID-19, gate screening

* Corresponding author at: Department of Internal Medicine, Korea University College of Medicine, Guro Hospital, Gurodong-ro 148, Guro-gu, Seoul 08308, Republic of Korea.

E-mail address: infection@korea.ac.kr (J.Y. Song).

Table 1
Safe hospital strategies in South Korea.

Individual strategy	Description
Hospital-level strategies	<ul style="list-style-type: none"> Education and training of hospital staff Universal masking and hand hygiene Inpatient monitoring for new-onset fever and respiratory symptoms
Governmental strategies	<ul style="list-style-type: none"> Hospital gate screening (strengthened triage in the emergency room and at the main hospital entrance) Specialized clinics for patients suspected to have COVID-19 Preemptive isolation of pneumonia patients High throughput diagnostic testing (RT-PCR) Strategies to reduce the hospital workload for COVID-19 Telemedicine

COVID-19, coronavirus disease 2019; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.




Figure 1. Timeline of the COVID-19 pandemic and safe hospital strategies in South Korea. A 'safe hospital' is a healthcare facility equipped with the infrastructure for infection control and prevention that meets the national standards: gate screening and separation of treatment areas between patients with respiratory illnesses and those with non-respiratory illnesses.

(strengthened triage in the emergency room and at the main hospital entrance), specialized clinics, and preemptive isolation of pneumonia patients were conducted from early during the pandemic period (Kim et al., 2020). Two types of specialized clinic have been in operation outside the main hospital building during this COVID-19 pandemic: a clinic for febrile respiratory illness and a clinic for patients with an epidemiological link to the COVID-19 outbreak. Second, the real-time PCR (RT-PCR) test method was validated in collaboration with academia and industry, allowing each hospital to perform a large number of tests and enabling the efficient use of limited isolation rooms. Third, considering the possibility of cross-transmission when cases

are concentrated in large hospitals all at one time, the inspection function of the public health center was strengthened and a residential treatment center was introduced to manage patients with mild COVID-19. In late February 2020 in South Korea, more than 500 cases of COVID-19 were reported daily, which was expected to reach unacceptable levels for the hospitals (Korea Ministry of Health and Welfare and Centers for Disease Control and Prevention, 2020). Thus, the first residential treatment center was opened on March 2, with further expansion to a total of 18 centers, which could accommodate 3500–4000 patients. As a result, hospitals could focus on the treatment of patients with severe COVID-19. Finally, telemedicine was introduced so that patients in

COVID-19 outbreak areas could be treated without face-to-face contact with HCP. These strategies should be implemented in an integrated manner and shared with hospital staff and visiting patients.

Funding source

No funding was provided for this manuscript.

Ethical approval

Ethical approval was not required.

Conflict of interest

We declare no conflict of interest.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.ijid.2020.06.094>.

References

- Choi WS, Kang CI, Kim Y, Choi JP, Joh JS, Shin HS, et al. Clinical presentation and outcomes of Middle East Respiratory Syndrome in the Republic of Korea. *Infect Chemother* 2016;48(2):118–26, doi:<http://dx.doi.org/10.3947/ic.2016.48.2.118>.
- Kim YJ, Jeong YJ, Kim SH, Kim YJ, Lee SY, Kim TY, et al. Preparedness for COVID-19 infection prevention in Korea: a single-centre experience. *J Hosp Infect* 2020; 105(2):370–2, doi:<http://dx.doi.org/10.1016/j.jhin.2020.04.018>.
- Korea Ministry of Health and Welfare and Centers for Disease Control and Prevention. Updates on coronavirus disease 2019 (COVID-19); for press release. 2020.
- Lee JK, Jeong HW. Wearing face masks regardless of symptoms is crucial for preventing spread of COVID-19 in hospitals. *Infect Control Hosp Epidemiol* 2020;1–4, doi:<http://dx.doi.org/10.1017/ice.2020.202>.
- McMichael TM, Currie DW, Clark S, Pogojans S, Kay M, Schwartz NG, et al. Epidemiology of Covid-19 in a long-term care facility in King County, Washington. *N Engl J Med* 2020;382(21) 2005–201.
- Nebehay S. Over 90,000 health workers infected with COVID-19 worldwide — nurses group. Thomson Reuters; 2020.
- CDC COVID-19 Response Team. Characteristics of health care personnel with COVID-19—United States, February 12–April 9, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69(15):477–81.
- Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020;323(11):1061–9, doi:<http://dx.doi.org/10.1001/jama.2020.1585>.