

Journal Pre-proof

Therapy of primary liver cancer

Mei Feng, Yisheng Pan, Ruirui Kong, Shaokun Shu



PII: S2666-6758(20)30032-1

DOI: <https://doi.org/10.1016/j.xinn.2020.100032>

Reference: XINN 100032

To appear in: *The Innovation*

Please cite this article as: Feng, M., Pan, Y., Kong, R., Shu, S., Therapy of primary liver cancer, *The Innovation* (2020), doi: <https://doi.org/10.1016/j.xinn.2020.100032>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020

Hepatocellular carcinoma (80%)


Risk factors:
HBV, HCV, fatty liver disease,
diabetes, alcohol consumption,
aflatoxin exposure

Clinical pathology:
well differentiated HCC,
moderately differentiated HCC,
poorly differentiated HCC

Genetic alterations:
TP53, AXIN1, RB1, MYC, MET,
TERT, CTNNB1, ARID1A, ARID2,
BAP1, MLL, EZH2, CCND1, CDKN2A

Targeted therapy:
VEGFR, CDK4/6,
TGF-beta inhibitors

Immunotherapy:
PD-1, PD-L1, CTLA-4
CAR-T, DC Vaccines



Intrahepatic cholangiocarcinoma (15%)



Risk factors:
Choledochal cyst, cholelithiasis,
choledocholithiasis, liver flukes,
viral hepatitis

Clinical pathology:
MF, PDI, IDG

Genetic alterations:
FGFR2 fusion, IDH1, IDH2,
KRAS, BRAF

Chemotherapy
CisGem, FOLFOX

Targeted therapy
FGFR, IDH1/2, VEGFR inhibitors
Immunotherapy
PD-1, PD-L1, CTLA-4




Cancer cell

Immune cell

Combined

Mixed



Combined HCC-ICC (2-5%)

Risk factors:
Overweight, obese, nonalcoholic steatohepatitis, liver cirrhosis

Clinical pathology:
Separate, Combined, Mixed

Genetic alterations:
TP53, TERT promoter, AXIN1

Therapy:
Combined—ICC like,
Mixed—HCC like

Therapy of primary liver cancer

Mei Feng^{1,4}, Yisheng Pan⁴, Ruirui Kong^{1,*}, Shaokun Shu^{1,2,3,*}

¹Translational Cancer Research Center, Peking University First Hospital, Beijing 100034, China; ²Department of Biomedical Engineering, Peking University, Beijing, 100871, China; ³Peking University Cancer Hospital, Beijing, 100142, China; ⁴Department of General Surgery, Peking University First Hospital, Beijing 100034, China.

*Correspondence to: Shaokun Shu and Ruirui Kong

E-mail: shaokun_shu@bjmu.edu.cn and ruiruikong@pkufh.com

Abstract

Primary liver cancer (PLC) is a fatal disease that affects millions of lives worldwide. PLC is the leading cause of cancer-related deaths and the rate of incidence is predicted to rise in the coming decades. PLC can be categorized into three major histological subtypes: hepatocellular carcinoma (HCC), intrahepatic cholangiocarcinoma (ICC), and combined HCC-ICC (cHCC-ICC). These subtypes are distinct with respect to epidemiology, clinicopathological features, genetic alterations, and clinical managements, which are thoroughly summarized in this review. The state of treatment strategies for each subtype, including the currently approved drugs and the potential novel therapies, are also discussed.

Key words: primary liver cancer, hepatocellular carcinoma, intrahepatic cholangiocarcinoma, combined HCC-ICC, PLC therapy

Introduction

Primary liver cancer (PLC) is a deadly malignancy with significant histological and biological heterogeneity, and ranks as the fourth leading cause of cancer-related death worldwide.^{1,2} Therefore, it has become a major public healthy challenge. Over the past decades, the morbidity and mortality associated with PLC have steadily risen.

According to Globocan's latest Global Cancer Statistics Report, 841,080 cases of liver cancer were reported worldwide in 2018, accounting for 4.7% of the total cancer cases in the same period, while deaths totaled 781,631, accounting for 8.2% of total cancer deaths.³ On the basis of annual projections, the World Health Organization estimates that 1,276,679 patients will die from liver cancer in 2040. Incidence and mortality of PLC differ widely between regions. The highest incidence of PLC was observed in East Asia and in sub-Saharan Africa.⁴ In particular, China experiences the highest number of cases of PLC, with a high incidence rate (18.3 cases/100,000 inhabitants).⁵

PLC manifests as three subtypes: HCC, ICC, and cHCC-ICC, which differ notably in epidemiology, clinicopathological morphology, genetic alteration and appropriate therapeutic responses. HCCs are primarily related to viral infection, alcohol abuse, and metabolic syndrome⁶, whereas ICCs are mainly associated with liver chronic inflammation and biliary tract diseases.^{7,8} Risk factors for development of cHCC-ICC include overweight, obesity, nonalcoholic steatohepatitis and liver cirrhosis.^{9,10} HCCs show a solid and trabecular pattern with local invasion restricted to the liver¹¹⁻¹³, whereas ICCs are ductular, papillary or solid tumor structures with high metastasis to distal organs¹⁴⁻¹⁶. cHCC-ICCs are the combination of the HCC and ICC phenotypes present in liver tissue, and are classified into separate, combined, and mixed cHCC-ICC subclasses, which are more aggressive and have a poorer prognosis.^{2,17-21}

The three PLC subtypes have distinct genetic alterations and molecular patterns. HCCs are associated with genetic alterations in specific chromosomal regions and genes, including TERT promoter mutation, TP53 deletion, and WNT signaling (CTNNB1 and AXIN1) activation.²²⁻²⁹ ICCs show a unique mutational landscape with recurrent mutations, with the genetic alterations in TP53, KRAS, IDH1/2 and FGFR gene fusions.³⁰⁻³⁵ Combined cHCC-ICCs showed strong ICC-like features, whereas mixed cHCC-ICCs showed HCC-like features.^{36,37} Understanding the molecular alterations that initiate various PLCs subtypes is of great importance for us to determine the mechanisms of tumorigenesis. Genetic alterations can be transformed into biomarkers that may represent new therapeutic targets, affect the treatment decisions, and ultimately improve the treatment of liver cancer patients. HCCs mainly respond to

targeted therapy, immunotherapy, and antiviral agents, while ICC patients benefit from classical chemotherapy, targeted therapy, and immunotherapy. Based on the pathological classification and the molecular features of cHCC-ICCs, combined cHCC-ICCs should be treated with the therapies resembled to ICCs, whereas mixed cHCC-ICCs are treated more like HCCs. In this review, we systematically summarize the epidemiology, pathogenesis, genetic alteration, and treatment for each subtype and comprehensively describe current therapy drugs and the potential novel therapies for PLC.

1. Epidemiology and risk factors

• HCC

HCC represents the major histologic subtype, accounting for approximately 80% of all cases of primary liver cancer. The risk factors for HCC includes hepatitis B/C viral infection (HBV and HCV), aflatoxin B1, alcoholic abuse, and non-alcoholic, metabolic symptoms, such as diabetes and obesity.⁶ According to the Global Burden of Disease from 1990 to 2015, HBV and HCV accounted for 432, 000 liver cancer deaths (54%), alcohol for 245, 000 (30%), and other causes for 133, 000 (16%) deaths. In particular, 55% of all HCC cases worldwide are reported from China³⁸ due to the locally high prevalence of hepatitis B virus (HBV) infection.

• ICC

As the second most common liver carcinoma following HCC, ICC accounts for around 15% of PLC cases with a high incidence of 2 per 100,000 population worldwide annually.³⁹ The most common risk factors for ICC are biliary tract diseases including choledochal cysts, cholelithiasis, choledocholithiasis, liver flukes, viral hepatitis, metabolic syndrome, and other risk factors including tobacco and alcohol use, and cirrhosis.⁷ Recently, the incidence of ICC has been increasing more rapidly owing to risk factors⁸ including increasing chronic liver disease and environmental toxins, and is found more often due to improved diagnostic tools and imaging.

• cHCC-ICC

cHCC-ICC presents as a heterogeneous tumor showing both hepatocyte and cholangiocyte differentiation, and has a poor prognosis.⁴⁰ cHCC-ICC is a rather rare tumor with an incidence rate less than 5%.¹ The poor prognosis associated with cHCC-ICC is due to the limited treatment options and difficulty of diagnosis. To date, the largest cohort analysis which included 529 patients diagnosed with cHCC-ICC between 2004 and 2014 across 18 registries⁴¹ reported that the incidence of cHCC-ICC in men and women was 0.08 and 0.03 per 100,000 per year respectively, with the average age of 63 y at diagnosis. One- and five-year cause-specific survival rate for cHCC-ICC was 41.9% and 17.7%, respectively, with the median survival of 8 m. Among racial groups, cHCC-ICCs are most common in Asian and Pacific islanders. Obesity, nonalcoholic steatohepatitis and liver cirrhosis were observed in some cHCC-ICC cohorts^{9,10} and are potential risk factors for cHCC-ICC.

2. Clinicopathological features

- **HCC**

HCC shows a solid, trabecular and pseudoglandular pattern with a high density of tumor cells. It has three subtypes: well differentiated HCC, moderately differentiated HCC, and poorly differentiated HCC.¹¹⁻¹³ Well-differentiated HCCs are often small (less than 2 cm in diameter) and are composed of cells with a higher nuclear to cytoplasmic ratio, arranged in a thin trabecular pattern with rare pseudoglandular structures. Moderately differentiated HCCs are usually larger tumors (larger than 3 cm) showing polygonal tumor cells in a thick trabecular arrangement with a frequent pseudoglandular pattern. Poorly differentiated HCCs are composed of pleomorphic tumor cells in a solid or compact growth pattern.

- **ICC**

ICC can be divided into two subtypes: a small duct type which originates from small intrahepatic ductules with no or minimal mucin production, and a large bile duct type which arises from large intrahepatic ducts proximal to the bifurcation of the right and left hepatic ducts, with high mucin production ability.¹⁴⁻¹⁶ Further, ICC shows three different growth patterns: mass-forming (MF), periductal infiltrating (PI), and

intraductal growth (IG).⁴² MF ICC is a firm, multilobulated, unencapsulated, white-gray tumor, owing to its extensive desmoplastic stroma. The PI subtype shows extensive infiltration along the intrahepatic hilum structure, and the IG subtype is usually restricted to tubes with papillary structures. MF ICC is the most common type associated with a poor prognosis while IG type is rare but has a favorable prognosis.¹⁷

- **cHCC-ICC**

Though the phenomenon of HCC and ICC being present in the same liver was first described in 1903.¹⁷ cHCC-ICC was not systematically described until 1949, when it was classified into three subtypes depending on the location of HCC and ICC: type A (separate type) has separate nodules of hepatocellular and bile duct carcinoma; type B (combined type) shows contiguity with intermingling but with clearly defined areas; type C (mixed type) presents as intimate association without clear boundaries.¹⁸ In 1985,

another classification system with three subtypes was established: Type I (collision tumors) — simultaneous occurrence of both HCC and ICC in the same patient; Type II (transitional tumors) — with an identifiable intermediate transition between HCC and ICC; Type III (fibrolamellar tumors) — which resembled the fibrolamellar variant of HCC but also contained mucin-producing pseudoglands.¹⁹ Presently, the WHO 2010 classification is commonly used, in which cHCC-ICC is classified into two main types, the classic type and the SC type (subtypes with stem cell (SC) features), with the SC type subdivided into three subtypes including the typical subtype (TS), intermediate subtype (INT), and cholangiolocellular type (CLC)⁴³.

The lack of a unified classification system greatly adds to the difficulty for cHCC-ICC research and the clinicopathological characteristics of cHCC-ICC remain ill-defined. cHCC-ICC can exhibit stem/progenitor cell phenotypes consisting of small cells with scant cytoplasm, hyperchromatic nuclei embedded within a thick, desmoplastic stroma, a high nuclear/cytoplasmic ratio, and the increased mitotic activity.¹ And the IHC (immunohistochemistry) identified stemness-related markers (KRT19, CD56, EpCAM, CD117, CD113, OV6).^{1,20} cHCC-ICC clinicopathologic characteristics include more frequent multifocal lesions, more microvascular emboli, and portal vein and lymph node invasion, all of which indicate a poor prognosis.²¹

3. Genetic alterations

- **HCC**

Wide-scale genomic studies have revealed that hundreds of somatic DNA alterations accrue in HCC, including chromosome aberrations and mutations. High-level DNA amplifications are enriched in chromosome locations 6p21 and 11q13 location in HCC⁴⁴, which occur in 5-10% of cases. Recently, some oncogenic genes were identified in the regions of frequent DNA gain. For example, LINC01138 is an oncogenic long intergenic non-coding RNA located in this region which has been identified as a driver of HCC.⁴⁵ VEGFA and CCND1/FGF19 have also identified in these regions and are potential therapeutic targets.⁴⁶ Loss of heterozygosity on chromosome 8p is a frequent event in HCC.⁴⁷ These DNA alterations are often associated with cancer progression due to the deletion of tumor suppressor genes. Intriguingly, in these regions, a variety of vulnerability genes have been recently identified. For example, TSLNC8 was characterized as a tumor suppressor gene on chromosome 8p12, the region that shows allelic loss in HCC and was shown to inhibit the proliferation and metastasis of HCC.⁴⁸ The genetic mutations of HCC have been well-studied. Mutations in the TERT promoter occur in approximately 60% of cases and cause recurrent viral insertion of HBV.⁴⁹ Deletion mutations in TP53 are the most frequent genetic alterations, accounting for about 30% of cases²²⁻²⁹, and are thought to be the initiating event driving the formation of precursor lesions. Mutated genes in WNT signaling (CTNNB1 and AXIN1) and chromatin remodeling (ARID1A) account for approximately 27-40% of cases.²²⁻²⁹ Accumulation of activating mutations in oncogenes, including activation of AKT or mTOR and of the oxidative stress pathway activation, occurs throughout tumor progression, and could be potentially targeted with molecular therapies in the future.

- **ICC**

ICC shows a unique mutational landscape with recurrent mutations, compared with HCC. It harbors the genetic alterations in TP53, KRAS, ARID1A, BAP1, IDH1, IDH2, PIK3CA, SMARCB1, EPHA2, SMAD4, GNAS and PBRM1 as well as FGFR gene fusions.³⁰⁻³⁵ Gain-of-function of IDH1 and IDH2 mutation on R132 and R172 two hotpot codons was observed in 10-28% of ICC cases.³² Fusions, amplifications, translocations and rearrangements of FGFR genes are found in ICC and are closely related to the initiation and progression of ICC.⁵⁰ The activating mutation of KRAS (15-20%) is another most frequently genomic alteration in ICC.^{31,51,52} The KRAS

mutation often exists concurrently with FGFR2 fusions and IDH mutations, suggesting a possible cooperative role in ICC pathogenesis.^{53,54} In addition, recent studies have shown that BRAF and Notch are considerably more prevalent in ICC and function in ICC pathogenesis.⁵⁵

- **cHCC-ICC**

cHCC-ICC are genetically complex tumors. The combined subtype of cHCC-ICC shows strong ICC-like features, with the high expression of EPCAM, KRT19, PRDM5 and KRAS. The mixed subtype of cHCC-ICC shows HCC-like features with the high expression levels of AFP, GPC3, APOE, SALL4 and AFP81.³⁶

The most frequent mutation observed in cHCC-ICCs is TP53 with a strikingly high 49.2% mutation frequency, much higher than that in HCC (20%-35%) and ICC (18%-38%).^{36,56} Interestingly, several studies have found that the disruption of Trp53 alone in livers of mice can induce the formation of cHCC-ICC,^{37,57} which further implies that TP53 may be the driver gene in cHCC-ICC. It is notable that Nestin - a type VI intermediate filament (IF) protein which is commonly used as a neuroectodermal stem cell marker, is highly expressed in cHCC-ICC and is strongly associated with poorer prognosis.³⁶ Hence, Nestin may be a promising biomarker for cHCC-ICC.

4. Challenges and limitations of current treatment strategies

4a. Resection, transplantation, local and regional therapies

- **HCC**

The commonly used staging system for HCC is the Barcelona Clinic Liver Cancer staging system (Figure 1). HCC in the very early stage or intermediate stage can be treated with the local regional therapies, which includes radiofrequency ablation (RFA), resection (Da Vinci surgery, laparoscopic surgery or traditional surgery), transplantation (orthotopic liver transplantation, piggyback transplantation, split liver transplantation, auxiliary liver transplantation), percutaneous ethanol injections (PEI), or transcatheter arterial chemoembolization (TACE), etc.⁵⁸

- **ICC**

Surgery is currently the only curative cure for ICCs but only a minority of patients in early stages are considered candidates for resection. In surgery, ICC is usually treated with hepatic resection to achieve negative resection margins.⁵⁹ For patients with locally unresectable ICC, tumor ablation such as RFA, or hepatic artery-based therapies like yttrium-90 radioembolization, appear promising.⁵⁹⁻⁶⁴

- **cHCC-ICC**

An accurate diagnosis is of paramount importance for the treatment of cHCC-ICC. Currently, major hepatectomy is the optimal management for cHCC-ICC.⁶⁵ The rarity of this cancer as well as the lack of biomarkers have made this cancer difficult to diagnosis and manage. Surgical resection remains the only curative option for patients with cHCC-ICC.

The treatment options for cHCC-ICC are similar to those for HCC and ICC and include surgery, radiation, yttrium-90 radioembolization, chemotherapy, combined radiation and chemotherapy, combined surgery and chemotherapy, and triple therapy (surgery, radiation, and chemotherapy).^{41,66-69} A recently retrospective analysis from 2001 to 2015 of 623 PLC patients including 47 cHCC-ICC, 468 HCC, and 108 ICC patients who underwent resection found that although cHCC-ICC is more poorly differentiated than HCC and ICC, it had a similar five year survival rate (49.7%, 54.8% and 68.7%, respectively) and three year recurrence rate (57.9%, 61.5%, 56%, respectively).⁷⁰

4b. Systemic chemotherapy

- **HCC**

Systemic chemotherapy has limited efficacy on HCC: several clinical trials of chemotherapy has showed low response rate and worse toxicity without an significant improvement in the overall survival (OS), including Gemcitabine - and Doxorubicin-based treatment, FOLFOX (5-Fluorouracil, Leucovorin, Oxaliplatin) and PIAF (Cisplatin / Interferon alpha-2b / Doxorubicin / Fluorouracil).⁷¹⁻⁷⁴ This suggests a limited role for traditional chemotherapy in the treatment of advanced HCC.

- **ICC**

Current first-line standard of treatment for ICC is the combination of Gemcitabine and Platinum-derived chemotherapy (Figure 2B). With the poor prognosis, the median survival of advanced ICC patients is less than one year. Very limited effective treatments are available for patients who progress on first-line chemotherapy, so there is a high medical demand.

The adoption of combination of Gemcitabine and Platinum-derived chemotherapy have currently been the first line standard-of-care for patients with ICC (Figure 2B). Yet, the vast majority of patients are diagnosed with advanced disease, facing the median survival of less than one year. ICC therapy for patients who progress on first-line chemotherapy is of limited efficacy, thus, the novel and effective therapies are required.

First line treatment

Effective molecular targeted therapy and immunotherapy is lacking, so chemotherapy, with Gemcitabine, Platinum compounds and Fluoropyrimidines, is still the mainstream of standard treatment for unresectable ICC.

The most primary chemotherapy for ICC is Gemcitabine which was established as the first-line therapy for advanced biliary tract cancer in 1999. In 2010, the randomized, controlled, ABC-02 phase III clinical trial compared the benefit of Gemcitabine plus Cisplatin (CisGem) chemotherapy with the single agent Gemcitabine.⁷⁵ This study showed an advantage for CisGem in OS (11.7m vs 8.1m; HR: 0.64; 95% CI: 0.52-0.80) and progression-free survival (PFS) (8.0m vs 5.0m, $p < 0.001$). This effectiveness was confirmed in a Japanese randomized phase II study, BT22 (median OS 11.2m vs 7.7m; HR: 0.69).⁷⁶ Based on these promising results, CisGem is currently regarded as the standard of care in the first-line treatment for advanced CCA.

CisGem has been shown a survival advantage and is currently proposed as the standard practice of first-line treatment for advanced cholangiocarcinoma (CCA).

Other than Cisplatin, Gemcitabine plus other agents such as Oxaliplatin, S-1, Capecitabine, Bevacizumab, and Nab-paclitaxel have also been considered as the

first-line choices for advanced CCA based on the promising outcomes from several phase II or III trials.⁷⁷⁻⁹¹ A recent multicenter, randomized, phase III clinical trial (NCT01470443) results showed that Capecitabine plus Oxaliplatin (XELOX) was noninferior to Gemcitabine plus Oxaliplatin (GEMOX) in aspect of 6-month PFS rate (46.7% vs 44.5). No significant difference were seen in this two treatment groups in terms of tumor response, OS, and safety. Also, lower frequencies of hospital visits occurred in the XELOX group. Thus, XELOX could be an alternative first-line therapy for CCAs.⁹⁰

A recent multicenter, randomized, phase III clinical trial (NCT01470443) results showed that Capecitabine plus Oxaliplatin (XELOX) has the comparable efficacious effect to Gemcitabine plus Oxaliplatin (GEMOX) in terms of tumor response, survival rate (OS and PFS) and safety. Also, XELOX has an advantage of low hospital visits, compared to GEMOX. Thus, XELOX could be optional and alternative for CCA therapies.

Second-line treatment

There is no established standard second-line chemotherapy for advanced CCA, and all regimens have shown limited efficacy, with a median PFS of around 3 month and median OS about 7 months.⁹²

FOLFOX (L-folinic acid, 5 FU, and Oxaliplatin) is an optional second-line treatment option based on the randomized phase III, multi-center, open-label ABC-06 study (NCT01926236). FOLFOX showed increased benefit for the Median OS (months (m)), 6m and 12m and OS-rate (%): 6.2m, 50.6% and 25.9% compared to 5.3m, 35.5%, 11.4% for the control group (ASC arm).⁹²

Based on the randomized phase III, multi-center, open-label ABC-06 study (NCT01926236), FOLFOX (L-folinic acid, 5 FU, and Oxaliplatin) showed better benefit with the Median OS (months (m)), 6m and 12m and OS-rate (%): 6.2m, 50.6% and 25.9% compared to 5.3m, 35.5%, 11.4% for the control group (ASC arm).⁹² Currently, FOLFOX has regarded as the second-line treatment option.

Currently several phase II and III chemotherapy clinical trials are under way (Table 3). Combined therapy with chemotherapy shows promise in the treatment of CCA: elective internal radiotherapy (SIRT) plus chemotherapy or hepatic arterial infusion plus systemic chemotherapy both had antitumor activity and are promising for the treatment of ICC.^{93,94}

- **cHCC-ICC**

In contrast to surgery-based treatments for resectable cHCC-ICC, systemic therapy is the nonstandard options for advanced and unresectable cHCC-ICC, based on the standard treatment strategy for the unresectable HCC or ICC. Chemotherapy for advanced or unresectable cHCC-ICC is largely understudied, with only a few case reports and some retrospective studies have been published.^{9,10,95-101} Recently, a multicenter retrospective analysis has been conducted by Kobayashi, S. and his colleagues.¹⁰ They enrolled 36 patients and divided them into 4 groups treating (1) Gemcitabine plus Cisplatin (n=12); (2) Fluorouracil plus Cisplatin (n=11); (3) Sorafenib monotherapy (n = 5); (4) others group (n = 8). Platinum-containing reagents are proven more effective than Sorafenib monotherapy with the OS being 11.9 (95% CI: 4.9-18.8), 10.2 (95% CI: 3.9-16.6), 3.5 (95% CI: 0.0-7.6) and 8.1 (95% CI: 0.9-15.4) months, respectively.

According to divided-group treatment with (1) Gemcitabine plus Cisplatin (n=12); (2) Fluorouracil plus Cisplatin (n=11); (3) Sorafenib monotherapy (n = 5); (4) others (n = 8), they found that 36 patients with Platinum-containing treatment have longer overall survival time than those treated by sorafenib monotherapy, showing OS with 11.9 (95% CI: 4.9-18.8), 10.2 (95% CI: 3.9-16.6), 3.5 (95% CI: 0.0-7.6) and 8.1 (95% CI: 0.9-15.4) months, respectively.

A similar conclusion was drawn in another retrospective study of 123 cHCC-ICC patients, with 68 receiving Gemcitabine-based therapy (Gemcitabine + Platinum or Gemcitabine + 5-FU) or targeted agents (Sorafenib).⁹ Median PFS favored

Gemcitabine/Platinum and Gemcitabine/5-FU (8.0 and 6.6 months respectively) over Sorafenib monotherapy (4.8 months).

4c. Molecular targeted therapy

- **HCC**

First line drugs:

1. Sorafenib

Sorafenib was the first U.S. Food and Drug Administration (FDA) approved first-line systemic targeted drug for advanced HCC. It is an oral small-molecule multikinase inhibitor targeting VEGFR1, VEGFR2, VEGFR3, PDGFR β and the Raf. Two large, international multicenter clinical trials, SHARP and Asian-Pacific, have proven that Sorafenib can suppress tumor progression and prolong OS in patients with advanced HCC.^{102,103}

Sorafenib was first approved FDA (U.S. Food and Drug Administration) as the first-line targeted drug for advanced HCC. Sorafenib is an oral small-molecule multi-kinase inhibitor, which targets VEGFR1/2/3, PDGFR β and the Raf. Two large, international multicenter clinical trials, SHARP and Asian-Pacific, showed that Sorafenib has approved to increase ~3 months of progression-free and overall survival in patients with advanced HCC in western countries. As the first generation of targeted drugs for HCC, Sorafenib has been used for over a decade. During this time, many patients have benefited, though others quickly developed resistance to Sorafenib.¹⁰⁴

2. Lenvatinib

Lenvatinib is becoming an available option for HCC patients who develop Sorafenib resistance. Lenvatinib is an oral TKI, inhibiting VEGFR1-3, FGFR1-4, PDGFR, RET, and KIT. In August 2018, the Food and Drug Administration approved Lenvatinib for first-line treatment of patients with unresectable HCC after Lenvatinib was proven to be noninferior to Sorafenib in the phase 3 REFLECT trial.¹⁰⁵

In August 2018, Lenvatinib was approved by FDA for the first-line treatment of patients with unresectable HCC, based on its noninferior effect to Sorafenib in the phase III REFLECT trial.

Median overall survival in the Lenvatinib arm and Sorafenib arm was 13.6 months and 12.3 months (HR: 0.92; 95% CI: 0.79, 1.06), respectively. The adverse effects were hypertension (42%), diarrhoea (39%), decreased appetite (34%) with lenvatinib, and palmar-plantar erythrodysesthesia (52%), diarrhoea (46%), decreased weight (31%), hypertension (30%), decreased appetite (27%) with sorafenib.

3. Donafenib

Similar to Sorafenib, Donafenib is a novel multikinase inhibitor targeting RAF kinase and various receptor tyrosine kinases (RTKs) including VEGFR, BRAF.¹⁰⁶ According to the report from 2020 International Conference of the American Society of Clinical Oncology (ASCO), Donafenib significantly improves OS over Sorafenib (12.1 vs 10.3 months) with fewer side effects and higher patient tolerance for advanced HCC patients in its phase II/III open-label trial.¹⁰⁷ The grade 3 and above adverse reaction rates for Donafenib and Sorafenib were 57.4% and 67.5%, respectively. Thus, Donafenib was recommended as the first-line therapy in the ASCO guidelines for HCC.

Second line drugs:

1. Regorafenib

Regorafenib, as an oral multi-kinase inhibitor, inhibits the activity of protein kinases involved in multiple biological processes, such as tumorigenesis, tumor angiogenesis, distant metastasis and tumor immune escape. These kinases include VEGFR 1-3, TIE2, RAF1, KIT, RET, RAF, BRAF, PDGFR, FGFR, and CSF1R. The randomized, double-blind, multicenter, phase III clinical trial RESORCE study showed that regorafenib significantly improves the overall survival of patients, as compared with the placebo, from 7.8 to 10.6 months (HR: 0.63, $P < 0.0001$).¹⁰⁸ Grade 3-4 adverse events were reported in 40% of patients receiving the regorafenib and 11% of patients receiving the placebo. In 2017, regorafenib received FDA approval as the second-line

drug for the treatment of patients with advanced HCC who fail to respond to the Sorafenib treatment.

2. Cabozantinib

Cabozantinib is an oral inhibitor and targets multiple kinases, including VEGFR2, cMET, RET, ROS1, TYRO3, MER, KIT, TRKB, FLT3, TIE-2, as well as the GAS6 receptor (AXL)^{109,110}. It was originally approved for medullary thyroid cancer in 2012 and advanced renal carcinoma in 2016. According to the randomized, double-blind multicenter phase 2 clinical trial conducted across 95 centers in 19 countries, median OS was 10.2 months for patients receiving Cabozantinib, and 8 months for patients treated with placebo (HR = 0.76, P=0.005).¹¹¹ Median PFS was 5.2 months and 1.9 months, respectively. Grade 3 or 4 adverse events occurred in 68% of patients in the Cabozantinib arm and 36% in the placebo arm. The observed hepatotoxicity can be mostly controlled through dose modifications. Based on the encouraging results of prolonged OS and PFS, Cabozantinib received its FDA approval for HCC in 2018.

Initially, Cabozantinib was approved to treat medullary thyroid cancer in 2012 and advanced renal carcinoma in 2016. According to the randomized, double-blind multicenter phase II clinical trial conducted across 95 centers in 19 countries, median OS and PFS were respectively 10.2 months and 5.2 months for patients receiving the Cabozantinib, whereas for patients treated with the placebo, median OS and PFS were only 8 months and 1.9 months.

3. Ramucirumab

Ramucirumab is a completely human monoclonal antibody, which can specifically inhibit VEGFR-2.¹¹² For patients with alpha-fetoprotein ≥ 400 ng/ml and have been previously treated with Sorafenib, Ramucirumab was approved as a monotherapy by the FDA on May 10, 2019.

Ramucirumab was approved as a monotherapy by the FDA on May 10, 2019, for patients with high level of alpha-fetoprotein and patients who were previously treated with Sorafenib.

Approval was based on REACH 2 (NCT02435433), a randomized, double-blind, multicenter phase III study of 292 patients with AFP \geq 400 ng/mL who had disease progression after Sorafenib or were intolerant to Sorafenib.¹¹³ More recently, a study further confirmed the efficacy of Ramucirumab in elderly patients with HCC and elevated AFP after Sorafenib in REACH and REACH-2 with a survival benefit observed across all age subgroups and a tolerable safety profile, supporting its value irrespective of age, including for patients \geq 75 years.¹¹⁴

4. Apatinib

Apatinib, a tyrosine kinase inhibitor targeting VEGFR-2, significantly prolonged OS and PFS in Chinese patients with advanced HCC who had previously been treated with Sorafenib and/or chemotherapy, according to the results of a randomized, placebo-controlled, phase III trial conducted in 31 sites in China.¹¹⁵ Median OS was almost 2 months longer for patients who received Apatinib compared with patients receiving the placebo (8.7m vs 6.8m), and median PFS was more than 2 months longer (4.5m vs 1.9m).¹¹⁵ The most common grade 3 or worse adverse events occurred at a rate of 69.2% in the Apatinib arm and 3.1% in the placebo arm. With the significantly prolonged OS and PFS and a manageable safety profile, Apatinib has potential to become a new second-line therapy for liver cancer.

5. Novel therapeutic targets

Even with all these available treatments (Table1), the median PFS for HCC patients remains less than a year. Thus, novel treatment is still a critical unmet need for treatment of HCC. Based on the genomic profile and biomarkers reported in HCC, several clinical trials targeting various pathways are currently ongoing (Table 2). Recently, a first-in-human phase I study (NCT02508467) of Fisogatinib (BLU-554) – an orally bioavailable inhibitor of human FGFR4 demonstrated its anti-tumor activity in HCC, and future validated that the aberrant FGF19–FGFR4 signaling pathway may

be a driver event.¹¹⁶ In addition, the TGF- β 1 Receptor Type I Inhibitor Galunisertib also showed an acceptable safety and prolonged OS outcome in combination with Sorafenib in a phase II trial (NCT01246986).^{117,118} Other potential candidates including the cyclin-dependent kinases (CDKs) inhibitors regulating the cell cycle pathways - Ribociclib, Palbociclib,^{119,120} Abemaciclib and Milciclib as well as the c-MET inhibitors Tepotinib¹²¹ and Tivantinib¹²² are being evaluated in HCC clinical trials.

- **ICC**

Molecular targeted therapy controls tumor cell proliferation, apoptosis, adhesion and movement by inhibiting the surface molecules of tumor cell membranes and thereby inhibiting intracellular signaling pathways. ICC genetic alterations primarily include fibroblast growth factor receptor (FGFR), isocitrate dehydrogenase (IDH), epidermal growth factor (EGFR), and breast cancer type 1 susceptible protein associated protein-1 (BAP1).¹²³⁻¹²⁵ Genetic alterations of these genes all have implications for therapy. At present, a variety of molecular targeted drugs are in the clinical research stage (Table 3), and some of which have made progress in the treatment of ICC (Table 1).

1. FGFR inhibitors

The most promising target therapy for CCA identified in recent years is the inhibitor of the fibroblast growth factor (FGF) signaling pathway, which consists 22 members labeled FGF1-23 (FGF15 = FGF19, called FGF15/19) and four interacting transmembrane receptors (FGFR1-4).¹²⁶ Fibroblast growth factor signals regulates cell proliferation, in which FGFR2 fusions occurred in 10–20% of ICC patients and are considered as a promising therapeutic target.^{33,51,127,128} Currently, several FGFR inhibitors are being evaluated in clinical trials for CCAs with FGFR genetic aberrations.

Pemigatinib (INCB054828)

Pemigatinib is the first and only targeted therapy so far approved (in 2020) by the FDA for the treatment of this rare cancer. It is a selective, potent oral inhibitor of fibroblast

growth factor receptor (FGFR) 1, 2, and 3.¹²⁹ Approval was based on findings from the phase II FIGHT-202 trial (NCT02924376), which enrolled 107 patients with locally advanced or metastatic cholangiocarcinoma with FGFR2 fusions or rearrangements (Cohort A), other FGF/FGFR genetic alterations (Cohort B), or no FGF/FGFR genetic alterations (Cohort C). For those in cohort A, treatment with Pemigatinib resulted in a median OS of 21.1 months and median PFS 6.9 months. The FIGHT-202 study suggests that locally advanced or metastatic cholangiocarcinoma patients with fibroblast growth factor receptor 2 (FGFR2) fusions or rearrangements may benefit from a potent oral FGFR1, 2, and 3 inhibitor treatment. Median progression-free survival was 6.9 months for patients with FGFR2 alterations, 2.1 months for patients with other FGF/FGFR alterations and 1.7 months for those with no alterations in these genes. Median overall survival was 21.1m, 6.7m and 4.0m for the respective cohorts.¹³⁰ With the promising results of phase II, the phase III clinical trial of Pemigatinib is currently underway (NCT03656536).

Infigratinib (BGJ-398)

Infigratinib (BGJ-398) was the first FGFR inhibitor investigated for treatment of CCA. It is an oral drug which selectively binds to FGFR 2 and shows impressive anti-tumor efficiency and a manageable safety profile in participants with advanced FGFR-altered CCA (NCT02150967).¹³¹ The FDA granted fast track designation to Infigratinib early in 2020 for first-line treatment of patients with unresectable advanced or metastatic CCA who harbor FGFR2 gene fusions or translocations. It is currently undergoing a phase III trial (NCT03773302) to assess the efficacy and safety of Infigratinib versus standard treatment chemotherapy CisGem (Gemcitabine plus Cisplatin) in first-line treatment of CCA patients. Patients will be randomized 2:1 to receive Infigratinib or CisGem.

Futibatinib (TAS-120)

Futibatinib (TAS-120) is a highly potent and selective irreversible pan-FGFR inhibitor for all four FGFR subtypes (FGFR1-4).¹³² Futibatinib demonstrated a clinically

promising benefit with a manageable toxicity profile in patients with cholangiocarcinoma harboring FGFR2 gene fusions in phase I/II (NCT02052778).^{133,134} Furthermore, Futibatinib can overcome acquired resistance to the ATP-competitive FGFR Inhibitors BGJ398 and Debio 1347 and still show promise to patients who had previously progressed on FGFR inhibitors.¹³⁵ A phase III, open-label, randomized study of Futibatinib versus Gemcitabine-Cisplatin Chemotherapy as first-line therapy of patients with advanced CCA harboring the FGFR2 gene rearrangement (FOENIX-CCA3) has been initiated (NCT04093362).

Derazantinib (BAL087, formerly ARQ 087)

Derazantinib is an orally-administered small molecule pan-FGFR kinase inhibitor with potent activity against FGFR1- 3. Derazantinib has demonstrated antitumor activity and a manageable safety profile in phase I study in ICC patients,¹³⁶⁻¹³⁸ and has received U.S. and EU orphan drug designation for ICC. Phase II clinical trials are currently ongoing. Basilea announced positive interim results of a phase II trial for Derazantinib in ICC patients in 2019. The interim analysis was conducted after 42 patients were included in the study, in which 29 patients received at least one post-baseline imaging assessment. The objective response rate (ORR) was 21% and the disease control rate (DCR) for patients with partial remission or stable disease was 83%. To date, the safety data obtained from all 42 patients are consistent with the results of previous clinical studies. So far, the results are encouraging.

Erdafitinib (JNJ-42756493)

Erdafitinib (JNJ-42756493) is a potent pan-FGFR 1-4 inhibitor with demonstrated anti-tumor activity in patients with metastatic urothelial cancer and CCA with FGFR alterations.¹³⁹ Asian advanced CCA patients with FGFR alterations treated with Erdafitinib in the phase IIa study (NCT02699606) had showed promising efficacy and manageable safety profile similar to that with other tumor types.¹⁴⁰

Debio 1347

Debio 1347 is a novel orally-administered small molecule, which is a highly selective FGFR 1-3 ATP competitive inhibitor. The preliminary phase I clinical trial result

showed an encouraging clinical activity and manageable treatment-emergent adverse events in solid tumors harboring a fusion of FGFR1, FGFR2 or FGFR3- (FUZE Clinical Trial) including 9 CCAs (NCT1948297).¹⁴¹ The phase II FUZE trial of Debio 1347 (NCT03834220) for patients with advanced solid tumors harboring FGFR fusions including a cohort for patients with CCA, is currently being assessed. Recent studies have reported that secondary single nucleotide variants (SNV) including p.E565A and p.L617M, appear in cells after FGFR inhibition, resulting in acquired resistance to these FGFR inhibitor therapies.¹⁴² The study confirmed the up-regulation of the PI3K/AKT/mTOR signaling pathway in drug-resistant cells, and proved that the combination of FGFR and mTOR inhibitor can desensitize cells to FGFR drug resistance.¹⁴²

2. IDH-1/2 inhibitors

Isocitrate dehydrogenase (IDH) catalyzes the conversion of isocitrate to α -ketoglutarate. The mutant forms of IDH1 and IDH2 catalyze the non-reversible accumulation of 2-hydroxyglutarate (2HG), an oncometabolite of α -ketoglutarate which is related to DNA methylation and can promote tumor cell proliferation, invasion and tumor angiogenesis.¹²⁴ Because IDH1 and IDH2 mutates in about 10-28% of ICC, small-molecule targeted inhibitors of mutant IDH1 and IDH2 have been developed and are undergoing pre-clinical and clinical trials.

Ivosidenib (AG120)

Ivosidenib is an oral IDH1 inhibitor developed by Agios and is currently approved to treat relapsed or refractory acute myeloid leukemia with an IDH1 mutation. It is now being evaluated for treatment of ICC. In the phase I trial, AG120 showed good tolerance and clinical benefit with 40% PFS rate at 6 months in patients with advanced CCA (NCT02073994). The phase III clinical trial of AG120, ClarIDHy is a global, multicenter, double-blind study randomizing 186 participants with IDH1 mutations in a 2:1 ratio to AG-120 or placebo (NCT02989857).¹⁴³ According to the report at the European Society for Medical Oncology (ESMO), AG120 (Ivosidenib) improved PFS from 1.4 months to 2.7 months compared to placebo (HR = 0.37; P < .001). 32% and 22%

of Ivosidenib-treated patients were progression-free at 6 months and 12 months respectively, while all patients receiving the placebo had disease progression at data cutoff. The risk of disease progression or death was reduced by 63% with AG120 in ClarIDHy.¹⁴⁴ Overall, Ivosidenib provides a significant improvement in PFS and OS. The IDH targeting drug Enasidenib (AG-221; a IDH2 inhibitor) which has been approved for IDH2 mutation-positive acute myeloid leukemia (AML), but has only been examined in one CCA clinical trial (NCT02273739). Although the trial began in October 2014 and was completed in June 2016, no literature has yet been published about the results of this trial so far.

FT-2102, BAY 1436032

FT-2102, BAY 1436032 which target IDH1 mutations are currently undergoing clinical trials in solid tumors with IDH1 R132 mutations (NCT03684811, NCT02746081).

3. Other novel targets

Vascular endothelial growth factors (VEGF) were found to be overexpressed in 53.8% ICCs.¹⁴⁵ Currently, several phase II or III clinical trials for treatments of biliary tract cancers (BTCs) targeting VEGF receptors (VEGFR) are ongoing, including Ramucirumab (NCT02520141), Apatinib (NCT03521219), and Surufatinib (NCT03873532). For those with BRCA1/2 mutations (3-5%) or BAP1 mutations (10%), Poly(ADP-ribose) polymerase (PARP) inhibitors Rucaparib (NCT03639935), Olaparib (NCT04042831) and Niraparib (NCT03207347) may provide some options. PARP inhibitors can compromise the DNA repair process, but this DNA single-strand damage can be converted into double-strand break (DSB) and hereby be repaired by homologous recombination (HR). If the tumor cells have defects in HR repair (including the BRCA1 /2 or BAP1 mutations), making DSB damage unable to repair, this can lead to the lethal effect of PARP inhibitors. For the approximately 5% of ICC patients who harbor PIK3CA mutation, the pan-class I PI3K inhibitor Copanlisib plus CisGem is in the phase II clinical trial (NCT02631590).¹⁴⁶ In addition, Binimetinib (MEK162), a potent inhibitor of MEK1/2, in combination with capecitabine was shown in a phase Ib clinical trial (NCT02773459) in RAS/RAF/MEK/ERK pathway activated BTC patients to have acceptable tolerability and encouraging antitumor

efficacy with the response rate and disease control rate of 17.6% and 76.5%, median PFS and OS 3.9 m and 8.0 m.¹⁴⁷

Other clinical trials assessing the efficiency of CD166 inhibitor CX-2009 (NCT03149549), NOTCH transcription complex inhibitor CB-103 (NCT03422679), and Proteasome inhibitor Bortezomib (NCT03345303) are ongoing.

- **cHCC-ICC**

To date, no standard molecular targeted therapy has been determined for cHCC-ICC. Sorafenib has been a standard of care for unresectable HCC. Because cHCC-ICC contains both the HCC and ICC elements, Sorafenib has been used in some cHCC-ICC patients. Some studies suggests outcomes with Sorafenib were poor compared with those with Platinum-containing regimens,^{9,10} However, in 2018, a clinical case reported a patient with advanced cHCC-ICC who achieved complete remission after long-term Sorafenib treatment and remained in remission after Sorafenib was withdrawn.¹⁴⁸ The efficacy of Sorafenib in cHCC-ICC needs to be further investigated in a large group of samples.

4d. Immunotherapy

PD-1/PD-L1 antibodies are immune checkpoint inhibitors that help T cells to uncover the hypocritical veil of tumor cells and restore their recognition and killing of tumor cells.¹⁴⁹ PD-1 is a negative costimulatory molecule on T cells, and PD-L1 is the ligand of PD-1 and is expressed on tumor cells. After binding, the inhibitory signals are generated, which induces T cell apoptosis, inhibits T cell activation, and prevents T cells from attacking the “invaders” with full force, acting like a brake.¹⁵⁰ Blocking PD-1 or PD-L1, the restraints on T cells will be lifted, ensuring that T cells can fully fight cancer cells.¹⁵¹

- **HCC**

1. PD-1 antibodies

Nivolumab

The PD-1 antibody Nivolumab is the first FDA approved checkpoint inhibitor for HCC. On September 22, 2017, the FDA granted accelerated approval to Nivolumab for HCC patients who have been previously treated with Sorafenib. The confirmed overall response rate assessed by RECIST 1.1, was 14.3% (95% CI: 9.2, 20.8).¹⁵² Currently, some phase III clinical trials of Nivolumab are underway including both Nivolumab monotherapy (NCT03383458, NCT02576509) or in combination with others (NCT04170556, NCT02423343, NCT03781960, NCT03510871, etc).

Pembrolizumab

Pembrolizumab is another PD-1 antibody that was granted accelerated approval for second-line therapy of advanced HCC in 2018 based on KEYNOTE 224 (NCT02702414), a single-arm, multicenter trial enrolling 104 patients with HCC.¹⁵³ Based on the excellent data of phase II keynote 224, Merck went on to conduct the phase III keynote 240 trial. The trial included patients with HCC who had not had success with previous Sorafenib treatment. The control group was treated with a placebo. The results showed that, compared with the control group, the OS results of the Keytruda group showed improvement, but did not reach a statistically significant difference (HR = 0.781 [95% CI: 0.611-0.998], p = 0.0238); PFS results also had an advantage, but did not reach statistically significant difference either (HR = 0.775 [95% CI: 0.609-0.987], p = 0.0186).¹⁵⁴ As the OS and PFS failed to achieve superiority, no formal evaluation of the key secondary endpoint ORR was performed.

The OS results

Although the phase III trial of Pembrolizumab is not satisfying, the researchers conducted an early trial (NCT03006926) of Pembrolizumab drug combined with Lenvatinib. The results are promising, with all patients except one showing tumor reduction.¹⁵⁵ In 2019, the FDA has granted the breakthrough therapy designation to the Pembrolizumab in combination with Lenvatinib for the potential first-line treatment of patients with advanced unresectable HCC who do not respond to locoregional treatment. The phase III clinical trial of Pembrolizumab plus Lenvatinib is ongoing (NCT03713593, NCT04246177).

Camrelizumab, Sintilimab, Tislelizumab, and Toripalimab

Camrelizumab, Sintilimab, Tislelizumab, and Toripalimab are four PD-1 inhibitors developed by Chinese pharmaceutical companies which all show great promise in recent clinical trials. According to the recent result published on *Lancet Oncology*, Camrelizumab showed antitumor activity in pretreated Chinese patients with advanced HCC in an open-label multicenter phase II trial (NCT02989922) and displayed manageable toxicities.¹⁵⁶ 14.7% of patients displayed an objective response (n = 32 of 217; 14.7%) and among all patients, 6-month overall survival rate was 74.4% (95% CI, 68.0-79.7). On March 4, 2020, Camrelizumab was officially approved by the National Medical Products Administration (NMPA) for patients with advanced HCC who have received Sorafenib treatment and/or oxaliplatin-containing chemotherapy. This is the first PD-1 inhibitor approved for liver cancer indications in China.

Sintilimab received its first approval for the treatment of classical Hodgkin's lymphoma in China in 2018.¹⁵⁷ Currently, Sintilimab in combination with Anlotinib (NCT04042805), Lenvatinib (NCT04042805) and IBI305 (NCT03794440) are undergoing clinical trials for the treatment of various solid tumors including HCC. Recently, Meihua Lin and his colleagues reported a case that an ICC patient who, after the first-line chemotherapy failed, achieved complete remission after three cycles of Sintilimab treatment with only mild adverse reactions.¹⁵⁸ Unlike other PD-1 antibodies, Tislelizumab is specifically designed to minimize binding to Fc γ R on macrophages¹⁵⁹ and to escape Fc γ R1-mediated effector function, because Fc γ R on macrophages impairs the anti-tumor activity of PD-1 antibodies by activating antibody-dependent macrophage-mediated T effector cell killing.¹⁶⁰ Phase Ia/Ib trials have shown promise for HCC.¹⁶¹ The global, phase III clinical trial (NCT03412773) designed to evaluate the efficacy and safety of Tislelizumab compared with Sorafenib as a potential first-line treatment of unresectable HCC has been initiated.¹⁶² Toripalimab, being developed by Shanghai Junshi Bioscience Co., Ltd, has received approval for the treatment of unresectable melanoma patients who failed previous systemic therapy in China.¹⁶³ Several clinical studies are currently being conducted to test the safety and efficiency of

Toripalimab in the treatment of HCC (NCT03412773, NCT04368078). These studies showed that the four domestic drugs showed great antitumor activity and efficiency and could be a first or second-line treatment option for advanced HCC patients, even for a population with a high proportion of patients with HBV infection.

2. PD-L1 antibodies

Atezolizumab, Bevacizumab

In July 2018, the PD-L1 monoclonal antibody Atezolizumab in combination Bevacizumab was awarded the FDA designation of a breakthrough therapy in the treatment of advanced HCC based on a phase Ib clinical study (NCT02715531). In this study, patients with advanced unresectable or metastatic hepatocellular carcinoma were included. Atezolizumab 1200mg + Bevacizumab 15mg/kg was given once every 3 weeks. The median overall survival (OS) was 17.1 months, 6-month OS was 82%, and 12-month OS was 63%.¹⁶⁴ Just recently, the NEJM published the exciting phase III trial result,¹⁶⁵ the combination of Atezolizumab and Bevacizumab significantly improved OS and PFS (6.8m vs 4.3 m) in patients with unresectable HCC with the similar toxicity to that of Sorafenib (Grade 3 or 4 adverse events 56.5% vs 55.1%). The 12-month overall survival rate of the patients increased to 67.2% compared to 54.6% with Sorafenib, breaking the long-standing bottleneck in liver cancer treatment. The combination is currently under review by the FDA, and it is possible that it will become the new standard of care later in 2020.

Durvalumab

Durvalumab is an FDA-approved immunotherapy first used for locally advanced or metastatic urothelial carcinoma and non-small cell lung cancer (NSCLC) developed by Medimmune/AstraZeneca. It is a human immunoglobulin G1 kappa monoclonal antibody that blocks the interaction of PD-L1 with PD-1. Based on the promising results in other solid tumors, several phase II/III clinical trials are being conducted for HCC. Durvalumab in combination with Bevacizumab (NCT03847428) and with Tremelimumab (NCT03298451) are currently under phase III evaluation.

3. CTLA-4 antibodies

Cytotoxic T lymphocyte-associated antigen-4 (CTLA-4) is another a co-inhibitory molecule that functions to regulate immune responses.¹⁶⁶ Antibodies that block the interaction of CTLA-4 with its ligands B7-1/ B7-2 can enhance T cell activation as well as anti-tumor efficacy.¹⁶⁷ Two CTLA-4 antibodies which are currently under clinical investigation are Tremelimumab and Ipilimumab.

Tremelimumab

Tremelimumab is a human monoclonal CTLA-4 antibody. In 2020 the FDA granted an orphan drug designation to Tremelimumab plus Durvalumab for the treatment of patients with HCC. The combination is being tested in a phase III clinical trial (NCT03298451) to evaluate Durvalumab alone and in combination with Tremelimumab compared with standard Sorafenib in 1310 patients with unresectable, advanced HCC who have not received prior systemic treatment and are ineligible for locoregional therapy.¹⁶⁸ Results from an early phase II trial combining Tremelimumab and Durvalumab demonstrated a safety and a promising antitumor activity both in HCC and BTC.¹⁶⁹ Tremelimumab and Durvalumab combination holds great promise in becoming a new first-line treatment for liver cancer.

Ipilimumab

Ipilimumab is another CTLA-4 monoclonal antibody intended to activate the immune system. On 10 March 2020, the FDA granted the first combination therapy accelerated approval for treatment of HCC to the combination of Ipilimumab and Nivolumab for HCC patients intolerant to Sorafenib. The approval was based on the favorable overall response rate (ORR) and duration of response (DoR) from cohort 4 of the CHECKMATE-040 (NCT01658878) trial which included a total of 49 patients who received Nivolumab in combination with Ipilimumab. The ORR was 33% (n=16; 95% CI: 20, 48) with 4 complete responses and 12 partial responses and DoR ranged from 4.6 to 30.5+ months, with 31% of responses lasting at least 24 months.

4. Neoantigen-based therapy

New and rapidly growing cancer immunotherapy treatments includes the development of personalized tumor vaccines which target neoantigens. In most tumor patients, there are certain specific T cells that can recognize short peptide antigens presented by MHC on the surface of cancer cells. This short peptide antigen, which can induce specific T cells to eliminate cancer cells, does not exist in normal tissues, and is thus called a tumor specific antigen or neoantigens (Neoantigens).¹⁷⁰ Unlike traditional vaccines, which are limited by the dual restrictions of human leukocyte antigen (HLA) diversity and expression and personalized neoantigen vaccines target each patient's tumor tissue mutation antigen, combining precise gene detection and tumor immunotherapy. This approach uses specific tumor gene mutations to design vaccines which stimulate patients' autoimmunity to kill eventually tumor cells.^{171,172} Neoantigens are mostly caused by errors in the DNA replication process of cancer cells, and some are caused by environmental factors such as viruses, radiation, and chemicals.¹⁷³ Although personalized tumor vaccines are still in the exploratory stage, the currently reported clinical trials of individualized neoantigen vaccines have shown encouraging results especially in treatment of melanoma with high accuracy and low side effects.¹⁷⁴⁻¹⁸⁰

Dendritic cells are the most effective antigen presenting cells in the body. After recognizing the antigen, dendritic cells are activated and enhance the anti-tumor immune response through T cells and NK cells.¹⁸¹ Currently, several neoantigens based on personalized dendritic cells vaccines for HCC patients are under investigation in multiple ongoing clinical trials (NCT03674073, NCT04147078, NCT03942328) (summarized in Table 2 - Immune cell). We expect that follow-up trials can achieve good results and realize its potential to bring patients efficient, safe and truly personalized tumor vaccine as soon as possible.

- **ICC**

Tumor with mismatch repair (MMR) pathway deficiency have been demonstrated to have favorable responses to PD-1 blockade immunotherapy.¹⁸² Mismatch repair

deficiency (dMMR) tumors cause high levels of microsatellite instability (MSI) and can generate neoantigens which make the cancer cells susceptible to inhibition of the PD-L1/PD-1 interaction and sensitive to immunotherapy.¹⁸³ MSI is most commonly seen in colorectal and endometrial cancers, however, CCA has also been reported to exhibit MSI with a frequency above 10%.^{184,185} Several clinical immunotherapies for ICC are currently in use, including the PD-1 antibody Pembrolizumab and Nivolumab, the PD-L1 antibody Durvalumab and the CTLA-4 antibodies Ipilimumab and Tremelimumab (Figure2).¹⁸⁶

Pembrolizumab

On May 23, 2017, Pembrolizumab was granted accelerated approval by the FDA for the treatment of patients with microsatellite instability-high (MSI-H), or dMMR solid tumors. This was the first time that FDA approved a drug based on the genetic profile instead of the primary tumor site.¹⁸⁷ A phase II, multicohort KEYNOTE-158 study (NCT02628067) evaluated the antitumor activity and safety of Pembrolizumab in patients with advanced solid tumors including 104 CCA patients.¹⁸⁸ Median PFS was 1.9 m vs 2.1 m, median OS was 7.2 m vs 9.6 m and ORR was 6.6% vs 2.9% in patients with PD-L1 combined positive score CPS ≥ 1 vs CPS < 1 . All responders were not MSI-High. Pembrolizumab in another phase Ib study (NCT02054806) with 24 BTC also showed durable antitumor activity regardless of PD-L1 CPS and had manageable toxicity.¹⁸⁸ Currently, several Pembrolizumab clinical trials are ongoing, both monotherapy (NCT03110328) and in combination with others therapies which include the standard first-line care drug - CisGem (NCT03260712), the RARP inhibitor - Olaparib (NCT04306367), the bispecific antibody that simultaneously targets immune checkpoint receptors CTLA-4 and LAG-3 - XmAb22841 (NCT03849469), and the immune cell therapy (NCT03937895).

Nivolumab

In a Japanese multicenter, open-label, phase I trial, researchers found Nivolumab showed activity against BTC that have progressed on prior systemic therapies, with a

manageable safety profile in patients with unresectable or recurrent BTC.¹⁸⁹ The median patient age was 64.5 years old. Two-thirds of the patients (64.7%) had intrahepatic cholangiocarcinoma, 2.9% had extrahepatic cholangiocarcinoma, and 32.4% had tumors of the gallbladder. The median OS was 5.2 months in Nivolumab monotherapy and 15.4 months in Nivolumab plus CisGem, with the median PFS 1.4 months and 4.2 months, respectively. A phase II study (NCT02829918) also found Nivolumab to have promising efficacy with tolerated toxicity including durable responses lasting 2 years in BTC.¹⁹⁰ These initial assessments of Nivolumab for the treatment of advanced BTC provides supportive evidence for future larger randomized studies of Nivolumab in this refractory cancer.

Durvalumab (D) and Tremelimumab (T)

A phase I, open-label, multicenter Study (NCT01938612) evaluated the safety, tolerability, and pharmacokinetics of Durvalumab and Tremelimumab in Asian patients with advanced solid tumors including BTC.¹⁹¹ Patients were enrolled in Durvalumab D (n = 42) and Durvalumab plus Tremelimumab (D+T) cohorts (n = 65). Promising clinical efficacy was observed in both groups with no unexpected toxicities. Currently, a phase II trial of Durvalumab and Tremelimumab (NCT04238637) is ongoing.

The first randomized, double-blind, international phase III clinical trial to evaluate immunotherapy plus chemotherapy in patients with BTC in the first-line setting is also in progress, testing Durvalumab in combination with Gemcitabine plus Cisplatin (NCT03875235).¹⁹²

- **cHCC-ICC**

Currently, few studies in the literature or clinical trials have focused on the use of immunotherapy for treatment of cHCC-ICC. Therefore, it will not be discussed in this review.

Currently, the drug development field for liver cancer is mainly dominated by antibody drugs, of which PD-1/PD-L1 and CTLA-4 are the main targets, and VEGFR and BRAF are the main small molecule inhibitor targets. Among the first-line treatment research and development drugs, the combination of Roche's Atezolizumab and Bevacizumab is the most promising first-line treatment for HCC globally, while Suzhou Zelgen Biopharmaceuticals's Donafenib is expected to become the first-line treatment in China. Among the drugs developed for second-line treatment, Hengrui Medicine's Apatinib can significantly improve the overall survival (OS) of HCC patients, and is expected to become a new second-line therapy for HCC.

5. Future perspective

Although many clinical drugs have been approved or tested in advanced HCC and ICC, the median PFS and OS remain dismal. One of the reasons is the acquired drug resistance due to the intra-tumor heterogeneity or the continuous diversification during treatment which allows certain tumor cells to survival and eventually develop a drug-resistant phenotype. This remains the huge hurdle for the long-term use of targeted therapies for PLC.^{193,194} It is therefore necessary to further explore the mechanism of drug resistance. Recently, Tang, J., et al. reported a novel somatic mutation in OCT4 (c.G52C) associated with Sorafenib resistance.¹⁹⁵ Further work in this vein will allow us to understand the mechanism and the exact gene mutation responsible for the drug resistance, allowing for targeting of specific mutation sites, thereby hopefully overcoming drug resistance.

Another challenge for targeted therapies in PLC is lack of precise targets and biomarkers. Unlike breast cancer, which has the precise biomarker HER2, PLC has a high degree of heterogeneity and genomic diversity and with no accurate biomarkers. Although many high-frequency mutant genes such as TERT, TP53, CTNNB1, and KRAS have been confirmed in PLC, it is still not clear whether they play the role of "driver gene" or "passenger gene" in the progression of liver cancer, which limits the development of targeted drugs. There is therefore an unmet need to comprehensively understand the genomic architecture, define the mutation landscape, and identify novel biomarkers and driver genes in order to develop new therapeutic interventions. With

this information, future clinical trials could employ precision medicine to treat patients based on specific genetic mutation and drivers. Another point of concern is that PLC has a high recurrence rate; more than 70% of patients will relapse within five years after surgery.¹⁹⁶ Thus, whether the genetic features remain the same in the primary and recurrent tumors is also worth exploring.

In recent years, immune checkpoint inhibitors (ICIs) have been of increased research interest. The 2018 Nobel Prize in Physiology or Medicine was awarded to American immunologist James Allison and Japanese immunologist Tasuku Honjo for their contributions to the tumor immunity field, leading to the development and progression of PD-1 / CTLA-4 inhibitors and other immunotherapy drugs. However, the overall response rate for ICIs has not been very high (10 - 20% in PLCs).^{152,153} which means the majority of patients cannot benefit from ICIs. It has been the main issue for ICIs. Fortunately, recent studies suggest that ICIs combined with other treatments, especially VEGF/VEGFR inhibitors, can significantly improve the overall response rate, with the prolonged median PFS and OS. For example, the overall response rate of the Atezolizumab (a PD-L1 inhibitor) and Bevacizumab's (a VEGF inhibitor) combination was 62% in the phase Ib clinical trial and 27% in the phase III trial. Combined therapies are therefore under more study currently. Combination therapies (Table2; Combined therapies) including Pembrolizumab (PD-1 inhibitor) plus Lenvatinib (VEGFR inhibitor) (NCT03713593), Atezolizumab (PD-L1 inhibitor) plus Cabozantinib (VEGFR inhibitor) (NCT03755791), Durvalumab (PD-L1 inhibitor) plus Bevacizumab (VEGF inhibitor) (NCT03847428), CS1003 (PD-1 inhibitor) plus Lenvatinib (VEGFR inhibitor) (NCT04194775) and Camrelizumab (PD-1 inhibitor) plus Apatinib (VEGFR inhibitor) (NCT03764293) are all in phase III clinical trials for HCC, we are eagerly waiting for the results.

The immunotolerance of the liver protects it from autoimmune damage caused by foreign antigens,¹⁹⁷ but also helps liver cancer cells to escape immune cells hunting. A decrease in NK cell number or impairment of function, accumulation of regulatory T cells and exhausted CD8+ T cells have all been seen in HCC tumors, implicating an immunosuppressive microenvironment.¹⁹⁸ Many patients cannot respond to immunotherapy with a low response rate due to an insufficient immune activation. Thus, how to turn a “cold tumor” (immune tolerant) into a “hot tumor” (immunogenic)

remains a major challenge for current tumor immunotherapy research and development. Future efforts in immunotherapy should be made in two directions: boosting the existing immune response and stimulating a *de novo* immune response.

For the first, more combination strategies such as ICIs combined with VEGFR, CTLA-4, or CDKs should be developed. In addition, immunoregulating the function of Treg and CD8⁺ T cells function is also of great importance. For example, previous studies have shown that TGF- β promotes tumor immune escape by inducing Treg cell differentiation. The mouse model confirmed that TGF β inhibitor SM-16 administration reduces Treg cell frequency, resulting in a reduced development of HCC,¹⁹⁹ providing a mechanistic rationale for the combination of TGF- β inhibitor and ICI in liver cancer.

Cyclic peptide C25 targeting human LAG-3 protein is reported to be able to significantly stimulate CD8⁺ T cell activation in human PBMCs, resulting in inhibition of tumor growth in CT26, B16, and B16-OVA bearing mice.²⁰⁰ The use of C25 and blockade of the LAG-3/HLA-DR interaction may also provide an alternative method for cancer immunotherapy. The bispecific anti-PD-1/LAG-3 antibodies are also promising in the future cancer treatment.

For stimulating the *de novo* immune response, cell-based immunotherapies such as adoptive cell therapies, including CAR-T cell therapies (e.g. NCT03993743, NCT04121273, and NCT03941626), TCR-T cell therapies (e.g. NCT03441100), and the vaccine-based therapies such as neoantigen based vaccines (e.g. NCT03674073), peptide vaccines and oncolytic virus drugs (NCT03071094, etc.) are currently being evaluated in HCC. We look forward to seeing the evaluation. Also, more research and studies are expected to be conducted in the future.

Recent studies have begun to unveil the complex hepatic immune microenvironment. Further work is required to decipher the intricate immune microenvironment of liver cancer such as the function and subtype of diverse immune cell subsets in liver including T cells, B cells, macrophages, neutrophils, DCs, myeloid-derived suppressor cells, NK cells, and cancer-associated fibroblasts, as well as the dynamic interaction between the immune cells and the tumor ecosystem. Solving these will help us take a

deep look inside the tumor microenvironment and understand patient's responses to immune therapy, and develop more immunotherapy options.

Aside from these mainstream treatments, some novel therapies have also been proposed in the management of PLC. For example, René Bernards and his team recently elaborated some new idea about combined therapies by devising a “one - two punch” method (named after the effective combination of two rapid consecutive moves in boxing).²⁰¹ The “first punch” makes use of a specific mutations (like TP53) in tumor cells to specifically induce it to a certain state like cell senescence, and then the next "second punch" precisely removes aging tumor cells. Therefore, although these two drugs are not used at the same time, they have synergistic effect with reduced toxicity and high precision.

The treatment of PLC is gradually shifting away from traditional chemotherapy, and toward targeted therapy including immunotherapy and especially the combination therapy. These new approaches have shown great potential in the clinical trials, and there is a need to develop more combination strategies or try novel combinations of the previously studied drugs.

Despite these past and ongoing trials investigating PLC treatments, publications and clinical trials regarding systematic treatments of the rare cHCC-ICC are still extremely limited. Further study is undoubtedly required to further improve current diagnosis, as well as to better understand the genomic profile and pathogenesis of cHCC-ICC in order to develop novel therapeutics.

Acknowledgments

We thank Zichen Xu for assistance in the preparation of the figures. This work was supported by the National Natural Science Foundation of China (81972735, 31671452, 81970459) and Clinical Medicine Plus X-Young Scholars Project of Perking University (71006Y2435).

References :

1. Brunt, E., Aishima, S., Clavien, P.-A., et al. (2018). cHCC-CCA: Consensus terminology for primary liver carcinomas with both hepatocytic and cholangiocytic differentiation. *Hepatology*. 68, 113-126.
2. Nagtegaal, I. D., Odze, R. D., Klimstra, D., et al. (2020). The 2019 WHO classification of tumours of the digestive system. *Histopathology*. 76, 182-188.
3. Bray, F., Ferlay, J., Soerjomataram, I., et al. (2018). Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 68, 394-424.
4. Yang, J. D., Hainaut, P., Gores, G. J., et al. (2019). A global view of hepatocellular carcinoma: trends, risk, prevention and management. *Nat Rev Gastroenterol Hepatol*. 16, 589-604.
5. Zheng, R., Qu, C., Zhang, S., et al. (2018). Liver cancer incidence and mortality in China: Temporal trends and projections to 2030. *Chinese journal of cancer research = Chung-kuo yen cheng yen chiu*. 30, 571-579.
6. McGlynn, K. A., Petrick, J. L., El-Serag, H. B. (2020). Epidemiology of Hepatocellular Carcinoma. *Hepatology*.
7. Clements, O., Eliahoo, J., Kim, J. U., et al. (2020). Risk factors for intrahepatic and extrahepatic cholangiocarcinoma: A systematic review and meta-analysis. *J Hepatol*. 72, 95-103.
8. Valle, J. W., Borbath, I., Khan, S. A., et al. (2016). Biliary cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 27, v28-v37.
9. Trikalinos, N. A., Zhou, A., Doyle, M. B. M., et al. (2018). Systemic Therapy for Combined Hepatocellular-Cholangiocarcinoma: A Single-Institution Experience. *J Natl Compr Canc Netw*. 16, 1193-1199.
10. Kobayashi, S., Terashima, T., Shiba, S., et al. (2018). Multicenter retrospective analysis of systemic chemotherapy for unresectable combined hepatocellular and cholangiocarcinoma. *Cancer Sci*. 109, 2549-2557.
11. Goodman, Z. D. (2007). Neoplasms of the liver. *Mod Pathol*. 20 Suppl 1, S49-60.
12. Jiang, K., Al-Diffhala, S., Centeno, B. A. (2018). Primary Liver Cancers-Part 1: Histopathology, Differential Diagnoses, and Risk Stratification. *Cancer Control*. 25, 1073274817744625.
13. Jain, D. (2014). Tissue diagnosis of hepatocellular carcinoma. *J Clin Exp Hepatol*. 4, S67-73.
14. Komuta, M., Govaere, O., Vandecaveye, V., et al. (2012). Histological diversity in cholangiocellular carcinoma reflects the different cholangiocyte phenotypes. *Hepatology*. 55, 1876-1888.
15. Razumilava, N., Gores, G. J. (2014). Cholangiocarcinoma. *Lancet*. 383, 2168-2179.
16. Kendall, T., Verheij, J., Gaudio, E., et al. (2019). Anatomical, histomorphological and molecular classification of cholangiocarcinoma. *Liver Int*. 39 Suppl 1, 7-18.
17. Lee, A. J., Chun, Y. S. (2018). Intrahepatic cholangiocarcinoma: the AJCC/UICC 8th edition updates. *Chinese clinical oncology*. 7, 52.

18. Allen, R. A., Lisa, J. R. (1949). Combined liver cell and bile duct carcinoma. *Am J Pathol.* 25, 647-655.
19. Goodman, Z. D., Ishak, K. G., Langloss, J. M., et al. (1985). Combined hepatocellular-cholangiocarcinoma. A histologic and immunohistochemical study. *Cancer.* 55, 124-135.
20. Zhang, F., Chen, X. P., Zhang, W., et al. (2008). Combined hepatocellular cholangiocarcinoma originating from hepatic progenitor cells: immunohistochemical and double-fluorescence immunostaining evidence. *Histopathology.* 52, 224-232.
21. Vilchez, V., Shah, M. B., Daily, M. F., et al. (2016). Long-term outcome of patients undergoing liver transplantation for mixed hepatocellular carcinoma and cholangiocarcinoma: an analysis of the UNOS database. *HPB (Oxford).* 18, 29-34.
22. Totoki, Y., Tatsuno, K., Yamamoto, S., et al. (2011). High-resolution characterization of a hepatocellular carcinoma genome. *Nat Genet.* 43, 464-469.
23. Fujimoto, A., Totoki, Y., Abe, T., et al. (2012). Whole-genome sequencing of liver cancers identifies etiological influences on mutation patterns and recurrent mutations in chromatin regulators. *Nat Genet.* 44, 760-764.
24. Guichard, C., Amaddeo, G., Imbeaud, S., et al. (2012). Integrated analysis of somatic mutations and focal copy-number changes identifies key genes and pathways in hepatocellular carcinoma. *Nat Genet.* 44, 694-698.
25. Cleary, S. P., Jeck, W. R., Zhao, X., et al. (2013). Identification of driver genes in hepatocellular carcinoma by exome sequencing. *Hepatology.* 58, 1693-1702.
26. Totoki, Y., Tatsuno, K., Covington, K. R., et al. (2014). Trans-ancestry mutational landscape of hepatocellular carcinoma genomes. *Nat Genet.* 46, 1267-1273.
27. Amaddeo, G., Cao, Q., Ladeiro, Y., et al. (2015). Integration of tumour and viral genomic characterizations in HBV-related hepatocellular carcinomas. *Gut.* 64, 820-829.
28. Romualdo, G. R., Grassi, T. F., Goto, R. L., et al. (2017). An integrative analysis of chemically-induced cirrhosis-associated hepatocarcinogenesis: Histological, biochemical and molecular features. *Toxicol Lett.* 281, 84-94.
29. Calderaro, J., Couchy, G., Imbeaud, S., et al. (2017). Histological subtypes of hepatocellular carcinoma are related to gene mutations and molecular tumour classification. *J Hepatol.* 67, 727-738.
30. Ong, C. K., Subimerb, C., Pairojkul, C., et al. (2012). Exome sequencing of liver fluke-associated cholangiocarcinoma. *Nat Genet.* 44, 690-693.
31. Chan-On, W., Nairismagi, M. L., Ong, C. K., et al. (2013). Exome sequencing identifies distinct mutational patterns in liver fluke-related and non-infection-related bile duct cancers. *Nat Genet.* 45, 1474-1478.
32. Jiao, Y., Pawlik, T. M., Anders, R. A., et al. (2013). Exome sequencing identifies frequent inactivating mutations in BAP1, ARID1A and PBRM1 in intrahepatic cholangiocarcinomas. *Nat Genet.* 45, 1470-1473.
33. Sia, D., Losic, B., Moeini, A., et al. (2015). Massive parallel sequencing uncovers actionable FGFR2-PPHLN1 fusion and ARAF mutations in intrahepatic cholangiocarcinoma. *Nat Commun.* 6, 6087.

34. Shiao, M.-S., Chiablaem, K., Charoensawan, V., et al. (2018). Emergence of Intrahepatic Cholangiocarcinoma: How High-Throughput Technologies Expedite the Solutions for a Rare Cancer Type. *Frontiers in Genetics*. 9
35. Liu, Z. H., Lian, B. F., Dong, Q. Z., et al. (2018). Whole-exome mutational and transcriptional landscapes of combined hepatocellular cholangiocarcinoma and intrahepatic cholangiocarcinoma reveal molecular diversity. *Biochim Biophys Acta Mol Basis Dis*. 1864, 2360-2368.
36. Xue, R., Chen, L., Zhang, C., et al. (2019). Genomic and Transcriptomic Profiling of Combined Hepatocellular and Intrahepatic Cholangiocarcinoma Reveals Distinct Molecular Subtypes. *Cancer Cell*. 35, 932-947 e938.
37. Katz, S. F., Lechel, A., Obenauf, A. C., et al. (2012). Disruption of Trp53 in livers of mice induces formation of carcinomas with bilineal differentiation. *Gastroenterology*. 142, 1229-1239 e1223.
38. Zhu, R. X., Seto, W.-K., Lai, C.-L., et al. (2016). Epidemiology of Hepatocellular Carcinoma in the Asia-Pacific Region. *Gut Liver*. 10, 332-339.
39. Valery, P. C., Laversanne, M., Clark, P. J., et al. (2018). Projections of primary liver cancer to 2030 in 30 countries worldwide. *Hepatology*. 67, 600-611.
40. Cazals-Hatem, D., Rebouissou, S., Bioulac-Sage, P., et al. (2004). Clinical and molecular analysis of combined hepatocellular-cholangiocarcinomas. *J Hepatol*. 41, 292-298.
41. Ramai, D., Ofosu, A., Lai, J. K., et al. (2019). Combined Hepatocellular Cholangiocarcinoma: A Population-Based Retrospective Study. *Am J Gastroenterol*. 114
42. Tang, R., Latchana, N., Rahnemai-Azar, A. A., et al. (2018) Guidelines for Resection of Intrahepatic Cholangiocarcinoma. in *Primary and Metastatic Liver Tumors: Treatment Strategy and Evolving Therapies* (Cardona, K., Maithel, S. K. eds.), Springer International Publishing, Cham. pp 99-110
43. Bosman FT, C. F., Hruban R, Theise ND. (2010). WHO Classification of Tumours of the Digestive System. IARC Press: Lyon; 2010.
44. Schulze, K., Nault, J. C., Villanueva, A. (2016). Genetic profiling of hepatocellular carcinoma using next-generation sequencing. *J Hepatol*. 65, 1031-1042.
45. Li, Z., Zhang, J., Liu, X., et al. (2018). The LINC01138 drives malignancies via activating arginine methyltransferase 5 in hepatocellular carcinoma. *Nature Communications*. 9, 1572-1572.
46. Llovet, J. M. (2014). Focal gains of VEGFA: candidate predictors of sorafenib response in hepatocellular carcinoma. *Cancer cell*. 25, 560-562.
47. Cai, Y., Crowther, J., Pastor, T., et al. (2016). Loss of Chromosome 8p Governs Tumor Progression and Drug Response by Altering Lipid Metabolism. *Cancer Cell*. 29, 751-766.
48. Zhang, J., Li, Z., Liu, L., et al. (2018). Long noncoding RNA TSLNC8 is a tumor suppressor that inactivates the interleukin-6/STAT3 signaling pathway. *Hepatology*. 67, 171-187.
49. Nault, J. C., Calderaro, J., Di Tommaso, L., et al. (2014). Telomerase reverse transcriptase promoter mutation is an early somatic genetic alteration in the

- transformation of premalignant nodules in hepatocellular carcinoma on cirrhosis. *Hepatology*. 60, 1983-1992.
50. Javle, M., Bekaii-Saab, T., Jain, A., et al. (2016). Biliary cancer: Utility of next-generation sequencing for clinical management. *Cancer*. 122, 3838-3847.
51. Nakamura, H., Arai, Y., Totoki, Y., et al. (2015). Genomic spectra of biliary tract cancer. *Nat Genet*. 47, 1003-1010.
52. Jusakul, A., Cutcutache, I., Yong, C. H., et al. (2017). Whole-Genome and Epigenomic Landscapes of Etiologically Distinct Subtypes of Cholangiocarcinoma. *Cancer Discov*. 7, 1116-1135.
53. O'Dell, M. R., Huang, J. L., Whitney-Miller, C. L., et al. (2012). Kras(G12D) and p53 mutation cause primary intrahepatic cholangiocarcinoma. *Cancer Res*. 72, 1557-1567.
54. Saha, S. K., Parachoniak, C. A., Ghanta, K. S., et al. (2014). Mutant IDH inhibits HNF-4alpha to block hepatocyte differentiation and promote biliary cancer. *Nature*. 513, 110-114.
55. Keller, J. W., Doyle, M. M., Wang-Gillam, A., et al. (2014). Analysis of the genomic profile of biphenotypic tumors compared to cholangiocarcinoma and hepatocellular carcinoma. *J Clin Oncol*. 32, 226-226.
56. Sasaki, M., Sato, Y., Nakanuma, Y. (2017). Mutational landscape of combined hepatocellular carcinoma and cholangiocarcinoma, and its clinicopathological significance. *Histopathology*. 70, 423-434.
57. Cai, X., Li, H., Kaplan, D. E. (2020). Murine hepatoblast-derived liver tumors resembling human combined hepatocellular-cholangiocarcinoma with stem cell features. *Cell & bioscience*. 10, 38.
58. Vogel, A., Cervantes, A., Chau, I., et al. (2019). Hepatocellular carcinoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Annals of oncology : official journal of the European Society for Medical Oncology*. 30, 871-873.
59. Wang, K., Zhang, H., Xia, Y., et al. (2017). Surgical options for intrahepatic cholangiocarcinoma. *Hepatobiliary Surg Nutr*. 6, 79-90.
60. Ibrahim, S. M., Mulcahy, M. F., Lewandowski, R. J., et al. (2008). Treatment of Unresectable Cholangiocarcinoma Using Yttrium-90 Microspheres Results From a Pilot Study. *Cancer*. 113, 2119-2128.
61. Hyder, O., Marsh, J. W., Salem, R., et al. (2013). Intra-arterial therapy for advanced intrahepatic cholangiocarcinoma: a multi-institutional analysis. *Ann Surg Oncol*. 20, 3779-3786.
62. Boehm, L. M., Jayakrishnan, T. T., Miura, J. T., et al. (2015). Comparative effectiveness of hepatic artery based therapies for unresectable intrahepatic cholangiocarcinoma. *J Surg Oncol*. 111, 213-220.
63. Han, K., Ko, H. K., Kim, K. W., et al. (2015). Radiofrequency ablation in the treatment of unresectable intrahepatic cholangiocarcinoma: systematic review and meta-analysis. *J Vasc Interv Radiol*. 26, 943-948.

64. Edeline, J., Toucheffeu, Y., Guiu, B., et al. (2020). Radioembolization Plus Chemotherapy for First-line Treatment of Locally Advanced Intrahepatic Cholangiocarcinoma A Phase 2 Clinical Trial. *Jama Oncology*. 6, 51-59.
65. Garancini, M., Goffredo, P., Pagni, F., et al. (2014). Combined hepatocellular-cholangiocarcinoma: a population-level analysis of an uncommon primary liver tumor. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 20, 952-959.
66. Cai, X., Zhai, J., Kaplan, D. E., et al. (2012). Background progenitor activation is associated with recurrence after hepatectomy of combined hepatocellular-cholangiocarcinoma. *Hepatology*. 56, 1804-1816.
67. Song, P., Midorikawa, Y., Nakayama, H., et al. (2019). Patients' prognosis of intrahepatic cholangiocarcinoma and combined hepatocellular-cholangiocarcinoma after resection. *Cancer Med*. 8, 5862-5871.
68. De Martin, E., Rayar, M., Golse, N., et al. (2020). Analysis of Liver Resection versus Liver Transplantation on Outcome of Small Intrahepatic Cholangiocarcinoma and Combined Hepatocellular-Cholangiocarcinoma in the setting of cirrhosis. *Liver Transpl*.
69. Malone, C. D., Gibby, W., Tsai, R., et al. (2020). Outcomes of Yttrium-90 Radioembolization for Unresectable Combined Biphenotypic Hepatocellular-Cholangiocarcinoma. *J Vasc Interv Radiol*. 31, 701-709.
70. Holzner, M. L., Tabrizian, P., Parvin-Nejad, F. P., et al. (2020). Resection of Mixed Hepatocellular-Cholangiocarcinoma, Hepatocellular Carcinoma, and Intrahepatic Cholangiocarcinoma. *Liver Transpl*. 26, 888-898.
71. Louafi, S., Boige, V., Ducreux, M., et al. (2007). Gemcitabine plus oxaliplatin (GEMOX) in patients with advanced hepatocellular carcinoma (HCC): results of a phase II study. *Cancer*. 109, 1384-1390.
72. Qin, S., Bai, Y., Lim, H. Y., et al. (2013). Randomized, multicenter, open-label study of oxaliplatin plus fluorouracil/leucovorin versus doxorubicin as palliative chemotherapy in patients with advanced hepatocellular carcinoma from Asia. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 31, 3501-3508.
73. Abou-Alfa, G. K., Niedzwieski, D., Knox, J. J., et al. (2016). Phase III randomized study of sorafenib plus doxorubicin versus sorafenib in patients with advanced hepatocellular carcinoma (HCC): CALGB 80802 (Alliance). *J Clin Oncol*. 34, 192-192.
74. Liu, L., Zheng, Y. H., Han, L., et al. (2016). Efficacy and safety of the oxaliplatin-based chemotherapy in the treatment of advanced primary hepatocellular carcinoma: A meta-analysis of prospective studies. *Medicine (Baltimore)*. 95, e4993.
75. Valle, J., Wasan, H., Palmer, D. H., et al. (2010). Cisplatin plus gemcitabine versus gemcitabine for biliary tract cancer. *N Engl J Med*. 362, 1273-1281.
76. Okusaka, T., Nakachi, K., Fukutomi, A., et al. (2010). Gemcitabine alone or in combination with cisplatin in patients with biliary tract cancer: a comparative multicentre study in Japan. *Br J Cancer*. 103, 469-474.

- 77.Raderer, M., Hejna, M. H., Valencak, J. B., et al. (1999). Two consecutive phase II studies of 5-fluorouracil/leucovorin/mitomycin C and of gemcitabine in patients with advanced biliary cancer. *Oncology*. 56, 177-180.
- 78.Andre, T., Tournigand, C., Rosmorduc, O., et al. (2004). Gemcitabine combined with oxaliplatin (GEMOX) in advanced biliary tract adenocarcinoma: a GERCOR study. *Ann Oncol*. 15, 1339-1343.
- 79.Cho, J. Y., Paik, Y. H., Chang, Y. S., et al. (2005). Capecitabine combined with gemcitabine (CapGem) as first-line treatment in patients with advanced/metastatic biliary tract carcinoma. *Cancer*. 104, 2753-2758.
- 80.Knox, J. J., Hedley, D., Oza, A., et al. (2005). Combining gemcitabine and capecitabine in patients with advanced biliary cancer: a phase II trial. *J Clin Oncol*. 23, 2332-2338.
- 81.Harder, J., Riecken, B., Kummer, O., et al. (2006). Outpatient chemotherapy with gemcitabine and oxaliplatin in patients with biliary tract cancer. *Br J Cancer*. 95, 848-852.
- 82.Manziona, L., Romano, R., Germano, D. (2007). Chemotherapy with gemcitabine and oxaliplatin in patients with advanced biliary tract cancer: a single-institution experience. *Oncology*. 73, 311-315.
- 83.Koeberle, D., Saletti, P., Borner, M., et al. (2008). Patient-reported outcomes of patients with advanced biliary tract cancers receiving gemcitabine plus capecitabine: A multicenter, phase II trial of the Swiss Group for Clinical Cancer Research. *J Clin Oncol*. 26, 3702-3708.
- 84.Nehls, O., Oettle, H., Hartmann, J. T., et al. (2008). Capecitabine plus oxaliplatin as first-line treatment in patients with advanced biliary system adenocarcinoma: a prospective multicentre phase II trial. *Br J Cancer*. 98, 309-315.
- 85.Andre, T., Reyes-Vidal, J. M., Fartoux, L., et al. (2008). Gemcitabine and oxaliplatin in advanced biliary tract carcinoma: a phase II study. *Br J Cancer*. 99, 862-867.
- 86.Wagner, A. D., Buechner-Steudel, P., Moehler, M., et al. (2009). Gemcitabine, oxaliplatin and 5-FU in advanced bile duct and gallbladder carcinoma: two parallel, multicentre phase-II trials. *Br J Cancer*. 101, 1846-1852.
- 87.Zhu, A. X., Meyerhardt, J. A., Blaszkowsky, L. S., et al. (2010). Efficacy and safety of gemcitabine, oxaliplatin, and bevacizumab in advanced biliary-tract cancers and correlation of changes in 18-fluorodeoxyglucose PET with clinical outcome: a phase 2 study. *Lancet Oncol*. 11, 48-54.
- 88.Mizusawa, J., Morizane, C., Okusaka, T., et al. (2016). Randomized Phase III study of gemcitabine plus S-1 versus gemcitabine plus cisplatin in advanced biliary tract cancer: Japan Clinical Oncology Group Study (JCOG1113, FUGA-BT). *Jpn J Clin Oncol*. 46, 385-388.
- 89.Sahai, V., Catalano, P. J., Zalupski, M. M., et al. (2018). Nab-Paclitaxel and Gemcitabine as First-line Treatment of Advanced or Metastatic Cholangiocarcinoma A Phase 2 Clinical Trial. *Jama Oncology*. 4, 1707-1712.
- 90.Kim, S. T., Kang, J. H., Lee, J., et al. (2019). Capecitabine plus oxaliplatin versus gemcitabine plus oxaliplatin as first-line therapy for advanced biliary tract cancers: a

- multicenter, open-label, randomized, phase III, noninferiority trial. *Ann Oncol.* 30, 788-795.
91. Shroff, R. T., Javle, M. M., Xiao, L., et al. (2019). Gemcitabine, Cisplatin, and nab-Paclitaxel for the Treatment of Advanced Biliary Tract Cancers: A Phase 2 Clinical Trial. *JAMA Oncol.* 5, 824-830.
92. Lamarca, A., Palmer, D. H., Wasan, H. S., et al. (2019). ABC-06 | A randomised phase III, multi-centre, open-label study of active symptom control (ASC) alone or ASC with oxaliplatin / 5-FU chemotherapy (ASC+mFOLFOX) for patients (pts) with locally advanced / metastatic biliary tract cancers (ABC) previously-treated with cisplatin/gemcitabine (CisGem) chemotherapy. *J Clin Oncol.* 37, 4003-4003.
93. Edeline, J., Toucheffeu, Y., Guiu, B., et al. (2020). Radioembolization Plus Chemotherapy for First-line Treatment of Locally Advanced Intrahepatic Cholangiocarcinoma: A Phase 2 Clinical Trial. *Jama Oncology.* 6, 51-59.
94. Cercek, A., Boerner, T., Tan, B. R., et al. (2020). Assessment of Hepatic Arterial Infusion of Floxuridine in Combination With Systemic Gemcitabine and Oxaliplatin in Patients With Unresectable Intrahepatic Cholangiocarcinoma: A Phase 2 Clinical Trial. *Jama Oncology.* 6, 60-67.
95. Hayashi, H., Beppu, T., Ishiko, T., et al. (2006). [A 42-month disease free survival case of combined hepatocellular-cholangiocarcinoma with lymph node metastases treated with multimodal therapy]. *Gan to kagaku ryoho. Cancer & chemotherapy.* 33, 1941-1943.
96. Kim, G. M., Jeung, H. C., Kim, D., et al. (2010). A case of combined hepatocellular-cholangiocarcinoma with favorable response to systemic chemotherapy. *Cancer Res Treat.* 42, 235-238.
97. Tani, S., Murata, S., Tamura, M., et al. (2011). [Effectiveness of systemic chemotherapy of GEM+CBDCA+5-FU/LV and hepatic arterial infusion of CDDP in a case of advanced, combined hepatocellular-cholangiocarcinoma with multiple lung metastases]. *Nihon Shokakibyō Gakkai Zasshi.* 108, 1892-1901.
98. Chi, M., Mikhitarian, K., Shi, C., et al. (2012). Management of combined hepatocellular-cholangiocarcinoma: a case report and literature review. *Gastrointest Cancer Res.* 5, 199-202.
99. Fowler, K., Saad, N. E., Brunt, E., et al. (2015). Biphenotypic Primary Liver Carcinomas: Assessing Outcomes of Hepatic Directed Therapy. *Ann Surg Oncol.* 22, 4130-4137.
100. Rogers, J. E., Bolonesi, R. M., Rashid, A., et al. (2017). Systemic therapy for unresectable, mixed hepatocellular-cholangiocarcinoma: treatment of a rare malignancy. *J Gastrointest Oncol.* 8, 347-351.
101. Sano, T., Ishikawa, T., Imai, M., et al. (2018). [Unresectable combined hepatocellular-cholangiocellular carcinoma treated with transcatheter arterial chemoembolization and gemcitabine: a case study]. *Nihon Shokakibyō Gakkai Zasshi.* 115, 1069-1077.
102. Cheng, A.-L., Kang, Y.-K., Chen, Z., et al. (2009). Efficacy and safety of sorafenib in patients in the Asia-Pacific region with advanced hepatocellular

carcinoma: a phase III randomised, double-blind, placebo-controlled trial. *The Lancet Oncology*. 10, 25-34.

103.Llovet, J. M., Ricci, S., Mazzaferro, V., et al. (2008). Sorafenib in Advanced Hepatocellular Carcinoma. *N Engl J Med*. 359, 378-390.

104.Wei, L., Lee, D., Law, C.-T., et al. (2019). Genome-wide CRISPR/Cas9 library screening identified PHGDH as a critical driver for Sorafenib resistance in HCC. *Nature Communications*. 10, 4681.

105.Kudo, M., Finn, R. S., Qin, S., et al. (2018). Lenvatinib versus sorafenib in first-line treatment of patients with unresectable hepatocellular carcinoma: a randomised phase 3 non-inferiority trial. *Lancet*. 391, 1163-1173.

106.Liu, J., Li, X., Zhang, H., et al. (2019). Safety, pharmacokinetics and efficacy of donafenib in treating advanced hepatocellular carcinoma: report from a phase 1b trial. *Pharmazie*. 74, 688-693.

107.Bi, F. (2020) Donafenib versus sorafenib as first-line therapy in advanced hepatocellular carcinoma: An open-label, randomized, multicenter phase II/III trial. (Feng Bi, S. Q. S. G. Y. B. Z. C. Z. W. J. Y. Y. L. Z. M. H. P. P. Y. H. Z. X. C. A. X. X. L. Q., Department of Medical Oncology, W. C. H. S. U. C. C., Pla Cancer Center, N. B. H. N. C., et al. eds.), American Society of Clinical Oncology, ASCO Virtual Scientific Program

108.Bruix, J., Qin, S., Merle, P., et al. (2017). Regorafenib for patients with hepatocellular carcinoma who progressed on sorafenib treatment (RESORCE): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet (London, England)*. 389, 56-66.

109.Yakes, F. M., Chen, J., Tan, J., et al. (2011). Cabozantinib (XL184), a novel MET and VEGFR2 inhibitor, simultaneously suppresses metastasis, angiogenesis, and tumor growth. *Mol Cancer Ther*. 10, 2298-2308.

110.Xiang, Q., Chen, W., Ren, M., et al. (2014). Cabozantinib suppresses tumor growth and metastasis in hepatocellular carcinoma by a dual blockade of VEGFR2 and MET. *Clin Cancer Res*. 20, 2959-2970.

111.Abou-Alfa, G. K., Meyer, T., Cheng, A. L., et al. (2018). Cabozantinib in Patients with Advanced and Progressing Hepatocellular Carcinoma. *N Engl J Med*. 379, 54-63.

112.Spratlin, J. L., Cohen, R. B., Eadens, M., et al. (2010). Phase I pharmacologic and biologic study of ramucirumab (IMC-1121B), a fully human immunoglobulin G1 monoclonal antibody targeting the vascular endothelial growth factor receptor-2. *J Clin Oncol*. 28, 780-787.

113.Kudo, M., Okusaka, T., Motomura, K., et al. (2020). Ramucirumab after prior sorafenib in patients with advanced hepatocellular carcinoma and elevated alpha-fetoprotein: Japanese subgroup analysis of the REACH-2 trial. *J Gastroenterol*. 55, 627-639.

114.Kudo, M., Galle, P. R., Llovet, J. M., et al. (2020). Ramucirumab in elderly patients with hepatocellular carcinoma and elevated alpha-fetoprotein after sorafenib in REACH and REACH-2. *Liver Int*.

- 115.Li, Q., Qin, S., Gu, S., et al. (2020). Apatinib as second-line therapy in Chinese patients with advanced hepatocellular carcinoma: A randomized, placebo-controlled, double-blind, phase III study. *J Clin Oncol.* 38, 4507-4507.
- 116.Kim, R. D., Sarker, D., Meyer, T., et al. (2019). First-in-Human Phase I Study of Fisolatinib (BLU-554) Validates Aberrant FGF19 Signaling as a Driver Event in Hepatocellular Carcinoma. *Cancer Discov.* 9, 1696-1707.
- 117.Kelley, R. K., Gane, E., Assenat, E., et al. (2019). A Phase 2 Study of Galunisertib (TGF- β 1 Receptor Type I Inhibitor) and Sorafenib in Patients With Advanced Hepatocellular Carcinoma. *Clinical And Translational Gastroenterology.* 10, e00056.
- 118.Faivre, S., Santoro, A., Kelley, R. K., et al. (2019). Novel transforming growth factor beta receptor I kinase inhibitor galunisertib (LY2157299) in advanced hepatocellular carcinoma. *Liver Int.* 39, 1468-1477.
- 119.Huang, C.-Y., Hsieh, F.-S., Wang, C.-Y., et al. (2018). Palbociclib enhances radiosensitivity of hepatocellular carcinoma and cholangiocarcinoma via inhibiting ataxia telangiectasia–mutated kinase–mediated DNA damage response. *Eur J Cancer.* 102, 10-22.
- 120.Reiter, F. P., Denk, G., Ziesch, A., et al. (2019). Predictors of ribociclib-mediated antitumour effects in native and sorafenib-resistant human hepatocellular carcinoma cells. *Cell Oncol (Dordr).* 42, 705-715.
- 121.Bouattour, M., Raymond, E., Qin, S., et al. (2018). Recent developments of c-Met as a therapeutic target in hepatocellular carcinoma. *Hepatology.* 67, 1132-1149.
- 122.Rimassa, L., Assenat, E., Peck-Radosavljevic, M., et al. (2018). Tivantinib for second-line treatment of MET-high, advanced hepatocellular carcinoma (METIV-HCC): a final analysis of a phase 3, randomised, placebo-controlled study. *Lancet Oncol.* 19, 682-693.
- 123.Farshidfar, F., Zheng, S., Gingras, M. C., et al. (2017). Integrative Genomic Analysis of Cholangiocarcinoma Identifies Distinct IDH-Mutant Molecular Profiles. *Cell Rep.* 18, 2780-2794.
- 124.Waitkus, M. S., Diplas, B. H., Yan, H. (2018). Biological Role and Therapeutic Potential of IDH Mutations in Cancer. *Cancer Cell.* 34, 186-195.
- 125.Yang, T., Liang, L., Wang, M. D., et al. (2020). FGFR inhibitors for advanced cholangiocarcinoma. *Lancet Oncol.* 21, 610-612.
- 126.Roskoski, R. (2020). The role of fibroblast growth factor receptor (FGFR) protein-tyrosine kinase inhibitors in the treatment of cancers including those of the urinary bladder. *Pharmacol Res.* 151, 104567.
- 127.Wu, Y. M., Su, F., Kalyana-Sundaram, S., et al. (2013). Identification of targetable FGFR gene fusions in diverse cancers. *Cancer Discov.* 3, 636-647.
- 128.Arai, Y., Totoki, Y., Hosoda, F., et al. (2014). Fibroblast growth factor receptor 2 tyrosine kinase fusions define a unique molecular subtype of cholangiocarcinoma. *Hepatology.* 59, 1427-1434.
- 129.Liu, P. C. C., Wu, L. X., Koblisch, H., et al. (2015). Preclinical characterization of the selective FGFR inhibitor INCB054828. *Cancer Res.* 75

130. Abou-Alfa, G. K., Sahai, V., Hollebecque, A., et al. (2020). Pemigatinib for previously treated, locally advanced or metastatic cholangiocarcinoma: a multicentre, open-label, phase 2 study. *Lancet Oncol.* 21, 671-684.
131. Javle, M. M., Shroff, R. T., Zhu, A., et al. (2016). A phase 2 study of BGJ398 in patients (pts) with advanced or metastatic FGFR-altered cholangiocarcinoma (CCA) who failed or are intolerant to platinum-based chemotherapy. *J Clin Oncol.* 34, 335-335.
132. Ochiwa, H., Fujita, H., Itoh, K., et al. (2013). Abstract A270: TAS-120, a highly potent and selective irreversible FGFR inhibitor, is effective in tumors harboring various FGFR gene abnormalities. *Mol Cancer Ther.* 12, A270-A270.
133. Meric-Bernstam, F., Arkenau, H., Tran, B., et al. (2018). Efficacy of TAS-120, an irreversible fibroblast growth factor receptor (FGFR) inhibitor, in cholangiocarcinoma patients with FGFR pathway alterations who were previously treated with chemotherapy and other FGFR inhibitors. *Ann Oncol.* 29 Suppl 5, v100.
134. Goyal, L., Bahleda, R., Furuse, J., et al. (2019). FOENIX-101: A phase II trial of TAS-120 in patients with intrahepatic cholangiocarcinoma harboring FGFR2 gene rearrangements. *J Clin Oncol.* 37, TPS468-TPS468.
135. Goyal, L., Shi, L., Liu, L. Y., et al. (2019). TAS-120 Overcomes Resistance to ATP-Competitive FGFR Inhibitors in Patients with FGFR2 Fusion-Positive Intrahepatic Cholangiocarcinoma. *Cancer Discov.* 9, 1064-1079.
136. Papadopoulos, K. P., El-Rayes, B. F., Tolcher, A. W., et al. (2017). A Phase 1 study of ARQ 087, an oral pan-FGFR inhibitor in patients with advanced solid tumours. *Br J Cancer.* 117, 1592-1599.
137. Gourd, E. (2019). Derazantinib for intrahepatic cholangiocarcinoma. *Lancet Oncol.* 20, e11.
138. Mazzaferro, V., El-Rayes, B. F., Droz Dit Busset, M., et al. (2019). Derazantinib (ARQ 087) in advanced or inoperable FGFR2 gene fusion-positive intrahepatic cholangiocarcinoma. *Br J Cancer.* 120, 165-171.
139. Bahleda, R., Italiano, A., Hierro, C., et al. (2019). Multicenter Phase I Study of Erdafitinib (JNJ-42756493), Oral Pan-Fibroblast Growth Factor Receptor Inhibitor, in Patients with Advanced or Refractory Solid Tumors. *Clin Cancer Res.* 25, 4888-4897.
140. Park, J. O., Feng, Y.-H., Chen, Y.-Y., et al. (2019). Updated results of a phase IIa study to evaluate the clinical efficacy and safety of erdafitinib in Asian advanced cholangiocarcinoma (CCA) patients with FGFR alterations. *J Clin Oncol.* 37, 4117-4117.
141. Moeini, A., Sia, D., Bardeesy, N., et al. (2016). Molecular Pathogenesis and Targeted Therapies for Intrahepatic Cholangiocarcinoma. *Clin Cancer Res.* 22, 291-300.
142. Krook, M. A., Lenyo, A., Wilberding, M., et al. (2020). Efficacy of FGFR Inhibitors and Combination Therapies for Acquired Resistance in FGFR2-Fusion Cholangiocarcinoma. *Mol Cancer Ther.* 19, 847-857.
143. Lowery, M. A., Abou-Alfa, G. K., Valle, J. W., et al. (2017). ClarIDHy: A phase 3, multicenter, randomized, double-blind study of AG-120 vs placebo in patients with an advanced cholangiocarcinoma with an IDH1 mutation. *J Clin Oncol.* 35

- 144.(2019). Abou-Alfa GK et al. ESMO 2019 plenary session.
- 145.Simone, V., Brunetti, O., Lupo, L., et al. (2017). Targeting Angiogenesis in Biliary Tract Cancers: An Open Option. *Int J Mol Sci.* 18
- 146.Mehta, R. J., Kim, D. W., Soares, H. P., et al. (2018). Phase II study of copanlisib (BAY 80-6946) in combination with gemcitabine and cisplatin in advanced cholangiocarcinoma. *J Clin Oncol.* 36, TPS525-TPS525.
- 147.Kim, J. W., Lee, K.-H., Kim, J.-W., et al. (2018). Phase Ib study of binimetinib (MEK162) in combination with capecitabine in gemcitabine-pretreated advanced biliary tract cancer. *J Clin Oncol.* 36, 4079-4079.
- 148.Futsukaichi, Y., Tajiri, K., Kobayashi, S., et al. (2019). Combined hepatocellular-cholangiocarcinoma successfully treated with sorafenib: case report and review of the literature. *Clin J Gastroenterol.* 12, 128-134.
- 149.He, X., Xu, C. (2020). Immune checkpoint signaling and cancer immunotherapy. *Cell Res.*
- 150.Gandini, S., Massi, D., Mandalà, M. (2016). PD-L1 expression in cancer patients receiving anti PD-1/PD-L1 antibodies: A systematic review and meta-analysis. *Critical Reviews in Oncology/Hematology.* 100, 88-98.
- 151.Seidel, J. A., Otsuka, A., Kabashima, K. (2018). Anti-PD-1 and Anti-CTLA-4 Therapies in Cancer: Mechanisms of Action, Efficacy, and Limitations. *Front Oncol.* 8, 86.
- 152.El-Khoueiry, A. B., Sangro, B., Yau, T., et al. (2017). Nivolumab in patients with advanced hepatocellular carcinoma (CheckMate 040): an open-label, non-comparative, phase 1/2 dose escalation and expansion trial. *Lancet.* 389, 2492-2502.
- 153.Zhu, A. X., Finn, R. S., Edeline, J., et al. (2018). Pembrolizumab in patients with advanced hepatocellular carcinoma previously treated with sorafenib (KEYNOTE-224): a non-randomised, open-label phase 2 trial. *The Lancet Oncology.* 19, 940-952.
- 154.Finn, R. S., Ryoo, B.-Y., Merle, P., et al. (2020). Pembrolizumab As Second-Line Therapy in Patients With Advanced Hepatocellular Carcinoma in KEYNOTE-240: A Randomized, Double-Blind, Phase III Trial. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology.* 38, 193-202.
- 155.Ikeda, M., Sung, M. W., Kudo, M., et al. (2018). A phase 1b trial of lenvatinib (LEN) plus pembrolizumab (PEM) in patients (pts) with unresectable hepatocellular carcinoma (uHCC). *J Clin Oncol.* 36, 4076-4076.
- 156.Qin, S., Ren, Z., Meng, Z., et al. (2020). Camrelizumab in patients with previously treated advanced hepatocellular carcinoma: a multicentre, open-label, parallel-group, randomised, phase 2 trial. *Lancet Oncol.* 21, 571-580.
- 157.Hoy, S. M. (2019). Sintilimab: First Global Approval. *Drugs.* 79, 341-346.
- 158.Zhang, J., Wu, L., Liu, J., et al. (2020). A metastatic intrahepatic cholangiocarcinoma treated with programmed cell death 1 inhibitor: a case report and literature review. *Immunotherapy.* 12, 555-561.

- 159.Zhang, T., Song, X., Xu, L., et al. (2018). The binding of an anti-PD-1 antibody to FcγRI has a profound impact on its biological functions. *Cancer immunology, immunotherapy : CII.* 67, 1079-1090.
- 160.Dahan, R., Sega, E., Engelhardt, J., et al. (2015). FcγR3 Modulate the Anti-tumor Activity of Antibodies Targeting the PD-1/PD-L1 Axis. *Cancer Cell.* 28, 543.
- 161.Deva, S., Lee, J., Lin, C., et al. (2018). 700 A phase Ia/Ib trial of tislelizumab, an anti-PD-1 antibody (ab), in patients (pts) with advanced solid tumors. 29, mdy487. 042.
- 162.Qin, S., Finn, R. S., Kudo, M., et al. (2019). RATIONALE 301 study: tislelizumab versus sorafenib as first-line treatment for unresectable hepatocellular carcinoma. *Future oncology (London, England).* 15, 1811-1822.
- 163.Keam, S. J. (2019). Toripalimab: First Global Approval. *Drugs.* 79, 573-578.
- 164.Stein, S., Pishvaian, M. J., Lee, M. S., et al. (2018). Safety and clinical activity of 1L atezolizumab + bevacizumab in a phase Ib study in hepatocellular carcinoma (HCC). *J Clin Oncol.* 36, 4074-4074.
- 165.Finn, R. S., Qin, S., Ikeda, M., et al. (2020). Atezolizumab plus Bevacizumab in Unresectable Hepatocellular Carcinoma. *N Engl J Med.* 382, 1894-1905.
- 166.Korman, A. J., Peggs, K. S., Allison, J. P. (2006). Checkpoint blockade in cancer immunotherapy. *Advances in immunology.* 90, 297-339.
- 167.Wei, S. C., Duffy, C. R., Allison, J. P. (2018). Fundamental Mechanisms of Immune Checkpoint Blockade Therapy. *Cancer Discov.* 8, 1069-1086.
- 168.Furuse, J., Chan, S. L., Sangro, B., et al. (2018). A phase 3 study of durvalumab tremelimumab as first-line treatment in patients with unresectable hepatocellular carcinoma: himalaya study. *Liver Cancer.* 7, 171-.
- 169.Floudas, C. S., Xie, C., Brar, G., et al. (2019). Combined immune checkpoint inhibition (ICI) with tremelimumab and durvalumab in patients with advanced hepatocellular carcinoma (HCC) or biliary tract carcinomas (BTC). *J Clin Oncol.* 37, 336-336.
- 170.Schumacher, T. N., Schreiber, R. D. (2015). Neoantigens in cancer immunotherapy. *Science.* 348, 69-74.
- 171.Yarchoan, M., Johnson, B. A., 3rd, Lutz, E. R., et al. (2017). Targeting neoantigens to augment antitumour immunity. *Nat Rev Cancer.* 17, 209-222.
- 172.Desrichard, A., Snyder, A., Chan, T. A. (2016). Cancer Neoantigens and Applications for Immunotherapy. *Clin Cancer Res.* 22, 807-812.
- 173.Mardis, E. R. (2019). Neoantigens and genome instability: impact on immunogenomic phenotypes and immunotherapy response. *Genome Med.* 11, 71.
- 174.Lu, L., Jiang, J., Zhan, M., et al. (2020). Targeting Neoantigens in Hepatocellular Carcinoma for Immunotherapy: A Futile Strategy? *Hepatology.*
- 175.Shi, D., Shi, Y., Kaseb, A. O., et al. (2020). Chimeric Antigen Receptor-Glypican-3 T-Cell Therapy for Advanced Hepatocellular Carcinoma: Results of Phase 1 Trials. *Clin Cancer Res.*
- 176.Vormehr, M., Tureci, O., Sahin, U. (2019). Harnessing Tumor Mutations for Truly Individualized Cancer Vaccines. *Annu Rev Med.* 70, 395-407.

177. Wang, Y., Yang, X., Yu, Y., et al. (2018). Immunotherapy of patient with hepatocellular carcinoma using cytotoxic T lymphocytes ex vivo activated with tumor antigen-pulsed dendritic cells. *J Cancer*. 9, 275-287.
178. Chen, C., Ma, Y. H., Zhang, Y. T., et al. (2018). Effect of dendritic cell-based immunotherapy on hepatocellular carcinoma: A systematic review and meta-analysis. *Cytotherapy*. 20, 975-989.
179. Sahin, U., Derhovanessian, E., Miller, M., et al. (2017). Personalized RNA mutanome vaccines mobilize poly-specific therapeutic immunity against cancer. *Nature*. 547, 222-226.
180. Ott, P. A., Hu, Z., Keskin, D. B., et al. (2017). An immunogenic personal neoantigen vaccine for patients with melanoma. *Nature*. 547, 217-221.
181. Wculek, S. K., Cueto, F. J., Mujal, A. M., et al. (2020). Dendritic cells in cancer immunology and immunotherapy. *Nat Rev Immunol*. 20, 7-24.
182. Le, D. T., Durham, J. N., Smith, K. N., et al. (2017). Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science*. 357, 409-413.
183. Sidaway, P. (2020). MSI-H: a truly agnostic biomarker? *Nat Rev Clin Oncol*. 17, 68.
184. Rashid, A., Ueki, T., Gao, Y. T., et al. (2002). K-ras mutation, p53 overexpression, and microsatellite instability in biliary tract cancers: a population-based study in China. *Clin Cancer Res*. 8, 3156-3163.
185. Silva, V. W., Askan, G., Daniel, T. D., et al. (2016). Biliary carcinomas: pathology and the role of DNA mismatch repair deficiency. *Chin Clin Oncol*. 5, 62.
186. Rizvi, S., Khan, S. A., Hallemeier, C. L., et al. (2018). Cholangiocarcinoma - evolving concepts and therapeutic strategies. *Nat Rev Clin Oncol*. 15, 95-111.
187. Marcus, L., Lemery, S. J., Keegan, P., et al. (2019). FDA Approval Summary: Pembrolizumab for the Treatment of Microsatellite Instability-High Solid Tumors. *Clin Cancer Res*. 25, 3753-3758.
188. Bang, Y.-J., Ueno, M., Malka, D., et al. (2019). Pembrolizumab (pembro) for advanced biliary adenocarcinoma: Results from the KEYNOTE-028 (KN028) and KEYNOTE-158 (KN158) basket studies. *J Clin Oncol*. 37, 4079-4079.
189. Ueno, M., Ikeda, M., Morizane, C., et al. (2019). Nivolumab alone or in combination with cisplatin plus gemcitabine in Japanese patients with unresectable or recurrent biliary tract cancer: a non-randomised, multicentre, open-label, phase 1 study. *Lancet Gastroenterol Hepatol*. 4, 611-621.
190. Kim, R. D., Kim, D. W., Alese, O. B., et al. (2019). A phase II study of nivolumab in patients with advanced refractory biliary tract cancers (BTC). *J Clin Oncol*. 37, 4097-4097.
191. Ioka, T., Ueno, M., Oh, D.-Y., et al. (2019). Evaluation of safety and tolerability of durvalumab (D) with or without tremelimumab (T) in patients (pts) with biliary tract cancer (BTC). *J Clin Oncol*. 37, 387-387.
192. Oh, D. Y., Chen, L. T., He, A. R., et al. (2019). A phase III, randomized, double-blind, placebo-controlled, international study of durvalumab in combination with gemcitabine plus cisplatin for patients with advanced biliary tract cancers: TOPAZ-1. *Ann Oncol*. 30, v319.

- 193.Liu, J., Dang, H., Wang, X. W. (2018). The significance of intertumor and intratumor heterogeneity in liver cancer. *Exp Mol Med.* 50, e416.
- 194.Losic, B., Craig, A. J., Villacorta-Martin, C., et al. (2020). Intratumoral heterogeneity and clonal evolution in liver cancer. *Nat Commun.* 11, 291.
- 195.Tang, J., Sui, C. J., Wang, D. F., et al. (2020). Targeted sequencing reveals the mutational landscape responsible for sorafenib therapy in advanced hepatocellular carcinoma. *Theranostics.* 10, 5384-5397.
- 196.Ding, X., He, M., Chan, A. W. H., et al. (2019). Genomic and Epigenomic Features of Primary and Recurrent Hepatocellular Carcinomas. *Gastroenterology.* 157, 1630-1645 e1636.
- 197.Ringelhan, M., Pfister, D., O'Connor, T., et al. (2018). The immunology of hepatocellular carcinoma. *Nat Immunol.* 19, 222-232.
- 198.Rizvi, S., Wang, J., El-Khoueiry, A. B. (2020). Liver Cancer Immunity. *Hepatology.*
- 199.Shen, Y., Wei, Y., Wang, Z., et al. (2015). TGF-beta regulates hepatocellular carcinoma progression by inducing Treg cell polarization. *Cell Physiol Biochem.* 35, 1623-1632.
- 200.Zhai, W., Zhou, X., Wang, H., et al. (2020). A novel cyclic peptide targeting LAG-3 for cancer immunotherapy by activating antigen-specific CD8(+) T cell responses. *Acta Pharm Sin B.* 10, 1047-1060.
- 201.Wang, C., Vegna, S., Jin, H., et al. (2019). Inducing and exploiting vulnerabilities for the treatment of liver cancer. *Nature.* 574, 268-272.

Figure1 Barcelona Clinic Liver Cancer staging system and corresponding treatment options.

The schematic diagram illustrates therapeutic choice by which a treatment theoretically recommended for a different stage as best treatment option. 1L, first-line; 2L, second-line; ECOG, Eastern Cooperative Oncology Group; M, metastasis stage; N, nodal stage; PEI, percutaneous ethanol injection; PS, performance status; RFA, radiofrequency ablation; T, tumor stage; TACE, transarterial chemoembolization; TARE, Transarterialradioembolization Y-90, Y-90 Radioembolization

Figure2 Treatment strategy for advanced HCC and ICC.

The schematic illustration represents FDA approved drugs for treatment of advanced HCC and ICC. First line drugs for HCC includes Sorafenib, Lenvatinib, Atezolizumab plus Bevacizumab, Tremelimumab plus Durvalumab and Donafenib, whereas for ICC,

the combination of gemcitabine and cisplatin is currently proposed as first-line. The bottom row represents corresponding second line therapies which come in when patients are not suitable for their first line therapy.

Journal Pre-proof

Table1. Systemic therapies currently or promising approved for advanced HCC and ICC

Drugs	Target	Therapy Line	Approved Year	Trial
HCC				
Sorafenib (Nexavar)	VEGFR-2, VEGFR-3, PDGFR- β ,RAF kinases	1	2007	SHARP Asian-Pacific
Lenvatinib (Lenvima)	FGFR, VEGFR, PDGFR- α , RET,KIT	1	2018	REFLECT
Regorafenib (Stivarga)	Tie2, VEGFR, PDGFR, FGFR	2	2017	RESORCE
Nivolumab (Opdivo)	PD1	2	2017	CHECKMATE-040
Cabozantinib (Cabometyx)	c-Met, VEGFR-2, AXL, RET	2	2018	CELESTIAL
Pembrolizumab (Keytruda)	PD1	2	2018	KEYNOTE-224
Ramucirumab (CYRAMZA)	VEGFR-2	2	2019	REACH-2
Nivolumab plus ipilimumab (Opdivo plus Yervoy)	PD1,CTLA4	2	2020	Cohort 4 of CHECKMATE-040
Atezolizumab plus Bevacizumab	PD-L1,VEGF	1	Promising	IMbrave150
Tremelimumab plus Durvalumab	PD1,CTLA4	1	Promising	NCT02519348
Donafenib	VEGFR, BRAF	1	Promising	NCT02645981
Apatinib	VEGFR-2	2	Promising	NCT02329860
ICC				
Gemcitabine plus cisplatin	Chemotherapy	1	2010	ABC-02
Pemigatinib (Pemazyre)	FGFR1-3	2	2020	FIGHT-202
Ivosidenib	IDH-1/2	2	Promising	ClarIDHy

Table2. Selected ongoing systemic therapy clinical trials for advanced HCC

Drug	Target	Sponsor	Status	Phase	Enrollment	Trial Identifier
Targeted therapy						
Cabozantinib	VEGFR	Hospices Civils de Lyon	Recruiting	Phase 4	170	NCT03963206
Lenvatinib	VEGFR	Eisai Pharmaceuticals India Pvt. Ltd	Not yet recruiting	Phase 4	50	NCT04297254
Donafenib	VEGFR	Suzhou Zelgen Biopharmaceuticals	Completed	Phase 2 Phase 3	668	NCT02645981
Milciclib	CDK2	Tiziana LifeSciences	Active, not recruiting	Phase 2	31	NCT03109886
Palbociclib	CDK4/6	Pfizer	Active, not recruiting	Phase 2	23	NCT01356628
Ribociclib	CDK4/6	Texas University	Recruiting	Phase 2	40	NCT02524119
Galunisertib vs LY2157299 + Sorafenib vs Placebo + Sorafenib	TGF- β	Eli Lilly	Active, not recruiting	Phase 2	120	NCT02178358
Immunotherapy						
Tislelizumab vs Sorafenib	PD-1	BeiGene	Active, not recruiting	Phase 3	674	NCT03412773
Toripalimab vs Placebo	PD-1	Shanghai Junshi Bioscience	Recruiting	Phase 2 Phase 3	402	NCT03859128
Nivolumab vs Placebo	PD-1	Bristol-Myers Squibb	Recruiting	Phase 3	530	NCT03383458
Nivolumab vs Sorafenib	PD-1	Bristol-Myers Squibb	Active, not recruiting	Phase 3	1723	NCT02576509
Pembrolizumab vs Placebo	PD-1	Merck Sharp & Dohme	Recruiting	Phase 3	950	NCT03867084
Avelumab	PD-L1	Seoul National University Hospital	Active, not recruiting	Phase 2	30	NCT03389126
Combined therapy						
Lenvatinib + Pembrolizumab vs Lenvatinib + Placebo	VGFR,PD-1	Merck Sharp & Dohme	Active, not recruiting	Phase 3	750	NCT03713593
CS1003+Lenvatinib vs Placebo+Lenvatinib	VGFR,PD-1	CStone Pharmaceuticals	Recruiting	Phase 3	525	NCT04194775
Tislelizumab + Regorafenib vs Placebo + Regorafenib	VEGF,PD-1	National Taiwan University Hospital	Not yet recruiting	Phase 2	125	NCT04183088
Toripalimab + Lenvatinib	VGFR,PD-1	Peking Union Medical College Hospital	Not yet recruiting	Phase 2	76	NCT04368078
Durvalumab + Bevacizumab vs Placebo	VEGF,PD-L1	AstraZeneca	Recruiting	Phase 3	888	NCT03847428
Atezolizumab + Bevacizumab vs Sorafenib	VEGF,PD-L1	Hoffmann-La Roche	Recruiting	Phase 3	480	NCT03434379
Atezolizumab + Bevacizumab	VEGF,PD-L1	National Health Research Institutes, Taiwan	Not yet recruiting	Phase 2	48	NCT04180072
cabozantinib + Atezolizumab vs sorafenib	VEGF,PD-L1	Exelixis	Recruiting	Phase 3	740	NCT03755791

Drug	Target	Sponsor	Status	Phase	Enrollment	Trial Identifier
Atezolizumab + Bevacizumab vs Active Surveillance	VEGF,PD-L1	Hoffmann-La Roche	Recruiting	Phase 3	662	NCT04102098
Regorafenib + Nivolumab	VEGF,PD-1	Fundacion Clinic per a la Recerca Biomédica	Not yet recruiting	Phase 1 Phase 2	60	NCT04170556
Lenvatinib + Pembrolizumab vs Lenvatinib + Placebo	VEGFR,PD-1	Merck Sharp & Dohme	Recruiting	Phase 3	750	NCT03713593
Camrelizumab + Apatinib	VEGFR,PD-1	Zhejiang University	Recruiting	Phase 1 Phase 2	120	NCT04035876
Camrelizumab + Apatinib vs Sorafenib	VEGFR,PD-1	Jiangsu HengRui	Recruiting	Phase 3	510	NCT03764293
Sintilimab + Lenvatinib	VEGFR,PD-1	Beijing Cancer Hospital	Not yet recruiting	Phase 2	56	NCT04042805
Sintilimab + IBI305 vs Sorafenib	VEGF,PD-1	Innovent Biologics	Recruiting	Phase 2 Phase 3	566	NCT03794440
Regorafenib + Avelumab	VEGF,PD-L1	Institut Bergonié	Recruiting	Phase 1 Phase 2	362	NCT03475953
Sorafenib + Toripalimab	VEGF,PD-1	Sichuan University	Not yet recruiting	Phase 1 Phase 2	39	NCT04069949
Galunisertib + Nivolumab	TGF-beta,PD-1	Eli Lilly	Active, not recruiting	Phase 2	75	NCT02423343
Fisogatinib + CS1001	FGFR4,PD-L1	CStone Pharmaceuticals	Recruiting	Phase 1 Phase 2	52	NCT04194801
AK105 + Anlotinib vs Sorafenib	RTK,PD-1	Chia Tai Tianqing	Not yet recruiting	Phase 3	648	NCT04344158
Anlotinib + Sintilimab	RTK,PD-1	Nanjing Medical University First hospital	Recruiting	Phase 2	20	NCT04052152
Abemaciclib + Nivolumab	CDK4/6,PD-1	Abramson Cancer Center Pennsylvania University	Suspended (COVID-19)	Phase 2	27	NCT03781960
Durvalumab + Tremelimumab vs Durvalumab vs Sorafenib	PD-L1,CTLA-4	AstraZeneca	Active, not recruiting	Phase 3	1310	NCT03298451
Nivolumab + Ipilimumab vs Sorafenib/lenvatinib	PD-1,CTLA-4	Bristol-Myers Squibb	Recruiting	Phase 3	1084	NCT04039607
Durvalumab + Tremelimumab vs Durvalumab monotherapy vs Tremelimumab monotherapy vs Durvalumab + Bevacizumab	VEGF,PD-L1,CTLA-4	MedImmune LLC	Active, not recruiting	Phase 2	433	NCT02519348
Galunisertib vs Galunisertib +Sorafenib/Ramucirumab	TGF-β,VEGF,VEGFR	Eli Lilly	Active, not recruiting	Phase 2	193	NCT01246986
Lenvatinib + Pembrolizumab + TACE vs Placebo + TACE	VEGFR,PD-1,TACE	Merck Sharp & Dohme	Not yet recruiting	Phase 3	950	NCT04246177
TAI+ lenvatinib vs Lenvatinib	VEGFR,Chemoinfusion	Sun Yat-sen University	Recruiting	Phase 3	206	NCT04053985
SBRT +Sintilimab vs SBRT	PD-1,Radiation	Mian XI, Sun Yat-sen University	Recruiting	Phase 2 Phase 3	116	NCT04167293
Donafenib + Anti-PD-1 antibody	VEGFR, PD-L1	Zhejiang University	Recruiting	Phase 1	30	NCT04418401
Others						

Drug	Target	Sponsor	Status	Phase	Enrollment	Trial Identifier
ALT-803 + Avelumab	PD-L1,IL-15 superagonist	Altor BioScience	Recruiting	Phase 2	611	NCT03228667
KY1044 monotherapy vs KY1044 + Atezolizumab	PD-L1,T cell CO-stimulator	Kymab Limited	Recruiting	Phase 1 Phase 2	412	NCT03829501
Pexa-Vec + Nivolumab	PD-1,oncolyticvirus	Transgene	Active, not recruiting	Phase 1 Phase 2	30	NCT03071094
Nivolumab + BMS-986253 vs Nivolumab + Cabiralizumab vs Nivolumab Monotherapy	VEGFR2,PD-1, interleukin-8	NYU Langone Health	Not yet recruiting	Phase 2	74	NCT04050462
Entecavir /Tenofovir Disoproxil monotherapy	Antiviral therapy(HBV)	West China Hospital	Recruiting	Phase 4	450	NCT04032860
Vemlidy vs Placebo	Antiviral therapy(HBV)	Taipei Veterans General Hospital, Taiwan	Not yet recruiting	Phase 4	402	NCT04290936
Pexastimogene Devacirepvec vs Sorafenib	Vaccinia virus-based oncolytic immunotherapy	SillaJen, Inc.	Active, not recruiting	Phase 3	600	NCT02562755
Immune cell						
CD147-CART	CAR-T therapy	Xijing Hospital	Recruiting	Phase 1	34	NCT03993743
Anti-DR5 CAR-T/TCR-T cells immunotherapy	CAR-T therapy	Shenzhen BinDeBio Ltd.	Recruiting	Phase 1 Phase 2	50	NCT03941626
CAR-GPC3 T Cells	CAR-T therapy	Zhejiang University	Recruiting	Phase 1	36	NCT03980288
GPC3 or TGFβ targeting CAR-T cell therapy	CAR-T therapy	Guangzhou Medical University Second Hospital	Recruiting	Phase 1	30	NCT03198546
GPC3-CAR (GLYCAR T cells) + Fludarabine and Cytosan	CAR-T therapy	Baylor College of Medicine	Recruiting	Phase 1	14	NCT02905188
c-Met/PD-L1 CAR-T cell injection	CAR-T therapy	Second Hospital Nanjing Medical University	Not yet recruiting	Early Phase 1	50	NCT03672305
IMA202 Product	TCR-T therapy	Immatics US, Inc.	Recruiting	Phase 1	16	NCT03441100
Microwave Ablation + Neoantigen Vaccines	Neoantigen DC Vaccines	Chinese PLA General Hospital	Recruiting	Phase 1	24	NCT03674073
DC vaccines	Neoantigen DC Vaccines	Sichuan University	Recruiting	Phase 1	80	NCT04147078
Autologous DC+ conjugate vaccine	Vaccine	Mayo Clinic	Recruiting	Early Phase 1	26	NCT03942328

TAI = Transarterial chemoinfusion; SBRT = stereotactic body radiotherapy


Table3. Selected ongoing systemic therapy clinical trials for advanced CCA

Drug	Target	Sponsor	Status	Condition or disease	Phase	Enrolment	Trial Identifier
Chemotherapy							
FOLFIRINOX vs GEMOX	Chemotherapy	Shi Ming, Sun Yat-sen University	Recruiting	Intrahepatic Cholangiocarcinoma	Phase 3	188	NCT03771846
Anlotinib vs Anlotinib +Levamisole	Chemotherapy	Zhengzhou University First hospital	Recruiting	Intrahepatic Cholangiocarcinoma	Phase 3	152	NCT03940378
Melphalan/PHP vs CisGem	Chemotherapy	Delcath Systems Inc.	Recruiting	Intrahepatic Cholangiocarcinoma	Phase 2 Phase 3	295	NCT03086993
Gemcitabine + Capecitabine vs Capecitabine	Chemotherapy	Tianjin Medical University	Recruiting	Cholangiocarcinoma	Phase 3	460	NCT03779035
Nab-paclitaxel, Cisplatin, Gemcitabine vs CisGem	Chemotherapy	Southwest Oncology Group	Recruiting	Intrahepatic Cholangiocarcinoma	Phase 3	268	NCT03768414
CisGem vs Capecitabine	Chemotherapy	niversitätsklinikum Hamburg-Eppendorf	Recruiting	Cholangiocarcinoma Gall Bladder Carcinoma	Phase 3	781	NCT02170090
Targeted therapy							
BGJ398 (Infigratinib)	FGFR2	QED Therapeutics, Inc.	Recruiting	Cholangiocarcinoma FGFR2 Gene Mutation	Phase 2	160	NCT02150967
Infigratinib vs CisGem	FGFR2	QED Therapeutics, Inc.	Recruiting	Cholangiocarcinoma FGFR2 Gene Mutation	Phase 3	384	NCT03773302
Pemigatinib vs CisGem	FGFR2	Incyte Corporation	Recruiting	Unresectable Cholangiocarcinoma	Phase 3	432	NCT03656536
Derazantinib	FGFR2	Basilea Pharmaceutica	Recruiting	Intrahepatic Cholangiocarcinoma	Phase 2	143	NCT03230318
TAS-120 vs CisGem	FGFR2	Taiho Oncology, Inc.	Not yet recruiting	Cholangiocarcinoma	Phase 3	216	NCT04093362
Ponatinib	FGFR, VEGFR	Sameek Roychowdhury	Recruiting	Solid Tumor with FGFR Mutations	Phase 2	45	NCT02272998
AG-120 vs placebo	IDH1	Agios Pharmaceuticals	Active, not recruiting	Advanced Cholangiocarcinoma	Phase 3	186	NCT02989857
FT-2102	IDH1	Forma Therapeutics, Inc.	Recruiting	Solid Tumors including ICC	Phase 1 Phase 2	200	NCT03684811
BAY 1436032	IDH1	Bayer	Active, not recruiting	Solid Tumors including ICC	Phase 1	81	NCT02746081
Ramucirumab	VEGFR2	M.D. Anderson Cancer Center	Recruiting	Cholangiocarcinoma	Phase 2	50	NCT02520141
Apatinib	VEGFR2	Zhengzhou University First Hospital	Recruiting	Intrahepatic Cholangiocarcinoma	Phase 2	30	NCT03521219
Surufatinib vs Capecitabine	VEGFR	Hutchison Medipharma	Recruiting	Biliary Tract Cancer	Phase 2 Phase 3	298	NCT03873532
Niraparib	PARP	University of Florida	Recruiting	Cholangiocarcinoma	Phase 2	57	NCT03207347
Olaparib	PARP	Academic and Community Cancer Research United	Not yet recruiting	Biliary tract cancer	Phase 2	36	NCT04042831

Drug	Target	Sponsor	Status	Condition or disease	Phase	Enrollment	Trial Identifier
CB-103	NOTCH	Cellestia Biotech AG	Recruiting	Cholangiocellular Carcinoma	Phase 1 Phase 2	165	NCT03422679
CX-2009	CD166	CytomX Therapeutics	Recruiting	Cholangiocarcinoma	Phase 1 Phase 2	150	NCT03149549
Bortezomib vs supportive care	Proteasome inhibitor	Zhengang Yuan	Recruiting	Intrahepatic Cholangiocarcinoma	Phase 3	50	NCT03345303
Immunotherapy							
Pembrolizumab + CisGem vs Placebo + CisGem	PD-1	Merck Sharp & Dohme	Recruiting	Biliary Tract Carcinoma	Phase 3	788	NCT04003636
Durvalumab + CisGem vs Placebo + CisGem	PD-L1	AstraZeneca	Recruiting	Biliary Tract Neoplasms	Phase 3	474	NCT03875235
Pembrolizumab	PD-1	Samsung Medical Center	Recruiting	biliary tract cancer	Phase 2	33	NCT03110328
Durvalumab + CisGem vs Placebo + CisGem	PD-L1	AstraZeneca	Recruiting	Biliary Tract Neoplasms	Phase 3	474	NCT03875235
Combined therapies							
Systemic Chemotherapy vs Chemotherapy and radiation	Chemotherapy, radiation	Tata Memorial Hospital	Recruiting	Cholangiocarcinoma	Phase 3	155	NCT02773485
CisGem + pembrolizumab	Chemotherapy, PD-1	EORTC	Recruiting	Biliary Tract Cancer	Phase 2	50	NCT03260712
Camrelizumab + Apatinib vs Camrelizumab + FOLFOX4 or GEMOX	PD-1, VEGF, chemotherapy	Jiangsu HengRui Medicine	Recruiting	Advanced Biliary Tract Carcinoma	Phase 2	152	NCT03092895
Lenvatinib + Pembrolizumab	PD-1, VEGF	Peking Union Medical College Hospital	Recruiting	Cholangiocarcinoma	Phase 2	50	NCT03895970
Bintrafusp alfa + CisGem vs Placebo + CisGem	PD-L1xTGF- β	EMD Serono Research & Development Institute	Recruiting	Biliary Tract Cancer Cholangiocarcinoma	Phase 2 Phase 3	512	NCT04066491
XmAb22841, XmAb22841 + Pembrolizumab	PD-1, CTLA-4 x LAG-3	Xencor, Inc.	Recruiting	Advanced Solid Tumors including ICC	Phase 1	242	NCT03849469
Rucaparib + Nivolumab	PD-1, PARP	University of Michigan Rogel Cancer Center	Recruiting	Biliary Tract Cancer	Phase 2	35	NCT03639935
Pembrolizumab + Olaparib	PD-1, PARP	Georgetown University	Recruiting	Cholangiocarcinoma	Phase 2	29	NCT04306367
Pembrolizumab + Sargramostim	PD-1, GM-CSF	Robin Kate Kelley	Active, not recruiting	Biliary Cancer	Phase 2	42	NCT02703714
Entinostat + Nivolumab	PD-1, HDAC1/3	Sidney Kimmel Comprehensive cancer Center	Recruiting	Cholangiocarcinoma	Phase 2	54	NCT03250273
Nivolumab + Ipilimumab	PD-1, CTLA-4	National Cancer Institute	Recruiting	Cholangiocarcinoma	Phase 2	818	NCT02834013

Drug	Target	Sponsor	Status	Condition or disease	Phase	Enrollment	Trial Identifier
Durvalumab + Tremelimumab vs Durvalumab	PD-L1, CTLA-4	Institut für Klinische Krebsforschung IKF GmbH	Recruiting	Intrahepatic Cholangiocarcinoma	Phase 2	50	NCT04238637
CisGem + Nivolumab vs Nivolumab + Ipilimumab	Chemotherapy, PD-1, CTLA-4	University of Michigan Rogel Cancer Center	Active, not recruiting	Biliary Tract Neoplasms	Phase 2	64	NCT03101566
Durvalumab + Tremelimumab vs Durvalumab + Tremelimumab + TACE vs Durvalumab + Tremelimumab+ RFA vs Durvalumab + Tremelimumab+ Cryo	PD-L1, CTLA-4, Ablativetherapies	National Cancer Institute	Recruiting	Biliary Tract Neoplasms	Phase 2	90	NCT02821754
Nivolumab + Radiotherapy vs Nivolumab + Ipilimumab + Radiotherapy	PD-1, CTLA-4, radiation	Herlev Hospital	Recruiting	Metastatic Biliary Tract Cancer	Phase 2	160	NCT02866383
M7824	anti-PD-LxTGFβ fusion protein	EMD Serono Research & Development Institute	Recruiting	Biliary Tract Cancer Cholangiocarcinoma	Phase 2	141	NCT03833661
Pembrolizumab + Oxaliplatin + Capecitabine	PD-1, Chemotherapy	National Cancer Institute (NCI)	Recruiting	Biliary Tract Neoplasms Cholangiocarcinoma	Phase 2	19	NCT03111732
Trastuzumab + CisGem	HER2 , Chemotherapy	Changhoon Yoo	Recruiting	Cholangiocarcinoma Biliary Tract Cancer	Phase 2	15	NCT03613168
Immune cell							
TC-210 T Cells	Genetically engineered T cells	TCR2 Therapeutics	Recruiting	Cholangiocarcinoma	Phase 1 Phase 2	70	NCT03907852
MUC-1 CART cell	Target abnormal glycosylation MUC-1	Zhejiang University Second Hospital	Recruiting	Intrahepatic Cholangiocarcinoma	Phase 1 Phase 2	9	NCT03633773
Tumor Infiltrating Lymphocytes (TIL)	Tumor Infiltrating Lymphocytes	Udai Kammula	Recruiting	Cholangiocarcinoma	Phase 2	59	NCT03801083

FOLFIRINOX = irinotecan + oxaliplatin + fluorouracil + leucovorin; CisGem = cisplatin + gemcitabine; GEMOX=gemcitabine + oxaliplatin; FOLFOX= leucovorin calcium (folinic acid)+ fluorouracil+ oxaliplatin; Cryo= Cryoablation; PARP = poly-ADP ribose polymerase; TACE = trans-arterial chemoembolisation; RFA = radiofrequency ablation; CAR-T = Chimeric antigen receptor T-cell.



Advanced HCC

1st line	Sorafenib*	Lenvatinib*	Atezolizumab plus Bevacizumab			Tremelimumab plus Durvalumab			Donafenib
2nd line	VEGF/R inhibitors Regorafenib* Cabozantinib* Ramucirumab* Bevacizumab Apatinib	TGF-β inhibitors Galunisertib	c-MET inhibitors Tepotinib Tivantinib	CDKs inhibitors Palbociclib Ribociclib Milciclib	FGFR4 inhibitors Fisogatinib	PD-1 inhibitors Nivolumab* Pembrolizumab Tislelizumab Toripalimab Sintilimab Camrelizumab Penpulimab CS1003	PD-L1 inhibitors Atezolizumab Durvalumab Avelumab CS1001	CLAT-4 inhibitors Ipilimumab* Tremelimumab	Other novel therapies ALT-803 (IL-15 superagonist) KY1044 (T cell CO-stimulator) Pexa-Vec(oncolyticvirus) BMS-986253 (IL-8 antibody) Entecavir /tenofovir DF /Vemlidy (antiviral) CAR-T DC vaccines

B

Advanced ICC

1st line	Gemcitabine plus Cisplatin*			Gemcitabine plus Oxaliplatin			Capecitabine plus Oxaliplatin		
2nd line	Chemotherapy FOLFOX	FGFR inhibitors Pemigatinib* Infigratinib Futibatinib Derazantinib Erdafitinib Debio 1347	IDH1/2 inhibitors Ivosidenib FT-2102 BAY 1436032	VEGFR inhibitors Ramucirumab Apatinib Surufatinib	PARP inhibitors Niraparib Olaparib Rucaparib	PD-1 inhibitors Pembrolizumab Camrelizumab Nivolumab	PD-L1 inhibitors Durvalumab	CLAT-4 inhibitors Tremelimumab Ipilimumab	Other novel therapies CB-103(Notch) CX-2009(CD166) Entinostat(HDAC1/3) Trastuzumab(HER2) Bortezomib (Proteasome) Bintrafusp alfa (PD-L1 xTGF- β) XmAb22841 (CTLA-4 x LAG-3) M7824(PD-L1XTGF β)

Public Summary

1. Primary liver cancer comprises hepatocellular carcinoma (HCC), intrahepatic cholangiocarcinoma (ICC), combined HCC-ICC (cHCC-ICC), which are markedly distinct in their epidemiology, clinical features and response to therapy.
2. HCC is viral infection-related malignancy with specific histological features, whereas ICC is associated with chronic liver inflammation, showing more specific signatures.
3. HCC is prone to respond to targeted therapy, immunotherapy and antiviral agents, whereas ICCs are benefit from chemotherapy, targeted therapy and immunotherapy.
4. Combined cHCC-ICC subclass shows strong ICC-like features and is considered to be treated like ICC, whereas mixed cHCC-ICC subclass is shown to resemble HCC and is treated like HCC.