

# Journal Pre-proof

Comorbidities, clinical signs and symptoms, laboratory findings, imaging features, treatment strategies, and outcomes in adult and pediatric patients with COVID-19: A systematic review and meta-analysis

Catherine R. Jutzeler, PhD, Lucie Bourguignon, BSc, Caroline V. Weis, MSc, Bobo Tong, MPH, Cyrus Wong, BSc, Bastian Rieck, PhD, Hans Pargger, MD, Sarah Tschudin-Sutter, MD, Adrin Egli, MD PhD, Karsten Borgwardt, PhD, Matthias Walter, MD PhD

PII: S1477-8939(20)30321-5

DOI: <https://doi.org/10.1016/j.tmaid.2020.101825>

Reference: TMAID 101825

To appear in: *Travel Medicine and Infectious Disease*

Received Date: 14 May 2020


Revised Date: 9 July 2020

Accepted Date: 27 July 2020

Please cite this article as: Jutzeler CR, Bourguignon L, Weis CV, Tong B, Wong C, Rieck B, Pargger H, Tschudin-Sutter S, Egli A, Borgwardt K, Walter M, Comorbidities, clinical signs and symptoms, laboratory findings, imaging features, treatment strategies, and outcomes in adult and pediatric patients with COVID-19: A systematic review and meta-analysis, *Travel Medicine and Infectious Disease*, <https://doi.org/10.1016/j.tmaid.2020.101825>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 The Author(s). Published by Elsevier Ltd.



**Comorbidities, clinical signs and symptoms, laboratory findings, imaging features,  
treatment strategies, and outcomes in adult and pediatric patients with COVID-19:  
A systematic review and meta-analysis**

Catherine R. Jutzeler PhD<sup>1,2,3\*#</sup>, Lucie Bourguignon BSc<sup>1,2#</sup>, Caroline V. Weis MSc<sup>1,2</sup>, Bobo Tong MPH<sup>4</sup>,  
Cyrus Wong BSc<sup>5</sup>, Bastian Rieck PhD<sup>1,2</sup>, Hans Pargger MD<sup>6</sup>, Sarah Tschudin-Sutter MD<sup>7,8</sup>, Adrin Egli MD  
PhD<sup>9,10</sup>, Karsten Borgwardt PhD<sup>1,2\$</sup>, Matthias Walter MD PhD<sup>3,4,11\$</sup>

<sup>1</sup>*Department of Biosystems Science and Engineering, ETH Zurich, Basel, Switzerland*

<sup>2</sup>*SIB Swiss Institute of Bioinformatics, Lausanne, Switzerland*

<sup>3</sup>*Spinal Cord Injury Center, Balgrist University Hospital, University of Zurich, Zurich, Switzerland*

<sup>4</sup>*International Collaboration on Repair Discoveries (ICORD), University of British Columbia, Vancouver, Canada*

<sup>5</sup>*Simon Fraser University, Vancouver, Canada*

<sup>6</sup>*Intensive Care Unit, University Hospital Basel, University Basel, Basel, Switzerland*

<sup>7</sup>*Division of Infectious Diseases & Hospital Epidemiology, University Hospital Basel and University of Basel,  
Switzerland*

<sup>8</sup>*Department of Clinical Research, University Hospital Basel and University of Basel, Switzerland*

<sup>9</sup>*Division of Clinical Bacteriology & Mycology, University Hospital Basel, Basel, Switzerland*

<sup>10</sup>*Applied Microbiology Research, Department of Biomedicine, University of Basel, Basel, Switzerland*

<sup>11</sup>*Swiss Paraplegic Center, Nottwil, Switzerland*

<sup>#</sup>These authors share the lead authorship

<sup>\$</sup>These authors share the senior authorship

*\*Corresponding author*

Catherine Jutzeler

Department of Biosystems Science and Engineering, ETH Zurich

Mattenstrasse 26

Basel 4058

Switzerland

Email: Catherine.Jutzeler@bsse.ethz.ch

**Key words:** SARS-CoV-2, COVID-19, meta-analysis, systematic review, comorbidities, clinical characteristics, laboratory findings, imaging features, treatment

## Abstract

### Introduction

Since December 2019, a novel coronavirus (SARS-CoV-2) has triggered a world-wide pandemic with an enormous medical and societal-economic toll. Thus, our aim was to gather all available information regarding comorbidities, clinical signs and symptoms, outcomes, laboratory findings, imaging features, and treatments in patients with coronavirus disease 2019 (COVID-19).

### Methods

EMBASE, PubMed/ Medline, Scopus, and Web of Science were searched for studies published in any language between December 1<sup>st</sup>, 2019 and March 28<sup>th</sup>. Original studies were included if the exposure of interest was an infection with SARS-CoV-2 or confirmed COVID-19. The primary outcome was the risk ratio of comorbidities, clinical signs and symptoms, imaging features, treatments, outcomes, and complications associated with COVID-19 morbidity and mortality. We performed random-effects pairwise meta-analyses for proportions and relative risks,  $I^2$ ,  $\tau^2$ , and Cochrane Q, sensitivity analyses, and assessed publication bias.

### Results:

148 studies met the inclusion criteria for the systematic review and meta-analysis with 12'149 patients (5'739 female) and a median age of 47.0 [35.0-64.6] years. 617 patients died from COVID-19 and its complication. 297 patients were reported as asymptomatic. Older age (SMD: 1.25 [0.78- 1.72];  $p < 0.001$ ), being male (RR = 1.32 [1.13-1.54],  $p = 0.005$ ) and pre-existing comorbidity (RR = 1.69 [1.48-1.94];  $p < 0.001$ ) were identified as risk factors of in-hospital mortality. The heterogeneity between studies varied substantially ( $I^2$ ; range: 1.5-98.2%). Publication bias was only found in eight studies (Egger's test:  $p < 0.05$ ).

### Conclusions:

Our meta-analyses revealed important risk factors that are associated with severity and mortality of COVID-19.

## 1. Introduction

The severe acute respiratory syndrome (SARS) coronavirus 2 (SARS-CoV-2) initially emerged in Wuhan, Hubei, People's Republic of China and has been identified as the causative agent of coronavirus disease 2019 (COVID-19). It's pandemic spread presents a substantial medical challenge with an enormous societal and economic toll<sup>1,2</sup>. Similar to influenza and SARS-CoV-1, SARS-CoV-2 is considered a "crowd disease" that spreads most easily when individuals are packed together at high densities. Phylogenetic data implicate a zoonotic origin<sup>3</sup> and the rapid spread suggests ongoing person-to-person transmission<sup>4</sup>. Additional factors contributing to the rapid spread constitute the duration of the incubation period<sup>5</sup> and infectiousness peaking on or before symptom onset<sup>6</sup> contribute to the rapid spread of SARS-CoV-2. Another factor contributing to the rapid spread and alarmingly high number of infected people is the SARS-CoV-2 nature of initial dormancy of symptoms. The most common symptoms associated with COVID-19 include a sudden onset of fever, coughing, and dyspnea<sup>2,7,8</sup>. Complications comprise acute respiratory distress syndrome (ARDS), pneumonia, kidney failure, bacterial superinfections, coagulation abnormalities and thromboembolic events, sepsis, and even death<sup>9,10</sup>. So far, only a few demographic and clinical factors, such as older age, diabetes, and cardiovascular diseases, have been linked with poor outcome and increased risk of mortality<sup>11,12</sup>. This knowledge gap extends to the risk of infections, disease progression, and outcome in vulnerable patient populations, including newborns, children, pregnant, and elderly patients. A better understanding of the risks for these vulnerable patient populations is critical in order to optimize their protection and tailor prevention and treatment strategies. Thus, the aim of our systematic review and meta-analysis was to gather available information in the literature and determine the most prevalent comorbidities, clinical signs and symptoms, imaging features, laboratory parameters, treatments, outcomes, and complications arising in patients with COVID-19. We stratified our systematic reviews and meta-analysis by different cohorts, namely pediatric/neonatal and adult COVID-19 patients

including pregnant women. Furthermore, we aimed to assess current evidence for the associations between risk factors and in-hospital mortality. Based on previous reports, we addressed the hypothesis that male sex, older age, as well as pre-existing hypertension and diabetes mellitus are risk factors of morbidity and mortality in patients with COVID-19.

## **2. Methods**

Our systematic review and meta-analysis adhere to the Preferred Reporting Items for Systematic reviews and Meta-Analysis (PRISMA) statement<sup>13</sup> and Meta-analysis of Observational Studies in Epidemiology (MOOSE) checklist<sup>14</sup>.

### **2.1. Search strategy and selection criteria**

Four bibliographic databases were systematically searched: EMBASE, PubMed/ Medline, Scopus, and Web of Science. Our search was not restricted by language. We searched for studies published from December 1st, 2019 to March 28<sup>th</sup>, 2020, with search terms related to COVID-19 ("COVID-19", "SARS-CoV-2", "coronavirus disease 2019", "severe acute respiratory syndrome coronavirus 2", "2019 novel coronavirus", "2019-nCoV", "coronavirus", and "corona virus"). The full search strategy is provided in **Appendix 1**. Manual searching was also performed, reviewing reference lists of relevant studies and comprehensive review articles. Records were managed by EndNote X 8.0 software to exclude duplicates.

### **2.2. Selection of studies**

Two investigators (CRJ and MW) independently screened the titles and abstracts to determine whether studies should be included. Eligibility criteria were also applied to the full-text articles during the final selection. In case multiple articles reported on a single study, the article that provided the most data was selected for further synthesis. We quantified the inter-rater agreement for study selection using Cohen's  $\kappa$  coefficient<sup>15</sup>. Articles written in Chinese were reviewed by our two native speaking authors

(BT and CW) and if the inclusion criteria were met, these authors also extracted the specified data. All disagreements were discussed and resolved at a consensus meeting.

### **2.3. Inclusion and exclusion criteria**

All full-text, peer-reviewed articles that described case-control, cohort studies, or case studies investigating the epidemiological and clinical features, comorbidities, laboratory parameters, imaging features, and/or treatment of patients that were diagnosed with COVID-19. We excluded duplicate publications, non-peer reviewed articles (e.g., preprints), reviews, meta-analyses, abstracts or conference proceedings, editorials, commentaries, letters with insufficient data, studies on non-human species, or out-of-scope studies (e.g., comparison with other infections, case-fatality reports). In case multiple studies published data from the same cohort, we included the article representing the most inclusive information on the population to avoid overlap. Lastly, studies that did not report demographics (i.e., age and sex) were also excluded. **Figure 1** outlines our search strategy and application of inclusion and exclusion criteria.

### **2.4. Data extraction and synthesis**

Data extraction tables were created with the following information: 1) publication information (i.e., author, date, language of article, country where the study was performed, study design [case study, case series, or cohort study]<sup>16</sup>, study population [pediatric/neonatal and adult COVID-19 patients including pregnant women]; 2) demographics (i.e., age, sex); 3) clinical signs and symptoms (e.g., cough, fatigue, fever, sputum); 3) comorbidities (e.g., hypertension, diabetes, cardiovascular diseases); 4) therapies administered to treat COVID-19 (e.g., antibiotics, antivirals, invasive mechanical ventilation); 5) clinical outcomes (e.g., death, survival, recovery); and 6) complications associated with COVID-19 (e.g., sepsis and shock, ARDS). In case studies provided data for multiple patient groups

(e.g., pediatric and adult patient), we extracted this information separately for each group. A full list of extracted variables is provided in **Supplementary Table 1**.

## 2.5. Statistical analysis

For the studies reporting mean and standard deviation (SD) for extracted variables, we computed the median and interquartile ranges (IQR) assuming a normal distribution (i.e., using the formula:  $IQR \sim SD * 1.35$ ). To test if there is a bias by including the studies for which we computed the median and IQR (i.e., quartiles,  $Q_1$  and  $Q_3$ ), we performed a sensitivity analyses in which we calculated the median and IQR under the assumption of right-skewed and left-skewed distribution (see **Appendix 2**). We compared the results of the different distributions to test the robustness of our findings. Descriptive statistics (median, IQR,  $n$ , and %) were used to characterize the studies and patients included as well as the laboratory parameters. Weighted by study sample size, the pooled median and 95% confidence interval (CI) were computed for continuous variables. Normality approximation of the binomial was used to construct an approximate confidence interval (R package *metamedian*<sup>17</sup>). Welch's two-sample  $t$ -test was employed to test if there are significant differences in the proportion of male and female patients across studies.

Our meta-analysis was structured in two parts. In the first part, we performed meta-analyses of all 148 studies to define the prevalence of comorbidities, clinical signs and symptoms, imaging features, treatments, outcomes, and complications associated with COVID-19. Using the *metaprop* function of the R package *metafor*<sup>18</sup>, we calculated the overall prevalence from studies reporting a single prevalence. Our meta-analysis was stratified by patient group (pediatric/neonatal [ $\leq 17$  years of age], pregnant, and adult COVID-19 patients). Heterogeneity between studies was assessed visually by Forest plots, and analytically by  $I^2$ , tau  $T^2$ , and Cochrane  $Q$ . Briefly put,  $I^2$  describes the percentage of variation across studies that is due to heterogeneity rather than chance<sup>19</sup>: 0% indicates no heterogeneity, whereas 25%, 50%, and 75% indicate low, moderate, and high heterogeneity,

respectively. CIs for  $I^2$  were calculated using the iterative non-central chi-squared distribution method of Hedges and Piggott<sup>20</sup>. Tau ( $T^2$ ) represents the absolute value of the true variance (heterogeneity) and is the estimated SD of underlying true effects across studies. Cochran's Q is the weighted sum of squared differences between individual study effects and the pooled effect across studies, with the weights being those used in the pooling method (i.e., sample size)<sup>21</sup>. The second part comprised meta-analyses to calculate the relative risk (RR) of certain comorbidities, clinical signs and symptoms, imaging features, laboratory parameters, complications, and outcomes in patients with severe vs. those with non-severe disease condition (12 studies) as well as deceased vs. survivors (7 studies). The categorization into severe and non-severe COVID-19 disease was consistent with the groups reported by the reviewed studies (**Supplementary Table 2**). Owing to our judgment that considerable clinical and statistical heterogeneity exists among the studies (statistical heterogeneity was confirmed by the computed  $I^2$ ,  $T^2$ , and Cochrane Q), we calculated pooled RRs with 95% CIs using a random-effects model with inverse-variance weighting (*metabin* function from R package *meta*). For continuous outcome data (e.g., age, laboratory parameters, and time from symptoms onset to hospital admission), we estimated the standardized mean difference (SMD) by means of a random-effect models with inverse variance weighting for pooling (*metacont* function from R package *meta*). To calculate the SMD, we converted medians, Q1s, and Q3s into means and standard deviations. The SMD, 95% CIs, and p values were reported. We produced Forest plots to visualize the results from the random-effect models (R function: *forest*). Publication bias was assessed visually by funnel plots (R function: *funnel*) and analytically by the Egger test (R function: *regtest*). An Egger test  $p < 0.05$  indicates a significant publication bias. All statistical analyses were performed in R (version 3.6.3) for MacOS X(Mojave, 10.14.4) with the packages *meta* (version 4.11-0) and *dmetar* (version 0.0.90)<sup>18</sup>. The code used for the analysis and to create figures and tables is provided in our GitHub repository (<https://github.com/jutzca/Corona-Virus-Meta-Analysis-2020>).



## 2.6. Role of funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

## 3. Results

### *Study selection and study characteristics*

Our systematic literature search yielded 5'049 articles (including articles identified by manual searching). Upon removal of duplicates and exclusion of studies on the basis of their abstracts or following screening their full text, 148 met the inclusion criteria and were considered for the review and meta-analysis (**Figure 1**)<sup>4,8,28,118-127,29,128-137,30,138-147,31,148-157,32,158-165,33-37,9,38-47,11,48-57,22,58-67,23,68-77,24,78-87,25,88-97,26,98-107,27,108-117</sup>. The inter-rater agreement for study selection was very high ( $\kappa=0.94$  [95% CI: 0.91 - 0.96], 97.0% agreement [11/ 362 studies with disagreement]). Detailed information on the included studies are provided in **Tables 1-3**. Included studies were conducted in 15 countries between December 1<sup>st</sup>, 2019 and March 28<sup>th</sup>, 2020 (**Supplementary Table 3**) and enrolled between 1 and 1'099 patients (median 12.5 [1.00 – 56.75]). The majority of the articles were written in English (123 studies, 83.1%) and the remainder in Chinese (25 studies, 16.9%). We classified studies according to their design<sup>16</sup>: cohort study (76 studies, 51.4%), case study/ report (41 studies, 27.7%), and case series (31 studies, 20.9%). While all studies reported information on demographics (148, 100%), the number of studies reporting information on comorbidities (84 studies, 56.8%), clinical sign and symptoms (130 studies, 87.8%), laboratory parameters (113 studies, 76.4%), imaging features (118 studies, 79.7%), treatments (91, 61.5%), outcomes (118 studies, 79.3%), and complications (59 studies, 39.9%) varied markedly.

In terms of study population, 114 studies included only adult participants, 6 only pregnant women, 22 only children and neonates, and 6 included mixed cohorts. Of the total 12'149 patients included, 6'410 (52.8%) were male and 5'739 were female (47.2%, **Figure 2A and 2B**). The median age of adult (11'058

patients, 91.0%), pregnant (35 patients, 0.3%), and pediatric (1'056 patients, 8.7%; including neonates) patients was 47.0 years [35.0-65.3] (**Figure 3A**), 30.0 [26.0-33.0] (**Figure 3B**), and 10.0 [2.0-13.0] (**Figure 3C**), respectively. Approximately 7.8% (297/ 3'822 patients) were reported to be asymptomatic and 7.7% (617/ 8'047) died during hospitalization due to complications related to the infection with SARS-CoV-2. With the exception of one 10-month old child, all deaths were non-pregnant adult COVID-19 patients.

#### *Adult patients*

Higher proportions of male than female patients were reported to be infected with SARS-CoV-2 ( $t = 2.678$ ,  $df = 202$ ,  $p\text{-value} = 0.008$ ; **Figure 2A**) across all studies. Comorbidities were present in ~31% of the adult patients (2'329/ 7'608), with hypertension being the most prevalent one (1'352/ 6'460 patients, 20.93%), followed by heart failure (37/ 354 patients, 10.5%), diabetes mellitus (678/ 6'535 patients, 10.4%), and coronary heart disease (194/ 2'388 patients, 8.5%) (**Figure 4A, Table 4, Supplementary Figure 1**). The most frequent clinical signs and symptoms were fever (6'955/ 8'859 patients, 78.5%), cough (4'778/ 8'885 patients, 53.8%), and fatigue (1'996/ 7'980 patients, 25.0%) (**Figure 4B, Table 4**). A little over five percent of the adult COVID-19 patients were asymptomatic (148/ 2'749 patients, 5.4%). Over 6'969 patients (89.6%) had abnormal CT imaging features. The most common patterns of CT abnormalities were indicating pneumonia (unilateral or bilateral; 6'620/ 7'917 patients, 83.6%), including air bronchogram (264/ 523 patients, 50.5%), and ground-glass opacity (GGO) with consolidation (153/ 323 patients, 47.4%) and without (2'446/ 5'591 patients, 43.8%) (**Table 4, Supplementary Figure 2**). In terms of laboratory parameters, inflammatory markers, such as interleukin 6 (22 pg/mL [4.68-51.8]), and erythrocyte sedimentation rate (32.5 mm/h [17.3-53.8]) were elevated across the adult population. Moreover, markers of coagulation, namely d-dimer (0.5 µg/mL [0.3-1.08]), fibrinogen (4.5 g/L [3.66-5.1]), and cell damage were also elevated (i.e., lactate dehydrogenase, U/L; 213 [173-268]). An overview of all laboratory parameters is provided in

**Supplementary Table 4.** As shown in **Figure 4D**, the most common treatments were antivirals (4'475/ 6'068, patients, 73.8%), oxygen therapy (1'300/ 1'872 patients, 69.4%), and antibiotics (2'518/ 4'825 patients, 52.2%). Detailed information on all treatments is provided in **Table 4**. Eight percent (616/ 7'727 patients) of the adults died during the hospitalization due to complications related to COVID-19. Amongst the survivors (7'111/ 7'727 patients, 92.0%), a total of 3'025 (68.7%) remained hospitalized, 1'751 (32.4%) were discharged, and 1'012 (27.1%) reportedly recovered (**Figure 4C, Table 4**). Important to note, for some patients it was stated that they both, recovered and were discharged (i.e., one patient can fall in multiple categories). The median duration between symptoms onset and hospitalization was 8 days [7 - 9.5]. A total of 195 (6.8%) patients were admitted to the intensive care unit (ICU). The most frequently reported complications associated with COVID-19 were pneumonia (1'032/ 1'489 patients, 69.2%), respiratory failure (141/ 413 patients, 34.1%), acute cardiac injury (242/ 1'250 patients, 19.4%), and ARDS (759/ 5'122 patients, 14.8%), (**Figure 4D, Table 4**).

#### *Pregnant woman*

Studies investigating the effect of COVID-19 in pregnant women reported that only five pregnant women had any history of comorbidities. Hypothyroidism, allergies, or influenza were reported each for one pregnant woman (**Supplementary Table 5**). Fever (25/ 35 patients, 71.4%), cough (12/29 patients, 41.4%), and myalgia (3/ 9 patients, 33.3%) were the three most common symptoms observed in pregnant women that were infected with SARS-CoV-2 (**Supplementary Figure 3, Supplementary Table 5**). Abnormal CT features were evident in 88.6% (31/ 35 patients) of pregnant women diagnosed with COVID-19. Pneumonia (unilateral or bilateral, 31/ 35 patients, 88.6%), GGO (29/ 34 patients, 85.3%), and consolidation (8/ 16 patients, 50.0%) were among the most common patterns of CT abnormalities (**Supplementary Figure 4, Supplementary Table 5**). Inflammatory markers, such as C-reactive protein (19.25 mg/L [12.35-25.7]), procalcitonin (0.187 ng/mL), and neutrophil count ( $9.14 \times$

$10^9/L$ ) were elevated in this patient population. Along this line, lactate dehydrogenase concentrations were increased (544 U/L) reflecting cellular damage. An overview of all laboratory parameters is provided in **Supplementary Table 4**. Moreover, antibiotics (14/ 14 patients, 100.0%), antivirals (11/ 14 patients, 78.6%) and oxygen therapy (high flow nasal cannula; 3/ 12 patients, 25.0%) were used to treat pregnant COVID-19 patients (**Supplementary Table 5**). None of the pregnant COVID-19 patients died. Lastly, one patient was admitted to the ICU (**Supplementary Table 5**).

#### *Pediatric and Neonatal Patients*

Similar to the adult cohort, the proportion between female and male patients were comparable in the pediatric/neonatal cohort ( $t = 1.169$ ,  $df = 26$ ,  $p\text{-value} = 0.253$ ; **Figure 2B**). Fourteen percent of the children and neonates were asymptomatic (149/ 1'054). With the exception of two children, no comorbidities were reported for any of the pediatric or neonatal patients (**Supplementary Table 6**). Similar to the adult and pregnant COVID-19 patients, children and neonates frequently presented with fever (170/ 320 patients, 53.1%), cough (149/ 311 patients, 47.9%), and sputum (14/ 51 patients, 27.5%) (**Supplementary Figure 6 and Supplementary Table 6**). Sixty-five percent of the pediatric and neonatal patients presented with CT abnormalities, including pneumonia (194/ 298 patients), GGO (108/ 278 patients, 38.9%), and local patchy shadowing (52/ 223 patients, 23.3%) (**Supplementary Figure 7, Supplementary Table 6**). An overview of all laboratory parameters is provided in **Supplementary Table 7**. As the reference values vary considerably within the pediatric/neonatal patient population, the results of the laboratory parameters have to be interpreted with caution. In terms of treatment, children and neonates received antibiotics (31/ 43 patients, 72.1%), oxygen therapy through high flow nasal cannula (5/ 9 patients, 55.6%), and alpha interferon aerosol inhalation therapy (31/ 52, 59.6%) to treat COVID-19 and its complications (**Supplementary Figure 8, Supplementary Table 6**). With the exception of a 10-month-old child that died four weeks after admission of multi-

organ failure, all children survived. Less than 30% remained hospitalized (90/ 293 patients), 74.5% were discharged (216/ 290 patients) and 87.4% reportedly recovered (236/ 270 patients) (**Supplementary Figure 9, Supplementary Table 6**). The median duration between symptoms onset and hospitalization was 6 days [4.0-8.5]. Fifteen percent (6/ 39 patients) had to be admitted to the ICU. Complications associated with COVID-19 comprised pneumonia (16/ 26 patients, 61.5%), secondary bacterial infection (12/ 21 patients, 57.1%), and respiratory failure (10/ 33 patients, 30.3%) (**Supplementary Table 6**).

#### *Non-severe vs. Severe*

Twelve studies (2'596 patients) provided separate data for patients with a severe (500 patients, 19.3%) and non-severe disease status (2'096, 80.7%). No differences regarding sex were found between severe ( $t = 0.604$ ,  $df = 16.645$ ,  $p\text{-value} = 0.554$ ; male: 278 patients [55.6%] and female: 210 patients [42.0%]; unknown sex: 12 patients [2.4%]) and non-severe disease status group ( $t = 0.217$ ,  $df = 16.393$ ,  $p\text{-value} = 0.831$ ; male: 1'059 patients [50.5%] and female: 925 patients [49.5%]) (**Supplementary Figure 10**). In terms of age, patients with non-severe COVID-19 were significantly younger (median age in years= 45.0 [34.0-57.0]) than those with a severe disease progression (61.4 [44.5-75.5], **Figure 5**). Our meta-analysis revealed that older age (SMD: 0.68 [0.40-0.97];  $p < 0.001$ ), being male (RR = 1.11 [1.01-1.22];  $p = 0.039$ ), and preexisting comorbidities (RR=2.11 [1.02-4.35],  $p = 0.046$ ) were associated with a higher risk of increased disease severity. Specifically, hypertension (RR = 2.15 [1.64-2.81],  $p < 0.001$ ), diabetes mellitus (RR = 2.56 [1.50-4.39],  $p = 0.005$ ), any heart condition (RR = 4.09 [2.45-6.84],  $p < 0.001$ ), and chronic obstructive pulmonary disease (COPD, RR = 5.10 [3.08-8.45],  $p < 0.001$ ) (**Figure 6, Table 5**) were associated with worse outcome (i.e., severe disease). To test if the increased risk of heart conditions is attributable to the study that has classified their patients into severe and non-severe based on the presence or absence of cardiac injuries, we conducted a sensitivity analysis excluding these studies<sup>44</sup>. The risk of any heart condition remained significantly elevated in the severe disease patient cohort (RR

= 3.87 [1.85 - 8.11],  $p = 0.005$ ). Numerous laboratory parameters were significantly different between the non-severe and severe patient cohorts. Patients with severe disease status presented with decreased levels of albumin (SMD = 1.60 [-2.97 - (-0.24)];  $p = 0.022$ ), hemoglobin (SMD = -0.23 [-0.41 - (-0.06)];  $p = 0.001$ ), and thrombocytes (SMD = -0.57 [-0.68 - (-0.45)];  $p < 0.001$ ) in comparison to patients with non-severe disease status. Additionally, C-reactive protein (SMD = 1.47 [0.88-2.07];  $p < 0.001$ ), lactate dehydrogenase (SMD = 1.71 [1.08-2.34];  $p < 0.001$ ), and aspartate transaminase levels (SMD = 0.85 [0.61-1.09];  $p < 0.001$ ) were elevated in patients with severe disease status. In terms of complications, patients with severe COVID-19 disease were at an elevated risk of developing ARDS (RR=10.59 [2.44-46.01],  $p = 0.014$ , **Figure 6**). The heterogeneity between the studies varied substantially (**Table 5**). Publication bias, measured by means of the Egger's test, was only evident in three analyses. However, Egger's test may lack the statistical power to detect bias when the number of studies is small (i.e., fewer than 10) as we only included 4-8 studies.

#### *Survivor vs. non-survivors*

Seven studies (957 patients) provided disaggregated data for COVID-19 survivors (617 patients, 64.5%) and non-survivors (340, 35.5%). No differences regarding sex were found in the survivor group ( $t = 0.258$ ,  $df = 11.879$ ,  $p\text{-value} = 0.801$ ; male: 326 patients [52.8%] and female: 291 patients [47.2%]), but a significantly higher proportion of male patients were amongst the deceased cohort ( $t = 4.30$ ,  $df = 12$ ,  $p\text{-value} = 0.001$ ; male: 236 patients [69.4%] and female: 104 patients [30.6%]) (**Supplementary Figure 10**). In terms of age, COVID-19 patients that survived were significantly younger (median age in years= 52.0 [35.0-66.0]) than non-survivors (68.0 [62.0-76.0], **Figure 5**). The meta-analysis yielded older age (SMD: 1.25 [0.78- 1.72];  $p < 0.001$ ), being male (RR = 1.32 [1.13-1.54],  $p = 0.005$ ), pre-existing comorbidities (RR = 1.69 [1.48-1.94];  $p < 0.001$ ) as potential risk factors of in-hospital mortality. Pre-existing cerebrovascular diseases (RR = 36.88 [8.50-160.04];  $p = 0.009$ ), heart conditions (RR = 3.95

[1.03-15.20],  $p = 0.047$ , **Figure 7A**), and hypertension ( $RR = 2.09$  [1.65-2.64];  $p = 0.001$ ) were found to be associated with the highest risks of mortality. Clinical signs and symptoms as well as imaging features were comparable between survivors and non-survivors. In terms of treatments, non-survivors were more frequently mechanically ventilated than survivors ( $RR = 6.05$  [1.41-26.05];  $p = 0.026$ , **Figure 7B**) and more commonly received extracorporeal membrane oxygenation ( $RR = 4.39$  [1.64-11.78],  $p = 0.014$ ). Non-survivors had higher risks of complications, particularly acute kidney injury ( $RR = 20.77$  [2.43-177.44],  $p = 0.017$ ; **Figure 7C**) and ARDS ( $RR = 4.24$  [1.30-13.83],  $p = 0.026$ , **Figure 7D**). Low levels of albumin ( $SMD = -1.13$  [-1.41 – (-0.85)];  $p < 0.001$ ) and lymphocytes ( $SMD = -0.92$  [-1.3 – (-0.55)];  $p < 0.001$ ) as well as elevated level of interleukin 6 ( $SMD = 1.21$  [0.93 – 1.5];  $p < 0.001$ ), leucocytes ( $SMD = 2.21$  [0.61 – 3.64];  $p = 0.06$ ), and prolonged prothrombin time ( $SMD = 7.99$  [4.64 – 11.34];  $p < 0.01$ ) were associated with death (**Table 5**). Publication bias, measured by means of the Egger's test, was only evident in five analyses.

#### 4. Discussion

As of May 1<sup>st</sup>, 2020, more than 3.3 million confirmed cases of COVID-19 and more than 230'000 deaths attributable to the disease, have been reported worldwide<sup>166,167</sup>. In-depth knowledge of clinical, laboratory, and imaging factors that are associated with the disease progression and outcome is critical to inform clinical decision making and pandemic preparedness initiatives. An ever-growing number of research studies have been performed, but thus far the meta-analytical evidence is sparse. To address this paucity, we conducted a systematic review and meta-analysis of 148 studies involving over 12'000 patients providing an unprecedentedly comprehensive overview of comorbidities, clinical signs and symptoms, laboratory parameters, CT imaging features, treatment, outcomes, and complications in adult, pregnant, and pediatric/neonatal COVID-19 patients. Approximately eight percent of the patients were reported to be asymptomatic. However, this low number does not appear to reflect the

reality as the vast majority of the included studies primarily reported on symptomatic patients and were not designed to screen for asymptomatic patients. Furthermore, over seven percent died from complications associated with COVID-19. Recent analysis suggests that up to 75% of the coronavirus infections caused no illness<sup>168–170</sup>. Presumably, the virus has been circulating for longer than generally believed and large swathes of the population have already been exposed. Although our fatality rate lies within previous estimates<sup>171,172</sup>, it is important to mention that only a limited number of studies reported on the outcome of COVID-19 (i.e., death, survival, recovery) and thus, caution has to be exercised when interpreting this number. Through our meta-analysis, we revealed several important risk factors that are associated with severe disease progression and mortality. Among these risk factors were two demographic factors, namely older age and being male. Well-studied consequences of ageing are the decline in the immune function (e.g., T-cell and B-cell function) and excess production of type 2 cytokines<sup>173,174</sup>. These age-dependent changes in the immune response are suspected to cause deficiency in control of viral replication and more prolonged proinflammatory responses, potentially leading to poor outcome<sup>175</sup>. Corroborative evidence stems from preclinical studies that found an age-dependent host innate responses to virus infection in non-human primates inoculated with SARS-CoV-1<sup>176</sup>. Confirming previous findings<sup>177,178</sup>, sex-specific differences in mortality and vulnerability to the disease were evident in the current study. Specifically, men were disproportionately affected by an infection with SARS-CoV-2 (i.e., proportion of men presented with COVID-19 was larger compared to women) and the in-hospital mortality amongst male patients was significantly higher compared to female patients. Emerging evidence pinpoints towards differences in the immune system<sup>140</sup>, genetic polymorphism<sup>179</sup>, life style factors including smoking<sup>180</sup>, personal hygiene habits<sup>181</sup>, pre-existing comorbidities<sup>182,183</sup>, and expression of angiotensin-converting enzyme 2 (ACE2)<sup>184,185</sup> as potential explanations for the increased vulnerability in men. This sex difference in vulnerability has also been observed for SARS and MERS<sup>186</sup>, two previously emerging coronavirus diseases. The lack of sex-



disaggregated data in the reviewed studies made it impossible to further explore these potential explanations for the discrepant findings in men and women. Overall, the preexisting comorbidities, namely hypertension, diabetes mellitus, and any heart condition, were found to be linked with both, more severe diseases status and increased in-hospital mortality. Smoking, by contrast, was not associated with disease severity or mortality. However, the low number of studies reporting smoking status (13/ 148) cautions against early assumptions. Clinical signs and symptoms were comparable between patients with non-severe and severe COVID-19 as well as survivors and non-survivors. Fever, cough, and myalgia were amongst the most frequent reported symptoms across all groups. Similarly, the present study revealed no differences in the CT imaging features. The majority of the COVID-19 patients presented with pneumonia (bilateral or unilateral) and GGO. These pathological findings are a hallmark of any viral pneumonia, and thus it is not surprising that asymptomatic patients had similar distinctive features<sup>187</sup>. In terms of laboratory parameters, elevated levels of interleukin 6, leucocytes, d-dimer, and lactate dehydrogenase as well as hypoalbuminemia and lymphopenia were more commonly seen in patients with severe COVID-19 illness and non-survivors. High levels of d-dimer have a reported association with 28-day mortality in patients with infections or sepsis admitted to the intensive care unit<sup>188</sup>. Systemic pro-inflammatory cytokine responses (e.g., interleukin 6 and other components) contribute to host defense against infections, such as SARS-CoV-2<sup>189–191</sup>. However, exaggerated synthesis of interleukin 6 can lead to an acute, severe systemic inflammatory response syndrome (SIRS) known as 'cytokine storm'<sup>192</sup>. In addition to SIRS, hypoalbuminemia and lymphopenia were previously shown to be associated with increased odds of severe infection and infection-related death<sup>193–195</sup>. Complications were very common amongst patients with severe COVID-19 disease (over 50%) and non-survivors (more than two thirds). Acute cardiac injury, ARDS, and acute kidney injury were strongly linked to the outcome. Widely used treatments for COVID-19 and associated complications comprised antibiotics, antivirals, and oxygen therapy. Patients with severe COVID-19 disease required more often

mechanical ventilation and renal replacement therapy compared to those with non-severe COVID-19. Moreover, corticosteroids have been commonly administered to hospitalized patients with severe illness, although their benefit is highly disputed. Evidence from MERS or influenza suggests that patients who were given corticosteroids had prolonged viral replication, receive mechanical ventilation, and have higher mortality<sup>196–199</sup>. Administration of antibiotics and antivirals was independent of disease-severity.

Pregnant women as well as pediatric and neonatal patients may be less vulnerable to complications of COVID-19. Comorbidities were almost non-existent in these patient cohorts. Clinical signs and symptoms, laboratory parameters, imaging features, and treatments were comparable to the adult (non-pregnant) cohort. While there was a considerable proportion of children and neonates with SARS-CoV-2 infections reported, most of these patients did not need hospitalization and recovered quite well. With the exception of a 10-month old neonate, no children were amongst the deaths reported. All pregnant women included in our study survived COVID-19 and associated complications.

#### **4.1. Limitations of review**

A limitation of the current review was that literature search was limited to articles listed in EMBASE, PubMed/ Medline, Scopus, Web of Science, or identified by hand searches. Considering the pace at which the research in this area is moving forward, it is likely that the findings of the publications described in this paper will be quickly complemented by further research. The literature search also excluded grey literature (e.g., preprints, reports, conference proceedings), the importance of which to this topic is unknown, and thus might have introduced another source of search bias. There is also a probability of publication bias, as well as potential for a search bias. Publication bias is likely to result in studies with more positive results being preferentially submitted and accepted for publication. Moreover, geographical bias cannot be rule out as the majority of the studies (129/ 148) were

conducted in China. While symptoms might be quite comparable across countries, comorbidities, treatments, and outcome potentially depends on the country (and its healthcare system). There is also a considerable risk for a reporting bias towards comorbidities, clinical signs and symptoms, laboratory parameters, imaging features, treatment, outcome, and complications that are present. Specifically, only a minority of studies reported a zero when this information was assessed, but absent in patients. Lack of data on absent clinical signs and symptoms might lead to distorted estimates of proportion. Furthermore, the low number of asymptomatic patients must be considered with caution. The meta-analysis of severity and mortality could only be performed with a small number of studies as the minority of the 148 provided data separately for different disease severity groups (e.g., non-severe, severe, survivors, non-survivors). This needs to be considered when interpreting the results, including the publication bias as the Egger's test may lack the statistical power to detect bias when the number of studies is small (i.e.,  $< 10$ ). Lastly, the criteria to classify patients in severe and non-severe COVID-19 disease cohorts varied between studies leading to additional heterogeneity between studies. By virtue of low number of studies available, we could not assess this heterogeneity nor adjust for it.

#### **4.2. Conclusion and future directions**

In conclusion, this unprecedentedly comprehensive systematic review and meta-analysis of the literature published during the first 120 days of the COVID-19 pandemic yields important information regarding the comorbidities, clinical signs and symptoms, laboratory parameters, imaging features, treatment, outcome, and complications. Male sex, older age, and pre-existing comorbidities are major risk factors for in-hospital mortality and complications. This study revealed a fatality rate of 7.7% and found that approximately 8% of the patients were reportedly asymptomatic. Based on recent reports, the latter number is likely 6- to 10-fold higher as only a few asymptomatic patients are captured by the health care system as they do not seek medical attention due to the lack of symptoms<sup>168</sup> or are not hospitalized and thus, included in studies. Unnoticed asymptomatic cases of COVID-19 are likely a

major source of ongoing transmission. Children and neonates appear to be the least vulnerable cohort. Forthcoming studies are needed that provide sex-disaggregated data to better characterize risk factors that affect both sexes or are specific to men or women, respectively.

#### **Authors' contribution**

**Catherine Jutzeler:** Substantial contributions to the conception and design of the study; acquisition, analysis, and interpretation of data, drafting the manuscript, final approval of version to be published.

**Lucie Bourguignon:** Substantial contributions to acquisition, analysis, and interpretation of data, drafting the manuscript, final approval of version to be published.

**Caroline Weis:** Acquisition and interpretation of data, revising the manuscript critically for important intellectual content, final approval of version to be published.

**Bastian Rieck:** Acquisition and interpretation of data, revising the manuscript critically for important intellectual content, final approval of version to be published.

**Bobo Tong:** Acquisition and interpretation of data, revising the manuscript critically for important intellectual content, final approval of version to be published.

**Cyrus Wong:** Acquisition and interpretation of data, revising the manuscript critically for important intellectual content, final approval of version to be published.

**Hans Pargger:** Substantial contributions to the interpretation of data, revising the manuscript critically for important intellectual content, final approval of version to be published.

**Sarah Tschudin-Sutter:** Substantial contributions to the interpretation of data, revising the manuscript critically for important intellectual content, final approval of version to be published.

**Adrian Egli:** Substantial contributions to the interpretation of data, revising the manuscript critically for important intellectual content, final approval of version to be published.

**Karsten Borgwardt:** Substantial contributions to the interpretation of data, revising the manuscript critically for important intellectual content, final approval of version to be published.

**Matthias Walter:** Substantial contributions to the conception and design of the study; acquisition, analysis, and interpretation of data, drafting the manuscript, final approval of version to be published.

#### **4.3. Conflict of interest**

The authors do not report any (financial or otherwise) conflict of interest.

#### **4.4. Acknowledgement**

This study was supported by the Alfried Krupp Prize for Young University Teachers of the Alfried Krupp von Bohlen und Halbach-Stiftung (Borgwardt), and the Swiss National Science Foundation (Ambizione

Grant [PZooP<sub>3</sub>\_186101, Jutzeler], SNSF Starting Grant [155913, Borgwardt}). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Journal Pre-proof

## 5. References

1. Tuite AR, Bogoch II, Sherbo R, Watts A, Fisman D, Khan K. Estimation of Coronavirus Disease 2019 (COVID-19) Burden and Potential for International Dissemination of Infection From Iran. *Ann Intern Med*. 2020. doi:10.7326/m20-0696
2. Wang C, Horby PW, Hayden FG, Gao GF. A novel coronavirus outbreak of global health concern. *Lancet*. 2020. doi:10.1016/S0140-6736(20)30185-9
3. Lu R, Zhao X, Li J, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet*. 2020. doi:10.1016/S0140-6736(20)30251-8
4. Chan JFW, Yuan S, Kok KH, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet*. 2020. doi:10.1016/S0140-6736(20)30154-9
5. Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med*. 2020. doi:10.1056/NEJMo2001316
6. He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med*. 2020. doi:10.1038/s41591-020-0869-5
7. Jiang F, Deng L, Zhang L, Cai Y, Cheung CW, Xia Z. Review of the Clinical Characteristics of Coronavirus Disease 2019 (COVID-19). *J Gen Intern Med*. 2020. doi:10.1007/s11606-020-05762-w
8. Xu XW, Wu XX, Jiang XG, et al. Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARS-Cov-2) outside of Wuhan, China: Retrospective case series. *BMJ*. 2020. doi:10.1136/bmj.m606
9. Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med*. 2020. doi:10.1016/S2213-2600(20)30079-5
10. Llitjos J-F, Leclerc M, Chochois C, et al. High incidence of venous thromboembolic events in anticoagulated severe COVID-19 patients. *J Thromb Haemost*. 2020. doi:10.1111/jth.14869
11. Chen H, Guo J, Wang C, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet*. 2020;395(10226):809-815. doi:10.1016/S0140-6736(20)30360-3
12. Xu. Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARS-Cov-2) outside of Wuhan, China: retrospective case series (vol 368, m606, 2020). *Bmj-British Med J*. 2020;368:1. doi:10.1136/bmj.m792
13. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *PLoS Med*. 2009;6(7). doi:10.1371/journal.pmed.1000100
14. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: A

- proposal for reporting. *J Am Med Assoc.* 2000. doi:10.1001/jama.283.15.2008
15. Viera AJ, Garrett JM. Understanding interobserver agreement: The kappa statistic. *Fam Med.* 2005.
  16. Mathes T, Pieper D. Clarifying the distinction between case series and cohort studies in systematic reviews of comparative studies: Potential impact on body of evidence and workload. *BMC Med Res Methodol.* 2017. doi:10.1186/s12874-017-0391-8
  17. McGrath S, Zhao XF, Qin ZZ, Steele R, Benedetti A. One-sample aggregate data meta-analysis of medians. *Stat Med.* 2019. doi:10.1002/sim.8013
  18. Balduzzi S, Rücker G, Schwarzer G. How to perform a meta-analysis with R: A practical tutorial. *Evid Based Ment Health.* 2019. doi:10.1136/ebmental-2019-300117
  19. Bollegala D, Matsuo Y, Ishizuka M. Measuring semantic similarity between words using web search engines. In: *16th International World Wide Web Conference, WWW2007.* ; 2007. doi:10.1145/1242572.1242675
  20. Hedges L V., Pigott TD. The power of statistical tests for moderators in meta-analysis. *Psychol Methods.* 2004. doi:10.1037/1082-989X.9.4.426
  21. Cochran WG. The Combination of Estimates from Different Experiments. *Biometrics.* 1954. doi:10.2307/3001666
  22. Xu X, Yu C, Qu J, et al. Imaging and clinical features of patients with 2019 novel coronavirus SARS-CoV-2. *Eur J Nucl Med Mol Imaging.* 2020. doi:10.1007/s00259-020-04735-9
  23. Ye G, Pan Z, Pan Y, et al. Clinical characteristics of severe acute respiratory syndrome coronavirus 2 reactivation. *J Infect.* 2020. doi:10.1016/j.jinf.2020.03.001
  24. Le HT, Nguyen L V, Tran DM, et al. The first infant case of COVID-19 acquired from a secondary transmission in Vietnam. *Lancet Child Adolesc Heal.* 2020. doi:10.1016/s2352-4642(20)30091-2
  25. Guan Q, Liu M, Zhuang YJ, et al. Epidemiological investigation of a family clustering of COVID-19. *Zhonghua Liu Xing Bing Xue Za Zhi.* 2020;41:629-633. doi:10.3760/cma.j.cn112338-20200223-00152
  26. Wang L, Gao YH, Lou LL, Zhang GJ. The clinical dynamics of 18 cases of COVID-19 outside of Wuhan, China. *Eur Respir J.* 2020. doi:10.1183/13993003.00398-2020
  27. Zhang MQ, Wang XH, Chen YL, et al. [Clinical features of 2019 novel coronavirus pneumonia in the early stage from a fever clinic in Beijing]. *Zhonghua Jie He He Hu Xi Za Zhi.* 2020;43(3):215-218. doi:10.3760/cma.j.issn.1001-0939.2020.03.015
  28. Ding D, Zhu C, Yao W. A cured patient with 2019-nCoV pneumonia. *Am J Med.* 2020. doi:10.1016/j.amjmed.2020.02.023
  29. Chen S, Huang B, Luo DJ, et al. [Pregnant women with new coronavirus infection: a clinical characteristics and placental pathological analysis of three cases]. *Zhonghua Bing Li Xue Za Zhi.*

- 2020;49(0):E005. doi:10.3760/cma.j.cn112151-20200225-00138
30. Hao W. Clinical Features of Atypical 2019 Novel Coronavirus Pneumonia with an initially Negative RT-PCR Assay. *J Infect.* 2020. doi:10.1016/j.jinf.2020.02.008
  31. Spiteri G, Fielding J, Diercke M, et al. First cases of coronavirus disease 2019 (COVID-19) in the WHO European Region, 24 January to 21 February 2020. *Euro Surveill.* 2020;25(9). doi:10.2807/1560-7917.Es.2020.25.9.2000178
  32. Ai T, Yang Z, Hou H, et al. Correlation of Chest CT and RT-PCR Testing in Coronavirus Disease 2019 (COVID-19) in China: A Report of 1014 Cases. *Radiology.* 2020:200642. doi:10.1148/radiol.2020200642
  33. Cui Y, Tian M, Huang D, et al. A 55-Day-Old Female Infant infected with COVID 19: presenting with pneumonia, liver injury, and heart damage. *J Infect Dis.* 2020. doi:10.1093/infdis/jiaa113
  34. Zhang JJ, Dong X, Cao YY, et al. Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy Eur J Allergy Clin Immunol.* 2020. doi:10.1111/all.14238
  35. Xu Y, Li XF, Zhu B, et al. Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nat Med.*:9. doi:10.1038/s41591-020-0817-4
  36. Peng YD, Meng K, Guan HQ, et al. Clinical characteristics and outcomes of 112 cardiovascular disease patients infected by 2019-nCoV. *Zhonghua Xin Xue Guan Bing Za Zhi.* 2020;48:E004. doi:10.3760/cma.j.cn112148-20200220-00105
  37. Lu X, Zhang L, Du H, et al. SARS-CoV-2 Infection in Children. *N Engl J Med.* 2020. doi:10.1056/NEJMc2005073
  38. Tong ZD, Tang A, Li KF, et al. Potential Presymptomatic Transmission of SARS-CoV-2, Zhejiang Province, China, 2020. *Emerg Infect Dis.* 2020;26(5). doi:10.3201/eid2605.200198
  39. Chen J, Qi T, Liu L, et al. Clinical progression of patients with COVID-19 in Shanghai, China. *J Infect.* 2020. doi:10.1016/j.jinf.2020.03.004
  40. Lin X, Gong Z, Xiao Z, Xiong J, Fan B, Liu J. Novel coronavirus pneumonia outbreak in 2019: Computed tomographic findings in two cases. *Korean J Radiol.* 2020;21(3):365-368. doi:10.3348/kjr.2020.0078
  41. Xu T, Chen C, Zhu Z, et al. Clinical features and dynamics of viral load in imported and non-imported patients with COVID-19. *Int J Infect Dis.* 2020. doi:10.1016/j.ijid.2020.03.022
  42. Yao N, Wang SN, Lian JQ, et al. Clinical characteristics and influencing factors of patients with novel coronavirus pneumonia combined with liver injury in Shaanxi region. *Zhonghua Gan Zang Bing Za Zhi.* 2020;28:E003. doi:10.3760/cma.j.cn501113-20200226-00070
  43. Tang A, Tong ZD, Wang HL, et al. Detection of Novel Coronavirus by RT-PCR in Stool Specimen from Asymptomatic Child, China. *Emerg Infect Dis.* 2020;26(6). doi:10.3201/eid2606.200301
  44. Shi S, Qin M, Shen B, et al. Association of Cardiac Injury With Mortality in Hospitalized Patients



- With COVID-19 in Wuhan, China. *JAMA Cardiol.* 2020. doi:10.1001/jamacardio.2020.0950
45. Wang Y, Liu Y, Liu L, Wang X, Luo N, Ling L. Clinical Outcomes in 55 Patients With Severe Acute Respiratory Syndrome Coronavirus 2 Who Were Asymptomatic at Hospital Admission in Shenzhen, China. *J Infect Dis.* 2020. doi:10.1093/infdis/jiaa119
  46. Dong L, Tian J, He S, et al. Possible Vertical Transmission of SARS-CoV-2 From an Infected Mother to Her Newborn. *JAMA.* 2020. doi:10.1001/jama.2020.4621
  47. Leung C. Clinical features of deaths in the novel coronavirus epidemic in China. *Rev Med Virol.* 2020. doi:10.1002/rmv.2103
  48. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet.* 2020. doi:10.1016/S0140-6736(20)30566-3
  49. Fan C, Lei D, Fang C, et al. Perinatal Transmission of COVID-19 Associated SARS-CoV-2: Should We Worry? *Clin Infect Dis.* 2020. doi:10.1093/cid/ciaa226
  50. Xiong Y, Sun D, Liu Y, et al. Clinical and High-Resolution CT Features of the COVID-19 Infection: Comparison of the Initial and Follow-up Changes. *Invest Radiol.* 2020. doi:10.1097/RLI.0000000000000674
  51. Shi H, Han X, Jiang N, et al. Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study. *Lancet Infect Dis.* 2020. doi:10.1016/S1473-3099(20)30086-4
  52. He XW, Lai JS, Cheng J, et al. Impact of complicated myocardial injury on the clinical outcome of severe or critically ill COVID-19 patients. *Zhonghua Xin Xue Guan Bing Za Zhi.* 2020;48:E011. doi:10.3760/cma.j.cn112148-20200228-00137
  53. Wei M, Yuan J, Liu Y, Fu T, Yu X, Zhang ZJ. Novel Coronavirus Infection in Hospitalized Infants under 1 Year of Age in China. *JAMA - J Am Med Assoc.* 2020. doi:10.1001/jama.2020.2131
  54. Lee NY, Li CW, Tsai HP, et al. A case of COVID-19 and pneumonia returning from Macau in Taiwan: Clinical course and anti-SARS-CoV-2 IgG dynamic. *J Microbiol Immunol Infect.* 2020. doi:10.1016/j.jmii.2020.03.003
  55. Yuan M, Yin W, Tao Z, Tan W, Hu Y. Association of radiologic findings with mortality of patients infected with 2019 novel coronavirus in Wuhan, China. *PLoS One.* 2020;15(3):e0230548. doi:10.1371/journal.pone.0230548
  56. Wang Z, Chen X, Lu Y, Chen F, Zhang W. Clinical characteristics and therapeutic procedure for four cases with 2019 novel coronavirus pneumonia receiving combined Chinese and Western medicine treatment. *Biosci Trends.* 2020;14(1):64-68. doi:10.5582/bst.2020.01030
  57. Hosoda T, Sakamoto M, Shimizu H, Okabe N. SARS-CoV-2 enterocolitis with persisting to excrete the virus for about two weeks after recovering from diarrhea: A case report. *Infect Control Hosp Epidemiol.* 2020;1-4. doi:10.1017/ice.2020.87
  58. Zhu Y, Liu YL, Li ZP, et al. Clinical and CT imaging features of 2019 novel coronavirus disease

- (COVID-19). *J Infect.* 2020. doi:10.1016/j.jinf.2020.02.022
59. Arentz M, Yim E, Klaff L, et al. Characteristics and Outcomes of 21 Critically Ill Patients With COVID-19 in Washington State. *JAMA.* 2020. doi:10.1001/jama.2020.4326
  60. Song Y, Liu P, Shi XL, et al. SARS-CoV-2 induced diarrhoea as onset symptom in patient with COVID-19. *Gut.* 2020. doi:10.1136/gutjnl-2020-320891
  61. Qu R, Ling Y, Zhang YH, et al. Platelet-to-lymphocyte ratio is associated with prognosis in patients with Corona Virus Disease-19. *J Med Virol.* 2020. doi:10.1002/jmv.25767
  62. Cheng Z, Lu Y, Cao Q, et al. Clinical Features and Chest CT Manifestations of Coronavirus Disease 2019 (COVID-19) in a Single-Center Study in Shanghai, China. *AJR Am J Roentgenol.* 2020;1-6. doi:10.2214/AJR.20.22959
  63. Jin X, Lian JS, Hu JH, et al. Epidemiological, clinical and virological characteristics of 74 cases of coronavirus-infected disease 2019 (COVID-19) with gastrointestinal symptoms. *Gut.* 2020. doi:10.1136/gutjnl-2020-320926
  64. Xu Z, Shi L, Wang Y, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir Med.* 2020. doi:10.1016/S2213-2600(20)30076-X
  65. Cai JH, Wang XS, Ge YL, et al. First case of 2019 novel coronavirus infection in children in Shanghai. *Zhonghua er ke za zhi = Chinese J Pediatr.* 2020;58:E002. doi:10.3760/cma.j.issn.0578-1310.2020.0002
  66. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet.* 2020;395(10223):497-506. doi:10.1016/S0140-6736(20)30183-5
  67. Chen C, Chen C, Yan JT, Zhou N, Zhao JP, Wang DW. [Analysis of myocardial injury in patients with COVID-19 and association between concomitant cardiovascular diseases and severity of COVID-19]. *Zhonghua Xin Xue Guan Bing Za Zhi.* 2020;48(0):E008. doi:10.3760/cma.j.cn112148-20200225-00123
  68. Deng Y, Liu W, Liu K, et al. Clinical characteristics of fatal and recovered cases of coronavirus disease 2019 (COVID-19) in Wuhan, China: a retrospective study. *Chin Med J.* 2020. doi:10.1097/cm9.0000000000000824
  69. COVID-19 National Emergency Response Center E and CMT, Korea Centers for Disease Control and Prevention Korea C. Early Epidemiological and Clinical Characteristics of 28 Cases of Coronavirus Disease in South Korea. *Osong Public Heal Res Perspect.* 2020;11(1):8-14. doi:10.24171/j.phrp.2020.11.1.03
  70. Wan S, Xiang Y, Fang W, et al. Clinical Features and Treatment of COVID-19 Patients in Northeast Chongqing. *J Med Virol.* 2020. doi:10.1002/jmv.25783
  71. Zhang X, Song W, Liu X, Lyu L. CT image of novel coronavirus pneumonia: a case report. *Jpn J Radiol.* 2020. doi:10.1007/s11604-020-00945-1
  72. Ding Q, Lu P, Fan Y, Xia Y, Liu M. The clinical characteristics of pneumonia patients co-infected

- with 2019 novel coronavirus and influenza virus in Wuhan, China. *J Med Virol.* 2020. doi:10.1002/jmv.25781
73. Xu YH, Dong JH, An WM, et al. Clinical and computed tomographic imaging features of novel coronavirus pneumonia caused by SARS-CoV-2. *J Infect.* 2020. doi:10.1016/j.jinf.2020.02.017
  74. Liu Y, Yang Y, Zhang C, et al. Clinical and biochemical indexes from 2019-nCoV infected patients linked to viral loads and lung injury. *Sci China Life Sci.* 2020;63(3):364-374. doi:10.1007/s11427-020-1643-8
  75. Chen R, Chen J, Meng QT. Chest computed tomography images of early coronavirus disease (COVID-19). *Can J Anesth.* 2020. doi:10.1007/s12630-020-01625-4
  76. Huang Y, Tu M, Wang S, et al. Clinical characteristics of laboratory confirmed positive cases of SARS-CoV-2 infection in Wuhan, China: A retrospective single center analysis. *Travel Med Infect Dis.* 2020. doi:10.1016/j.tmaid.2020.101606
  77. Lian J, Jin X, Hao S, et al. Analysis of Epidemiological and Clinical features in older patients with Corona Virus Disease 2019 (COVID-19) out of Wuhan. *Clin Infect Dis.* 2020. doi:10.1093/cid/ciaa242
  78. Zhao D, Yao F, Wang L, et al. A comparative study on the clinical features of COVID-19 pneumonia to other pneumonias. *Clin Infect Dis.* 2020. doi:10.1093/cid/ciaa247
  79. Young BE, Ong SWX, Kalimuddin S, et al. Epidemiologic Features and Clinical Course of Patients Infected With SARS-CoV-2 in Singapore. *JAMA.* 2020. doi:10.1001/jama.2020.3204
  80. Qian G, Yang N, Ma AHY, et al. A COVID-19 Transmission within a family cluster by presymptomatic infectors in China. *Clin Infect Dis.* 2020. doi:10.1093/cid/ciaa316
  81. Liu W, Tao ZW, Lei W, et al. Analysis of factors associated with disease outcomes in hospitalized patients with 2019 novel coronavirus disease. *Chin Med J (Engl).* 2020. doi:10.1097/CM9.0000000000000775
  82. Xie C, Jiang L, Huang G, et al. Comparison of different samples for 2019 novel coronavirus detection by nucleic acid amplification tests. *Int J Infect Dis.* 2020;93:264-267. doi:10.1016/j.ijid.2020.02.050
  83. Wang J, Wang D, Chen GC, Tao XW, Zeng LK. [SARS-CoV-2 infection with gastrointestinal symptoms as the first manifestation in a neonate]. *Zhongguo Dang Dai Er Ke Za Zhi.* 2020;22(3):211-214.
  84. Tian S, Hu W, Niu L, Liu H, Xu H, Xiao SY. Pulmonary Pathology of Early-Phase 2019 Novel Coronavirus (COVID-19) Pneumonia in Two Patients With Lung Cancer. *J Thorac Oncol.* 2020. doi:10.1016/j.jtho.2020.02.010
  85. Wu T, Kang SC, Feng W, et al. Biological characters analysis of COVID-19 patient accompanied with aplastic anemia. *Zhonghua Xue Ye Xue Za Zhi.* 2020;41:E003. doi:10.3760/cma.j.issn.0253-2727.2020.0003

86. Liu W, Zhang Q, Chen J, et al. Detection of Covid-19 in Children in Early January 2020 in Wuhan, China. *N Engl J Med*. 2020. doi:10.1056/NEJMc2003717
87. Feng K, Yun YX, Wang XF, et al. Analysis of CT features of 15 Children with 2019 novel coronavirus infection. *Zhonghua er ke za zhi = Chinese J Pediatr*. 2020;58:E007. doi:10.3760/cma.j.issn.0578-1310.2020.0007
88. Fang Z, Zhang Y, Hang C, Zhang W, Ai J, Li S. Comparisons of nucleic acid conversion time of SARS-CoV-2 of different samples in ICU and non-ICU patients. *J Infect*. 2020. doi:10.1016/j.jinf.2020.03.013
89. Sun Q, Xu X, Xie J, Li J, Huang X. Evolution of Computed Tomography Manifestations in Five Patients Who Recovered from Coronavirus Disease 2019 (COVID-19) Pneumonia. *Korean J Radiol*. 2020. doi:10.3348/kjr.2020.0157
90. Liu M, He P, Liu HG, et al. [Clinical characteristics of 30 medical workers infected with new coronavirus pneumonia]. *Zhonghua Jie He He Hu Xi Za Zhi*. 2020;43(3):209-214. doi:10.3760/cma.j.issn.1001-0939.2020.03.014
91. Silverstein WK, Stroud L, Cleghorn GE, Leis JA. First imported case of 2019 novel coronavirus in Canada, presenting as mild pneumonia. *Lancet*. 2020;395(10225):734. doi:10.1016/S0140-6736(20)30370-6
92. Dong Y, Mo X, Hu Y, et al. Epidemiological Characteristics of 2143 Pediatric Patients With 2019 Coronavirus Disease in China. *Pediatrics*. 2020. doi:10.1542/peds.2020-0702
93. Fang X, Zhao M, Li S, Yang L, Wu B. Changes of CT Findings in a 2019 Novel Coronavirus (2019-nCoV) pneumonia patient. *QJM*. 2020. doi:10.1093/qjmed/hcaa038
94. An P, Song P, Lian K, Wang Y. CT Manifestations of Novel Coronavirus Pneumonia: A Case Report. *Balkan Med J*. 2020. doi:10.4274/balkanmedj.galenos.2020.2020.2.15
95. Chen L, Liu HG, Liu W, et al. [Analysis of clinical features of 29 patients with 2019 novel coronavirus pneumonia]. *Zhonghua Jie He He Hu Xi Za Zhi*. 2020;43(3):203-208. doi:10.3760/cma.j.issn.1001-0939.2020.03.013
96. Li XQ, Cai WF, Huang LF, et al. Comparison of epidemic characteristics between SARS in 2003 and COVID-19 in 2020 in Guangzhou. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2020;41(5):634-637. doi:10.3760/cma.j.cn112338-20200228-00209
97. Zeng LK, Tao XW, Yuan WH, Wang J, Liu X, Liu ZS. First case of neonate infected with novel coronavirus pneumonia in China. *Zhonghua er ke za zhi = Chinese J Pediatr*. 2020;58:E009. doi:10.3760/cma.j.issn.0578-1310.2020.0009
98. Mo P, Xing Y, Xiao Y, et al. Clinical characteristics of refractory COVID-19 pneumonia in Wuhan, China. *Clin Infect Dis*. 2020. doi:10.1093/cid/ciaa270
99. Li YX, Wu W, Yang T, et al. Characteristics of peripheral blood leukocyte differential counts in patients with COVID-19. *Zhonghua nei ke za zhi*. 2020;59:E003. <http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L631125257>.

100. Wang Z, Yang B, Li Q, Wen L, Zhang R. Clinical Features of 69 Cases with Coronavirus Disease 2019 in Wuhan, China. *Clin Infect Dis*. 2020. doi:10.1093/cid/ciaa272
101. Chen F, Liu ZS, Zhang FR, et al. First case of severe childhood novel coronavirus pneumonia in China. *Zhonghua er ke za zhi = Chinese J Pediatr*. 2020;58(3):179-182. doi:10.3760/cma.j.issn.0578-1310.2020.03.003
102. Hu X, Chen J, Jiang X, et al. CT imaging of two cases of one family cluster 2019 novel coronavirus (2019-nCoV) pneumonia: inconsistency between clinical symptoms amelioration and imaging sign progression. *Quant Imaging Med Surg*. 2020;10(2):508-510. doi:10.21037/qims.2020.02.10
103. Shi H, Han X, Zheng C. Evolution of CT Manifestations in a Patient Recovered from 2019 Novel Coronavirus (2019-nCoV) Pneumonia in Wuhan, China. *Radiology*. 2020;295(1):20. doi:10.1148/radiol.2020200269
104. Han W, Quan B, Guo Y, et al. The course of clinical diagnosis and treatment of a case infected with coronavirus disease 2019. *J Med Virol*. 2020;92(5):461-463. doi:10.1002/jmv.25711
105. Dai WC, Zhang HW, Yu J, et al. CT Imaging and Differential Diagnosis of COVID-19. *Can Assoc Radiol J*. 2020;84:6537120913033. doi:10.1177/0846537120913033
106. Dong XC, Li JM, Bai JY, et al. Epidemiological characteristics of confirmed COVID-19 cases in Tianjin. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2020;41(5):638-642. doi:10.3760/cma.j.cn112338-20200221-00146
107. Wang D, Ju XL, Xie F, et al. Clinical analysis of 31 cases of 2019 novel coronavirus infection in children from six provinces (autonomous region) of northern China. *Zhonghua er ke za zhi = Chinese J Pediatr*. 2020;58(4):E011. doi:10.3760/cma.j.cn112140-20200225-00138
108. Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost*. 2020. doi:10.1111/jth.14768
109. Huang P, Liu T, Huang L, et al. Use of Chest CT in Combination with Negative RT-PCR Assay for the 2019 Novel Coronavirus but High Clinical Suspicion. *Radiology*. 2020;295(1):22-23. doi:10.1148/radiol.2020200330
110. Zhu ZW, Tang JJ, Chai XP, et al. Comparison of heart failure and 2019 novel coronavirus pneumonia in chest CT features and clinical characteristics. *Zhonghua Xin Xue Guan Bing Za Zhi*. 2020;48:E007. doi:10.3760/cma.j.cn112148-20200218-00093
111. Li YY, Wang WN, Lei Y, et al. [Comparison of the clinical characteristics between RNA positive and negative patients clinically diagnosed with 2019 novel coronavirus pneumonia]. *Zhonghua Jie He He Hu Xi Za Zhi*. 2020;43(0):E023. doi:10.3760/cma.j.cn112147-20200214-00095
112. Gross A, Thiemig D, Koch FW, Schwarz M, Glaser S, Albrecht T. CT appearance of severe, laboratory-proven coronavirus disease 2019 (COVID-19) in a Caucasian patient in Berlin, Germany. *Rofo*. 2020. doi:10.1055/a-1138-8783
113. Holshue ML, DeBolt C, Lindquist S, et al. First case of 2019 novel coronavirus in the United

- States. *N Engl J Med*. 2020;382(10):929-936. doi:10.1056/NEJMoa2001191
114. Li K, Fang Y, Li W, et al. CT image visual quantitative evaluation and clinical classification of coronavirus disease (COVID-19). *Eur Radiol*. 2020. doi:10.1007/s00330-020-06817-6
  115. Fang Y, Zhang H, Xu Y, Xie J, Pang P, Ji W. CT Manifestations of Two Cases of 2019 Novel Coronavirus (2019-nCoV) Pneumonia. *Radiology*. 2020;295(1):208-209. doi:10.1148/radiol.2020200280
  116. Sun D, Li H, Lu XX, et al. Clinical features of severe pediatric patients with coronavirus disease 2019 in Wuhan: a single center's observational study. *World J Pediatr*. 2020. doi:10.1007/s12519-020-00354-4
  117. Zhang S, Li H, Huang S, You W, Sun H. High-resolution CT features of 17 cases of Corona Virus Disease 2019 in Sichuan province, China. *Eur Respir J*. 2020. doi:10.1183/13993003.00334-2020
  118. Hill DKJ, Russell DCD, Clifford DS, et al. The index case of SARS-CoV-2 in Scotland: a case report. *J Infect*. 2020. doi:10.1016/j.jinf.2020.03.022
  119. Hu Z, Song C, Xu C, et al. Clinical characteristics of 24 asymptomatic infections with COVID-19 screened among close contacts in Nanjing, China. *Sci China Life Sci*. 2020. doi:10.1007/s11427-020-1661-4
  120. Guan CS, Lv ZB, Yan S, et al. Imaging Features of Coronavirus disease 2019 (COVID-19): Evaluation on Thin-Section CT. *Acad Radiol*. 2020. doi:10.1016/j.acra.2020.03.002
  121. Zhou Y, Yang GD, Feng K, et al. [Clinical features and chest CT findings of coronavirus disease 2019 in infants and young children]. *Zhongguo Dang Dai Er Ke Za Zhi*. 2020;22(3):215-220.
  122. Liao X, Yang H, Kong J, Yang H. Chest CT Findings in a Pregnant Patient with 2019 Novel Coronavirus Disease. *Balk Med J*. 2020. doi:10.4274/balkanmedj.galenos.2020.2020.3.89
  123. Wang X, Zhou Z, Zhang J, Zhu F, Tang Y, Shen X. A case of 2019 Novel Coronavirus in a pregnant woman with preterm delivery. *Clin Infect Dis*. 2020. doi:10.1093/cid/ciaa200
  124. Wen R, Sun Y, Xing QS. A patient with SARS-CoV-2 infection during pregnancy in Qingdao, China. *J Microbiol Immunol Infect*. 2020. doi:10.1016/j.jmii.2020.03.004
  125. Wu J, Liu J, Zhao X, et al. Clinical Characteristics of Imported Cases of COVID-19 in Jiangsu Province: A Multicenter Descriptive Study. *Clin Infect Dis*. 2020. doi:10.1093/cid/ciaa199
  126. Yang W, Cao Q, Qin L, et al. Clinical characteristics and imaging manifestations of the 2019 novel coronavirus disease (COVID-19): A multi-center study in Wenzhou city, Zhejiang, China. *J Infect*. 2020. doi:10.1016/j.jinf.2020.02.016
  127. Gautret P, Lagier JC, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int J Antimicrob Agents*. 2020:105949. doi:10.1016/j.ijantimicag.2020.105949
  128. Bai SL, Wang JY, Zhou YQ, et al. Analysis of the first cluster of cases in a family of novel

- coronavirus pneumonia in Gansu Province. *Zhonghua Yu Fang Yi Xue Za Zhi*. 2020;54:E005. doi:10.3760/cma.j.issn.0253-9624.2020.0005
129. Zheng F, Liao C, Fan QH, et al. Clinical Characteristics of Children with Coronavirus Disease 2019 in Hubei, China. *Curr Med Sci*. 2020. doi:10.1007/s11596-020-2172-6
  130. Ruan ZR, Gong P, Han W, Huang MQ, Han M. A case of 2019 novel coronavirus infected pneumonia with twice negative 2019-nCoV nucleic acid testing within 8 days. *Chin Med J (Engl)*. 2020. doi:10.1097/CM9.0000000000000788
  131. Chang, Lin M, Wei L, et al. Epidemiologic and Clinical Characteristics of Novel Coronavirus Infections Involving 13 Patients Outside Wuhan, China. *JAMA*. 2020. doi:10.1001/jama.2020.1623
  132. Pan Y, Guan H, Zhou S, et al. Initial CT findings and temporal changes in patients with the novel coronavirus pneumonia (2019-nCoV): a study of 63 patients in Wuhan, China. *Eur Radiol*. 2020. doi:10.1007/s00330-020-06731-x
  133. Lui K, Fang YY, Deng Y, et al. Clinical characteristics of novel coronavirus cases in tertiary hospitals in Hubei Province. *Chin Med J (Engl)*. 2020. doi:10.1097/CM9.0000000000000744
  134. Sun Y, Koh V, Marimuthu K, et al. Epidemiological and Clinical Predictors of COVID-19. *Clin Infect Dis*. 2020. doi:10.1093/cid/ciaa322
  135. Ji LN, Chao S, Wang YJ, et al. Clinical features of pediatric patients with COVID-19: a report of two family cluster cases. *World J Pediatr*. 2020. doi:10.1007/s12519-020-00356-2
  136. Xia H, Zhao S, Wu Z, Luo H, Zhou C, Chen X. Emergency Caesarean delivery in a patient with confirmed coronavirus disease 2019 under spinal anaesthesia. *Br J Anaesth*. 2020. doi:10.1016/j.bja.2020.02.016
  137. Han R, Huang L, Jiang H, Dong J, Peng H, Zhang D. Early Clinical and CT Manifestations of Coronavirus Disease 2019 (COVID-19) Pneumonia. *AJR Am J Roentgenol*. 2020;1-6. doi:10.2214/AJR.20.22961
  138. Guan WJ, Ni ZY, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. 2020. doi:10.1056/NEJMoa2002032
  139. Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients with 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA - J Am Med Assoc*. 2020;323(11):1061-1069. doi:10.1001/jama.2020.1585
  140. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395(10223):507-513. doi:10.1016/S0140-6736(20)30211-7
  141. Yoon SH, Lee KH, Kim JY, et al. Chest Radiographic and CT Findings of the 2019 Novel Coronavirus Disease (COVID-19): Analysis of Nine Patients Treated in Korea. *Korean J Radiol*. 2020;21(4):494-500. doi:10.3348/kjr.2020.0132
  142. Guan W, Liu J, Yu C. CT Findings of Coronavirus Disease (COVID-19) Severe Pneumonia. *AJR Am*



- J Roentgenol.* 2020;W1-w2. doi:10.2214/ajr.20.23035
143. Yao Y, Tian Y, Zhou J, Ma X, Yang M, Wang S. Epidemiological characteristics of 2019-nCoV infections in Shaanxi, China by February 8, 2020. *Eur Respir J.* 2020. doi:10.1183/13993003.00310-2020
  144. Duan YN, Qin J. Pre- and Posttreatment Chest CT Findings: 2019 Novel Coronavirus (2019-nCoV) Pneumonia. *Radiology.* 2020;295(1):21. doi:10.1148/radiol.202000323
  145. Zhang YH, Lin DJ, Xiao MF, et al. 2019-novel coronavirus infection in a three-month-old baby. *Zhonghua er ke za zhi = Chinese J Pediatr.* 2020;58:E006. doi:10.3760/cma.j.issn.0578-1310.2020.0006
  146. Liu H, Liu F, Li J, Zhang T, Wang D, Lan W. Clinical and CT imaging features of the COVID-19 pneumonia: Focus on pregnant women and children. *J Infect.* 2020. doi:10.1016/j.jinf.2020.03.007
  147. Liu K, Chen Y, Lin R, Han K. Clinical feature of COVID-19 in elderly patients: a comparison with young and middle-aged patients. *J Infect.* 2020. doi:10.1016/j.jinf.2020.03.005
  148. Tian S, Hu N, Lou J, et al. Characteristics of COVID-19 infection in Beijing. *J Infect.* 2020. doi:10.1016/j.jinf.2020.02.018
  149. Cheng JL, Huang C, Zhang GJ, et al. [Epidemiological characteristics of novel coronavirus pneumonia in Henan]. *Zhonghua Jie He He Hu Xi Za Zhi.* 2020;43(0):E027. doi:10.3760/cma.j.cn112147-20200222-00148
  150. Yu P, Zhu J, Zhang Z, Han Y, Huang L. A familial cluster of infection associated with the 2019 novel coronavirus indicating potential person-to-person transmission during the incubation period. *J Infect Dis.* 2020. doi:10.1093/infdis/jiaa077
  151. Xia W, Shao J, Guo Y, Peng X, Li Z, Hu D. Clinical and CT features in pediatric patients with COVID-19 infection: Different points from adults. *Pediatr Pulmonol.* 2020. doi:10.1002/ppul.24718
  152. Wang S, Guo L, Chen L, et al. A case report of neonatal COVID-19 infection in China. *Clin Infect Dis.* 2020. doi:10.1093/cid/ciaa225
  153. Liu Q, Wang RS, Qu GQ, et al. Gross examination report of a COVID-19 death autopsy. *Fa Yi Xue Za Zhi.* 2020;36(1):21-23. doi:10.12116/j.issn.1004-5619.2020.01.005
  154. Van Cuong L, Giang HTN, Linh LK, et al. The first Vietnamese case of COVID-19 acquired from China. *Lancet Infect Dis.* 2020. doi:10.1016/S1473-3099(20)30111-0
  155. Zhang X, Cai H, Hu J, et al. Epidemiological, clinical characteristics of cases of SARS-CoV-2 infection with abnormal imaging findings. *Int J Infect Dis.* 2020. doi:10.1016/j.ijid.2020.03.040
  156. Qian GQ, Yang NB, Ding F, et al. Epidemiologic and Clinical Characteristics of 91 Hospitalized Patients with COVID-19 in Zhejiang, China: A retrospective, multi-centre case series. *QJM.* 2020. doi:10.1093/qjmed/hcaa089



157. Cheng SC, Chang YC, Fan Chiang YL, et al. First case of Coronavirus Disease 2019 (COVID-19) pneumonia in Taiwan. *J Formos Med Assoc.* 2020;119(3):747-751. doi:10.1016/j.jfma.2020.02.007
158. Fan Q, Pan Y, Wu Q, et al. Anal swab findings in an infant with COVID-19. *Pediatr Investig.* 2020;4(1):48-50. doi:10.1002/ped4.12186
159. Albarello F, Pianura E, Di Stefano F, et al. 2019-novel Coronavirus severe adult respiratory distress syndrome in two cases in Italy: An uncommon radiological presentation. *Int J Infect Dis.* 2020;93:192-197. doi:10.1016/j.ijid.2020.02.043
160. Bernard Stoecklin S, Rolland P, Silue Y, et al. First cases of coronavirus disease 2019 (COVID-19) in France: surveillance, investigations and control measures, January 2020. *Euro Surveill.* 2020;25(6). doi:10.2807/1560-7917.Es.2020.25.6.2000094
161. Chen T, Wu D, Chen H, et al. Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *Bmj.* 2020;368:m1091. doi:10.1136/bmj.m1091
162. Zhao X, Liu B, Yu Y, et al. The characteristics and clinical value of chest CT images of novel coronavirus pneumonia. *Clin Radiol.* 2020. doi:10.1016/j.crad.2020.03.002
163. Park JY, Han MS, Park KU, Kim JY, Choi EH. First Pediatric Case of Coronavirus Disease 2019 in Korea. *J Korean Med Sci.* 2020;35(11):e124. doi:10.3346/jkms.2020.35.e124
164. Song F, Shi N, Shan F, et al. Emerging 2019 Novel Coronavirus (2019-nCoV) Pneumonia. *Radiology.* 2020;295(1):210-217. doi:10.1148/radiol.202000274
165. Ren LL, Wang YM, Wu ZQ, et al. Identification of a novel coronavirus causing severe pneumonia in human: a descriptive study. *Chin Med J (Engl).* 2020. doi:10.1097/CM9.0000000000000722
166. <https://www.worldometers.info/coronavirus/>.
167. World Health Organization. WHO Corona Virus Situation Report. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/>. Published 2020.
168. Day M. Covid-19: four fifths of cases are asymptomatic, China figures indicate. *BMJ.* 2020. doi:10.1136/bmj.m1375
169. Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. *Eurosurveillance.* 2020. doi:10.2807/1560-7917.Es.2020.25.10.2000180
170. Day M. Covid-19: identifying and isolating asymptomatic people helped eliminate virus in Italian village. *BMJ.* 2020. doi:10.1136/bmj.m1165
171. Verity R, Okell LC, Dorigatti I, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis.* 2020. doi:10.1016/s1473-3099(20)30243-7
172. Ruan S. Likelihood of survival of coronavirus disease 2019. *Lancet Infect Dis.* 2020. doi:10.1016/s1473-3099(20)30257-7

173. Montecino-Rodriguez E, Berent-Maoz B, Dorshkind K. Causes, consequences, and reversal of immune system aging. *J Clin Invest*. 2013. doi:10.1172/JCI64096
174. Fuentes E, Fuentes M, Alarcón M, Palomo I. Immune system dysfunction in the elderly. *An Acad Bras Cienc*. 2017. doi:10.1590/0001-3765201720160487
175. Opal SM, Girard TD, Ely EW. The Immunopathogenesis of Sepsis in Elderly Patients. *Clin Infect Dis*. 2005. doi:10.1086/432007
176. Smits SL, De Lang A, Van Den Brand JMA, et al. Exacerbated innate host response to SARS-CoV in aged non-human primates. *PLoS Pathog*. 2010. doi:10.1371/journal.ppat.1000756
177. Jin J-M, Bai P, He W, et al. Gender differences in patients with COVID-19: Focus on severity and mortality. *medRxiv*. 2020. doi:10.1101/2020.02.23.20026864
178. Wenham C, Smith J, Morgan R. COVID-19: the gendered impacts of the outbreak. *Lancet*. 2020. doi:10.1016/S0140-6736(20)30526-2
179. Delanghe JR, Speeckaert MM, De Buyzere ML. The host's angiotensin-converting enzyme polymorphism may explain epidemiological findings in COVID-19 infections. *Clin Chim Acta*. 2020. doi:10.1016/j.cca.2020.03.031
180. Liu S, Zhang M, Yang L, et al. Prevalence and patterns of tobacco smoking among Chinese adult men and women: Findings of the 2010 national smoking survey. *J Epidemiol Community Health*. 2017. doi:10.1136/jech-2016-207805
181. Judah G, Aunger R, Schmidt WP, Michie S, Granger S, Curtis V. Experimental pretesting of hand-washing interventions in a natural setting. *Am J Public Health*. 2009. doi:10.2105/AJPH.2009.164160
182. Nordström A, Hadrévi J, Olsson T, Franks PW, Nordström P. Higher prevalence of type 2 diabetes in men than in women is associated with differences in visceral fat mass. *J Clin Endocrinol Metab*. 2016. doi:10.1210/jc.2016-1915
183. Regitz-Zagrosek V, Lehmkuhl E, Weickert MO. Gender differences in the metabolic syndrome and their role for cardiovascular disease. *Clin Res Cardiol*. 2006. doi:10.1007/s00392-006-0377-8
184. Gupte M, Thatcher SE, Boustany-Kari CM, et al. Angiotensin converting enzyme 2 contributes to sex differences in the development of obesity hypertension in C57BL/6 mice. *Arterioscler Thromb Vasc Biol*. 2012. doi:10.1161/ATVBAHA.112.248559
185. Asselta R, Paraboschi EM, Mantovani A, Duga S. ACE2 and TMPRSS2 Variants and Expression as Candidates to Sex and Country Differences in COVID-19 Severity in Italy. *SSRN Electron J*. 2020. doi:10.2139/ssrn.3559608
186. Channappanavar R, Fett C, Mack M, Ten Eyck PP, Meyerholz DK, Perlman S. Sex-Based Differences in Susceptibility to Severe Acute Respiratory Syndrome Coronavirus Infection. *J Immunol*. 2017. doi:10.4049/jimmunol.1601896
187. Lin C, Ding Y, Xie B, et al. Asymptomatic novel coronavirus pneumonia patient outside Wuhan:

- The value of CT images in the course of the disease. *Clin Imaging*. 2020. doi:10.1016/j.clinimag.2020.02.008
188. Rodelo JR, De La Rosa G, Valencia ML, et al. D-dimer is a significant prognostic factor in patients with suspected infection and sepsis. *Am J Emerg Med*. 2012. doi:10.1016/j.ajem.2012.04.033
  189. Dinarello CA. Proinflammatory cytokines. *Chest*. 2000. doi:10.1378/chest.118.2.503
  190. Short KR, Veeris R, Leijten LM, et al. Proinflammatory Cytokine Responses in Extra-Respiratory Tissues during Severe Influenza. *J Infect Dis*. 2017. doi:10.1093/infdis/jix281
  191. Kube D, Sontich U, Fletcher D, Davis PB. Proinflammatory cytokine responses to *P. aeruginosa* infection in human airway epithelial cell lines. *Am J Physiol - Lung Cell Mol Physiol*. 2001. doi:10.1152/ajplung.2001.280.3.l493
  192. Toliver-Kinsky T, Kobayashi M, Suzuki F, Sherwood ER. The systemic inflammatory response syndrome. In: *Total Burn Care: Fifth Edition*. ; 2018. doi:10.1016/B978-0-323-47661-4.00019-8
  193. Xiong T-Y, Redwood S, Prendergast B, Chen M. Coronaviruses and the cardiovascular system: acute and long-term implications. *Eur Heart J*. 2020. doi:10.1093/eurheartj/ehaa231
  194. Warny M, Helby J, Nordestgaard BG, Birgens H, Bojesen SE. Lymphopenia and risk of infection and infection-related death in 98,344 individuals from a prospective Danish population-based study. *PLoS Med*. 2018. doi:10.1371/journal.pmed.1002685
  195. Minatoguchi S, Nomura A, Imaizumi T, et al. Low serum albumin as a risk factor for infection-related in-hospital death among hemodialysis patients hospitalized on suspicion of infectious disease: a Japanese multicenter retrospective cohort study. *Ren Replace Ther*. 2018. doi:10.1186/s41100-018-0173-8
  196. Arabi YM, Mandourah Y, Al-Hameed F, et al. Corticosteroid therapy for critically ill patients with middle east respiratory syndrome. *Am J Respir Crit Care Med*. 2018. doi:10.1164/rccm.201706-1172OC
  197. Russell CD, Millar JE, Baillie JK. Clinical evidence does not support corticosteroid treatment for 2019-nCoV lung injury. *Lancet*. 2020. doi:10.1016/S0140-6736(20)30317-2
  198. Zumla A, Hui DS, Perlman S. Middle East respiratory syndrome. *Lancet*. 2015. doi:10.1016/S0140-6736(15)60454-8
  199. Rodrigo C, Leonardi-Bee J, Nguyen-Van-Tam JS, Lim WS. Effect of corticosteroid therapy on influenza-related mortality: A systematic review and meta-analysis. *J Infect Dis*. 2015. doi:10.1093/infdis/jiu645

## Figure legend

### Figure 1. Flow-chart of the search strategy.

A total of 148 studies were eligible for the literature review and the first part of the meta-analysis (i.e., prevalence). Nineteen studies were included in the second part of the meta-analysis (i.e., severity and mortality).

### Figure 2. Proportion of female and male patients in adult (A) and pediatric/neonatal cohort (B).

All case studies/ reports were pooled together for visualization (CS\_adult, and CS\_children [pediatric/neonatal]). The key to the study identifier can be found in Tables 1 (adults) and Table 3 (children).

### Figure 3. Age of adult (A), pregnant (B), and pediatric/neonatal (C) COVID-19 patients included in eligible studies.

Median age and interquartile ranges (IQR) are represented by the midpoints and error bars, respectively. The studies have been sorted by patients' median age in years. The size of the midpoint (circle, square, triangle) indicates the study sample size. The red line indicates the pooled median age of the respective cohort. All adult case studies/ reports (CS\_adult) were pooled for the visualization reasons. The key to the study identifier can be found in Table 1 (adults), Table 2 (pregnant women), and Table 3 (children).

### Figure 4. Comorbidities (A), clinical signs and symptoms (B), outcomes (C), and treatments administered (D) to adult COVID-19 patients.

The colors indicated the proportion of patients (%; 0 = yellow, 100 = dark purple). Note: Missing values are colored in white.

### Figure 5. Age of non-severe (A), severe (B), survivor (C), and non-survivor (D) COVID-19 patients included in eligible studies.

The median age and Interquartile ranges (IQR) are represented by the midpoints and error bars, respectively. The studies have been sorted by patients' median age in years. The size of the midpoint indicates the study sample size. The red line indicates the pooled median age of the respective cohort. The key to the study identifier can be found in Table 1.

### Figure 6. Relative risks of comorbidities (i.e., hypertension, diabetes mellitus, and COPD) and complications (i.e., ARDS) in patients with a severe COVID-19 disease progression.

Funnel plots indicate the potential of publication bias. The key to the study identifier can be found in Table 1.

### Figure 7. Relative risks of comorbidity (i.e., any heart condition), treatment (i.e., mechanical ventilation), and complications (i.e., acute kidney injury and ARDS) in survivors and non-survivors.

Funnel plots indicate the potential of publication bias. The key to the study identifier can be found in Table 1.

**SUPPLEMENTARY FILE****APPENDICES**

- **Appendix 1:** Full Search Strategy
- **Appendix 2:** Sensitivity analyses in the adult population

**SUPPLEMENTARY TABLES**

- **Supplementary Table 1.** List of extracted variables
- **Supplementary Table 2.** Classification of patients in severe and non-severe COVID-19 cohort
- **Supplementary Table 3.** Number of studies performed in each country
- **Supplementary Table 4.** Laboratory findings reported for adult and pregnant COVID-19 patients
- **Supplementary Table 5.** Results of meta-analysis of prevalence in pregnant COVID-19 patients
- **Supplementary Table 6.** Results of meta-analysis of prevalence in pediatric and neonatal COVID-19 patients
- **Supplementary Table 7.** Laboratory findings reported for pediatric/neonatal COVID-19 patients

**SUPPLEMENTARY FIGURES**

- **Supplementary Figure 1.** All comorbidities in adult COVID-19 patients.
- **Supplementary Figure 2.** All imaging features in adult COVID-19 patients.
- **Supplementary Figure 3.** Symptoms and signs in pregnant COVID-19 patients.
- **Supplementary Figure 4.** Imaging features in pregnant COVID-19 patients.
- **Supplementary Figure 5.** Treatments administered to pregnant COVID-19 patients.
- **Supplementary Figure 6.** Symptoms and signs in pediatric and neonatal COVID-19 patients.
- **Supplementary Figure 7.** Imaging features in pediatric and neonatal COVID-19 patients.
- **Supplementary Figure 8.** Treatments administered to pediatric and neonatal COVID-19 patients.
- **Supplementary Figure 9.** Outcomes of pediatric and neonatal COVID-19 patients.
- **Supplementary Figure 10.** Proportion of male and female patients in the non-severe, severe, survivors, and non-survivor patient groups.

## APPENDIX

### Appendix 1: Full Search Strategy

#### Keywords

- COVID-19
- coronavirus disease 2019
- severe acute respiratory syndrome coronavirus 2
- SARS-CoV-2
- 2019 novel coronavirus
- 2019-nCoV
- coronavirus
- corona virus

Literature search performed on March 28<sup>th</sup>, 2020:

- **EMBASE – String**  
 #1 AND (2019:py OR 2020:py)  
 ('covid 19' OR (coronavirus AND disease AND 2019) OR (severe AND acute AND respiratory AND syndrome AND coronavirus AND 2) OR 'sars cov 2' OR (2019 AND novel AND coronavirus) OR '2019 ncov' OR coronavirus OR (corona AND virus)) AND [1-12-2019]/sd NOT [29-3-2020]/sd
- **Pubmed – String**  
 (((((((("COVID-19"[All Fields] OR "severe acute respiratory syndrome coronavirus 2"[Supplementary Concept] OR "severe acute respiratory syndrome coronavirus 2"[All Fields] OR "2019-nCoV"[All Fields] OR "SARS-CoV-2"[All Fields] OR "2019nCoV"[All Fields] OR ("Wuhan"[All Fields] AND ("coronavirus"[MeSH Terms] OR "coronavirus"[All Fields])) AND 2019/12[PDAT] : 2030[PDAT])) OR ("COVID-19"[Supplementary Concept] OR "COVID-19"[All Fields] OR "coronavirus disease 2019"[All Fields])) OR ("severe acute respiratory syndrome coronavirus 2"[Supplementary Concept] OR "severe acute respiratory syndrome coronavirus 2"[All Fields])) OR ("severe acute respiratory syndrome coronavirus 2"[Supplementary Concept] OR "severe acute respiratory syndrome coronavirus 2"[All Fields] OR "sars cov 2"[All Fields])) OR ("severe acute respiratory syndrome coronavirus 2"[Supplementary Concept] OR "severe acute respiratory syndrome coronavirus 2"[All Fields] OR "2019 novel coronavirus"[All Fields])) OR ("severe acute respiratory syndrome coronavirus 2"[Supplementary Concept] OR "severe acute respiratory syndrome coronavirus 2"[All Fields] OR "2019 ncov"[All Fields])) OR ("coronavirus"[MeSH Terms] OR "coronavirus"[All Fields])) OR (corona[All Fields] AND ("viruses"[MeSH Terms] OR "viruses"[All Fields] OR "virus"[All Fields])) OR ("severe acute respiratory syndrome coronavirus 2"[Supplementary Concept] OR "severe acute respiratory syndrome coronavirus 2"[All Fields] OR "2019 novel coronavirus"[All Fields]) AND ("2019/12/01"[PDAT] : "3000/12/31"[PDAT]))

- **Scopus – String**  
covid-19 OR coronavirus AND disease 2019 OR severe AND acute AND respiratory AND syndrome AND coronavirus 2 OR sars-cov-2 OR 2019 novel AND coronavirus OR 2019-ncov OR coronavirus OR corona AND virus AND ( LIMIT-TO ( PUBYEAR , 2020 ) OR LIMIT-TO ( PUBYEAR , 2019 ) )
- **Web of Science – String**  
ALL FIELDS: (COVID-19 OR coronavirus disease 2019 OR severe acute respiratory syndrome coronavirus 2 OR SARS-CoV-2 OR 2019 novel coronavirus OR 2019-nCoV OR coronavirus OR corona virus) Indexes=SCI-EXPANDED, SSCI, A&HCI, CPCI-S, CPCI-SSH, BKCI-S, BKCI-SSH, ESCI, CCR-EXPANDED, IC Timespan=2019-2020

## Appendix 2: Sensitivity analyses in the adult population

To assess the sensitivity with respect to studies that only reported a mean and standard deviation, we performed additional probabilistic simulations. Specifically, we assumed that mean and standard deviation were calculated from a skewed normal distribution with unknown shape parameter  $\alpha$  (where  $\alpha=0$  corresponds to the standard normal distribution). We then calculated the expectation of the median and the respective IQR under the assumption that  $\alpha \in (-\infty, 0]$ , i.e., a left-skewed distribution, and under the assumption that  $\alpha \in [0, \infty)$ , i.e., a right-skewed distribution. Both of these assumptions are the “agnostic” ones in the sense that we consider skewness, but do not assume boundedness. We find the overall median and IQR (of all studies) to be unaffected when including the results of these simulations, indicating that our analyses are stable.

Variable	All studies that reported mean and SD (assumption normal distribution)	All studies All studies that reported mean and SD (assumption left-skewed distribution)	All studies that reported mean and SD (assumption right-skewed distribution)
Age, years	49.50 [20.25]	52.33 [20.23]	46.67 [20.70]
Leucocytes	5.48 [3.24]	5.89 [3.24]	5.08 [3.31]
Lymphocytes	1.21 [0.76]	1.33 [0.76]	1.08 [0.77]
Neutrophils	3.29 [2.31]	3.63 [2.31]	2.98 [2.36]
Aspartate transaminase (AST)	30.08 [24.16]	34.17 [24.15]	25.99 [24.70]
Alanine aminotransferase (ALT)	29.37 [34.71]	34.53 [34.68]	24.21 [35.47]
Interleukin 6 (IL-6)	3.6 [0.67]	-	3.25 [0.75]
D-dimer	3.36 [0.73]	3.47 [0.73]	3.25 [0.75]

**Supplementary Table 1:** List of extracted variables

Category	Variable Name	Remark
Publication information	First Author Name, Year of Publication	
	Title	
	PMID or DOI	DOI: Digital object identifier; PMID: Pubmed ID
	Country	
	Language of article	Original language of article
Publication information	Study population	Adults, pregnant women, children/neonates
	Study type	Case study, case series (study with more than 1 patient providing data on patient level), cohort study (study with more than 1 patient providing data on group level)
Demographics	Sex	Count
	Age [years]	Median [first and third quartiles], mean (standard deviation)
Comorbidities	Any comorbidities	Count
	Aorta sclerosis	Count
	Arrhythmia	Count
	Asthma	Count
	Autoimmune disease	Count
	Carcinoma	Count
	Cardio cerebrovascular disease	Count
	Cardiovascular disease	Count
	Cerebrovascular disease	Count
	Cholecystitis	Count
	Cholelithiasis	Count
	Chronic lung disease	Count
	Chronic gastritis, gastric ulcer	Count
	Chronic kidney disease	Count
	Chronic liver disease	Count
	Chronic pharyngitis	Count
	Chronic renal insufficiency	Count
	Cirrhosis	Count
	COPD	Count
	Coronary heart disease	Count
	Dementia	Count
	Diabetes mellitus	Count
	Digestive system diseases	Count
	Drug hypersensitivity	Count
	Electrolyte imbalance	Count
	Endocrine diseases	Count
	End-stage kidney disease	Count



	Fatty liver	Count
	Food allergy	Count
	Gastrectomy	Count
	Heart failure (congestive, chronic)	Count
	Hepatitis B	Count
	Hepatitis C	Count
	HIV infection	Count
	Hypothyroidism	Count
	Hyperlipidemia	Count
	Hypertension	Count
	Hyperuricemia	Count
	Hypothyroidism	Count
	Immunodeficiency	Count
	Influenza	Count
	Liver disease	Count
	Malnutrition	Count
	Nervous system disease	Count
	Obstructive sleep apnea	Count
	Organ transplant	Count
	Respiratory system diseases	Count
	Rheumatologic disease	Count
	Secondary pulmonary	Count
	Tuberculosis	Count
	Current smoking status: No	Count
	Current smoking status: Yes	Count
	Stroke	Count
	Transplants	Count
	Thyroid disease	Count
	Urolithiasis	Count
	Urticaria	Count
<b>Signs and symptoms</b>	Abdominal pain	Count
	Anorexia	Count
	Anosmia	Count
	Belching	Count
	Chest pain	Count
	Chest tightness	Count
	Conjunctival congestion	Count
	Constipation	Count
	Cough	Count
	Diarrhea	Count
	Dizziness or confusion	Count
	Dysgeusia	Count
	Dysosmia	Count
	Dyspnea	Count
	Emesis / vomiting	Count
	Fatigue	Count
	Fever	Count
	Headache	Count
	Hemoptysis	Count

	Myalgia	Count
	Nasal congestion	Count
	Nausea	Count
	Heart palpitation	Count
	Rash	Count
	Running nose/ rhinorrhea	Count
	Sinusitis	Count
	Sore throat	Count
	Sputum	Count
Imaging features	Air bronchogram	Count
	Atelectasis	Count
	Bilateral patchy shadowing	Count
	Bilateral pneumonia	Count
	Bilateral pulmonary infiltration	Count
	Bronchiectasis	Count
	Bulla	Count
	Consolidation	Count
	Crazy paving pattern	Count
	Focal consolidation	Count
	Ground glass opacity (GGO)	Count
	GGO with consolidation	Count
	Interstitial abnormalities	Count
	Local patchy shadowing	Count
	Nodular lesions	Count
	Pathological findings are not further specified	Count
	Pathologic CT findings	
	Peribronchial thickening	Count
	Pleural effusion	Count
	Pneumonia	Count
	Pulmonary edema	Count
	Radiology reported	Count
	Reticulation/ thickened interlobular septa	Count
	Stripe	Count
	Subpleural opacities	Count
	Thickened intralobular septa	Count
	Thickened pleura	Count
	Unilateral pneumonia	Count
	Vascular thickening	Count
	Venous congestion	Count
Laboratory parameters	<b>Hematology</b>	Median [IQR] or mean (SD)
	Hemoglobin, g/L	Median [IQR] or mean (SD)
	Leucocyte count, $\times 10^9/L$	Median [IQR] or mean (SD)
	Lymphocyte count, $\times 10^9/L$	Median [IQR] or mean (SD)
	Neutrophils count, $\times 10^9/L$	Median [IQR] or mean (SD)
	Platelet count, $\times 10^9/L$	Median [IQR] or mean (SD)
	<b>Coagulation</b>	Median [IQR] or mean (SD)
	Prothrombin time, s	Median [IQR] or mean (SD)

Treatments	Activated partial thromboplastin time, s	Median [IQR] or mean (SD)
	Fibrinogen, g/L	Median [IQR] or mean (SD)
	<b>Electrolytes</b>	Median [IQR] or mean (SD)
	Sodium, mmol/L	Median [IQR] or mean (SD)
	Potassium, mmol/L	Median [IQR] or mean (SD)
	<b>Clinical chemistry labs</b>	Median [IQR] or mean (SD)
	Albumin, g/L	Median [IQR] or mean (SD)
	Alanine aminotransferase (ALT), U/L	Median [IQR] or mean (SD)
	Aspartate transaminase (AST), U/L	Median [IQR] or mean (SD)
	Total bilirubin, mmol/L	Median [IQR] or mean (SD)
	Lactate dehydrogenase, U/L	Median [IQR] or mean (SD)
	Creatine kinase, U/L	Median [IQR] or mean (SD)
	D-dimer, µg/mL	Median [IQR] or mean (SD)
	Creatinine, µmol/L	Median [IQR] or mean (SD)
	Blood urea nitrogen, mmol/L	Median [IQR] or mean (SD)
	<b>Inflammation</b>	Median [IQR] or mean (SD)
	C-reactive protein, mg/L	Median [IQR] or mean (SD)
	Interleukin 6, pg/mL	Median [IQR] or mean (SD)
	Procalcitonin, ng/mL	Median [IQR] or mean (SD)
	Erythrocyte sedimentation rate, mm/h	Median [IQR] or mean (SD)
	<b>Antibiotic treatment</b>	
	Any antibiotic	Not further specified, count
	Amoxicillin	Count
	Azithromycin	Count
	Carbapenem linezolid	Count
	Cefaclor	Count
	Cefazolin	Count
	Cefepime	Count
	Cefotiam hydrochloride	Count
	Ceftazidime	Count
	Ceftriaxone	Count
	Clavulanate	Count
	Levofloxacin	Count
	Moxifloxacin hydrochloride	Count
	Ornidazole	Count
	Piperacillin tazobactam	Count
	Sulbactam	Count
	Teicoplanin	Count
	Vancomycin	Count
	<b>Antiviral treatment</b>	
	Any antiviral	Not further specified, count
	Ganciclovir	Count
	Lopinavir	Count
	Oseltamivir	Count

	Remdesivir	Count
	Ribavirin	Count
	Ritonavir	Count
	Veletonavir	Count
	<b>Oxygen therapy</b>	
	Any oxygen therapy	Not further specified, count
	High flow nasal cannula	Count
	All mechanical ventilation together	Count
	Non-invasive mechanical ventilation	Count
	Invasive mechanical ventilation	Count
	ECMO	Count
	<b>Other</b>	
	Alpha interferon aerosol inhalation	Count
	Antifungal treatment	Count
	Corticosteroids	Count
	Hydroxychloroquine	Count
	Immune enhancing treatment	Count
	Intravenous immunoglobulin	Count
	Renal replacement therapy	Count
	Thymalfasin	Count
Outcome	Died	Count
	Survived	Count
	Discharged	Count
	Hospitalized	Count
	Recovery	Count
	Admission to ICU	Count
Complications	Acidosis	Count
	Acute cardiac injury	Count
	Acute liver injury	Count
	Acute liver or kidney injury	Count
	Acute kidney injury	Count
	Acute respiratory distress syndrome (ARDS)	Count
	Acute respiratory injury	Count
	Arrhythmia	Count
	Coagulopathy	Count
	Disseminated intravascular coagulation	Count
	Gastrointestinal hemorrhage	Count
	Heart Failure	Count
	Hyperglycemia	Count
	Hypoproteinemia	Count
	Pneumonia	Count
	Pneumothorax	Count
	Respiratory failure	Count

Rhabdomyolysis	Count
Secondary infections	Count
any	Count
bacterial	Count
fungal	Count
viral	Count
Sepsis	Count
Septic shock	Count
Others	
Hospital length of stay, days	Median [IQR] or mean (SD)
Time from symptom onset to admission, days	Median [IQR] or mean (SD)

**Supplementary Table 2:** Classification of patients in severe and non-severe CoVID-19 cohort

Author / year	Study reference Title	Severe vs non-severe classification
Guan et al, 2020	DOI: 10.1056/NEJMoa2002032	In accordance with American Thoracic Society guidelines for community-acquired pneumonia (Metlay JP et al., Am J Respir Crit Care Med 2019; 200(7): e45-e67.)
Tian et al, 2020	32112886	<b>Non-severe:</b> A mild case was defined as a confirmed case with fever, respiratory symptoms and radiographic evidence of pneumonia. <b>Severe:</b> A severe case was defined as a mild case with dyspnea or respiratory failure.
Wang et al, 2020	32031570	<b>Non-severe:</b> patients were not admitted to ICU <b>Severe:</b> patients were admitted to ICU
Wang et al, 2020	32179910	(1) mild, mild clinical symptoms without pneumonia seen at chest computed tomography; (2) ordinary, fever and other respiratory symptoms with pneumonia seen at imaging; (3) severe, respiratory distress, hypoxia (oxygen saturation, $\leq 93\%$ ), or abnormal results of blood gas analysis ( $\text{PaO}_2 < 60 \text{ mm Hg}$ or $\text{PaCO}_2 > 50 \text{ mm Hg}$ ); and (4) critical, respiratory failure requiring mechanical ventilation, shock, or other organ failure requiring intensive care unit monitoring and treatment. <b>Non-severe:</b> mild and ordinary <b>Severe:</b> severe and critical
Wan et al, 2020	32198776	The mild group had mild clinical symptoms and no pneumonia on imaging. The normal group had symptoms of fever, respiratory tract symptoms, and imaging showed pneumonia. The severe group had respiratory distress, $\text{RR} \geq 30$ beats/minute in a resting state, a mean oxygen saturation of $\leq 93\%$ , and an arterial blood oxygen partial pressure ( $\text{PaO}_2$ )/oxygen concentration ( $\text{FiO}_2$ ) $\leq 300 \text{ mmHg}$ . The critical group had respiratory failure and required mechanical ventilation, the occurrence of shock, and the combined failure of other organs that required intensive care unit (ICU) monitoring and treatment. <b>Non-severe:</b> mild and normal <b>Severe:</b> severe and critical
Wang et al, 2020	32176772	<b>Non-severe:</b> $\text{SpO}_2 \geq 90\%$ <b>Severe:</b> $\text{SpO}_2 < 90\%$
Qian et al, 2020	32181807	Patients were divided into the diagnosed as severe group and mild group according to national treatment guideline. (National Health Commission of the People's Republic of China. The notice of launching guideline on diagnosis and treatment of the novel coronavirus pneumonia (4 <sup>th</sup> edition) <a href="http://www.nhc.gov.cn/yzygj/s7653p/202001/4294563ed35b43209b31739bdo785e67.shtml">http://www.nhc.gov.cn/yzygj/s7653p/202001/4294563ed35b43209b31739bdo785e67.shtml</a> and (5 <sup>th</sup> edition) <a href="http://www.nhc.gov.cn/yzygj/s7653p/202002/3b09b894ac9b4204a79db5b8912d4440.shtml">http://www.nhc.gov.cn/yzygj/s7653p/202002/3b09b894ac9b4204a79db5b8912d4440.shtml</a> )
Qu et al, 2020	32181903	Clinical condition assessment criteria: according to the COVID-19

		<p>diagnosis and treatment plan issued by the National Health Committee of China, clinical classification is performed, which is divided into four types: light, ordinary, heavy, and critical. Mild: mild clinical symptoms, no pneumonia on imaging; common: fever, respiratory tract symptoms, and pneumonia on imaging; severe: meet any of the following: (a) shortness of breath, RR &gt; 30 times/min; (b) resting state, means oxygen saturation <math>\leq 93\%</math>; (c) partial pressure of arterial oxygen (PaO<sub>2</sub>)/oxygen concentration (FiO<sub>2</sub>) <math>\leq 30</math> mmHg (1 mmHg = 0.133 kPa), pulmonary imaging shows that the lesions have progressed significantly within 50 to 48 hours, and those who are more than 50% are managed as heavy; critical: meet any of the following: (a) have respiratory failure and require mechanical ventilation; (b) have body weight; (c) combined with other organ failure requires ICU monitoring and treatment.</p> <p><b>Non-severe:</b> light and ordinary</p> <p><b>Severe:</b> heavy and critical</p>
Chen et al, 2020	32141280	<p><b>Non-severe:</b> The clinical symptoms are mild, and there are no pneumonia manifestations on imaging.</p> <p><b>Severe:</b> Meets any of the following: shortness of breath, respiratory rate (RR) &gt; 30 times / min; at rest, refers to oxygen saturation &lt; 93%; arterial blood oxygen partial pressure (PaO<sub>2</sub>) / oxygen concentration (FiO<sub>2</sub>) &lt; 300 mmHg (1 mmHg = 0.133 kPa), high altitude (altitude &gt; 1 000 m) area should be corrected according to the formula, namely PaO<sub>2</sub> / FiO<sub>2</sub> <math>\times</math> [atmospheric pressure (mmHg) / 760]; lung imaging shows 24 The progress of the lesion was &gt; 50% at ~ 48 h.</p>
Xu et al, 2020	32109443	<p>Infection by the National Health Commission (Trial Version 5), 6 the NCP was classified into four types: mild with slight clinical symptoms but no imaging presentations of pneumonia; common with fever, respiratory symptoms and imaging presentations of pneumonia; severe type with any of the following: respiratory distress with RR &gt; 30 times/minutes, oxygen saturation at rest &lt; 93%, or PaO<sub>2</sub>/FiO<sub>2</sub> &lt; 300 mmHg (1 mmHg = 0.133 kPa); critically severe type with any of the following: respiratory failure needing mechanical ventilation, shock, or combination with other organ failure needing ICU intensive care.</p> <p><b>Non-severe:</b> mild and common</p> <p><b>Severe:</b> severe and critically severe</p>
Zhang et al, 2020	32077115	<p>Severity of COVID-19 was defined according to the diagnostic and treatment guideline for SARS-CoV-2 issued by Chinese National Health Committee (version 3-5).</p> <p><b>Severe</b> COVID-19 was designated when the patients had one of the following criteria: (a) respiratory distress with respiratory frequency <math>\geq 30</math>/min; (b) pulse oximeter oxygen saturation <math>\leq 93\%</math> at rest; and (c) oxygenation index (artery partial pressure of oxygen/inspired oxygen fraction, PaO<sub>2</sub>/ FiO<sub>2</sub>) <math>\leq 300</math> mm Hg.</p>

		<b>Non-severe:</b> all other patients
Shi et al, 2020	32211816	<b>Non-severe:</b> without cardiac injury <b>Severe:</b> with cardiac injury Cardiac injury was defined as blood levels of cardiac biomarkers above the 99th-percentile upper reference limit, regardless of new abnormalities in electrocardiography and echocardiography.



**Supplementary Table 3.** Number of studies performed in each country

Country	Number of studies (%)
Canada	1 (0.68)
China	129 (87.2)
Europe*	1 (0.67)
France	2 (1.35)
Germany	1 (0.67)
Italy	1 (0.67)
Japan	1 (0.67)
Scotland	1 (0.67)
Singapore	2 (1.35)
South Korea	3 (2.03)
Taiwan	1 (0.67)
USA	2 (1.35)
Vietnam	3 (2.03)

*\*different countries in Europe*

**Supplementary Table 4.** Laboratory findings reported for adult and pregnant COVID-19 patients

Laboratory parameters	Normal range	Adult	Pregnant
<b>Hematology</b>			
Hemoglobin, g/L	120-160	137 [119-151]	
Leucocyte count, $\times 10^9/L$	3.5-10	4.8 [3.76-6.03]	7.63 [6.15-9.34]
Lymphocyte count, $\times 10^9/L$	0.9-3.3	1 [0.7-1.4]	0.87 [0.66-1.53]
Neutrophils count, $\times 10^9/L$	1.3-6.7	3.00 [2.20-4.41]	9.14 [9.14-9.14]
Platelet count, $\times 10^9/L$	150-450	177 [132-223]	
<b>Coagulation</b>			
Prothrombin time, s	11-13	12.2 [10.6-13.9]	
Activated partial thromboplastin time, s	25-34	31.40 [29.40-41.07]	
Fibrinogen, g/L	1.7-4	4.5 [3.66-5.1]	
<b>Electrolytes</b>			
Sodium, mmol/L	135-145	138.33 [136.18-140.50]	
Potassium, mmol/L	3.6-4.8	3.83 [3.50-4.23]	
<b>Clinical chemistry labs</b>			
Albumin, g/L	32-52	40.5 [35.0-43.0]	24.6
Alanine aminotransferase (ALT), U/L	8-41	23 [15-38]	16 [12-54]
Aspartate transaminase (AST), U/L	11-34	29.35 [19-43]	24 [22-71]
Total bilirubin, mmol/L	<15	9.6 [7-13.5]	
Lactate dehydrogenase, U/L	135-214	213 [173-268]	544
Creatine kinase, U/L	38-157	73 [48-111]	
D-dimer, $\mu g/mL$	<0.5	0.5 [0.3-1.08]	0.84
Creatinine, $\mu mol/L$	42-80	66 [56-78]	
Blood urea nitrogen, mmol/L	2.6-6.7	3.95 [2.9-5.69]	
<b>Inflammation</b>			
C-reactive protein, mg/L	<10	8.8 [2.6-22]	19.25 [12.35-25.7]
Interleukin 6, pg/mL	<3.1	22 [4.68-51.8]	
Procalcitonin, $\mu g/mL$	<0.1	0.07 [0.04-0.1]	0.187
Erythrocyte sedimentation rate, mm/h	Female: <20 Male: <10	32.5 [17.3-53.8]	

**Supplementary Table 5:** Results of meta-analysis of prevalence in pregnant COVID-19 patients

Variable	Numbers of studies	Patients	Total patients	Crude Prevalence [%]	Random effect model (REM) Prevalence*	REM (lower CI)	REM (upper CI)	Tau	I <sup>2</sup>	Q
<b>Comorbidities</b>										
Any comorbidity	2	2	10	20	20	5.04	54.07	0	0	3.73
Hypothyroidism	1	1	1	100						
Influenza	1	1	9	11.11						
Allergies	1	1	1	100						
<b>Signs and symptoms</b>										
Fever	10	25	35	71.43	96.35	17.57	99.97	7.628	81.5	19.95
Cough	6	12	29	41.38	41.38	25.21	59.64	0	0	5.8
Fatigue	3	5	18	27.78	58.94	1.45	99.29	2.893	64.1	5.83
Diarrhea	3	32	26	123.08	10.45	0.19	87.7	8.865	82.8	7.82
Sore throat	1	2	9	22.22						
Chest tightness	1	1	9	11.11						
Myalgia	1	3	9	33.33						
Dyspnea	2	3	19	15.79	15.79	5.18	39.15	0	0	0.7
Nasal congestion	1	1	1	100						
Dizziness or confusion	1	2	9	22.22						
<b>Imaging features</b>										
Pathologic findings	10	31	35	88.57	88.57	73.23	95.64	0	0	3.16
Pneumonia	10	31	35	88.57	88.57	73.23	95.64	0	0	3.16
Ground glass opacity (GGO)	9	29	34	85.29	85.29	69.18	93.74	0	0	4.12
Bilateral pneumonia	6	15	21	71.43	71.43	49.24	86.57	0	0	3.96
Unilateral pneumonia	2	4	17	23.53	23.53	9.12	48.55	0	0	3.11
Consolidation	1	8	16	50						
GGO with consolidation	3	9	18	50	50	28.42	71.58	0	0	3.02
Local patchy shadowing	1	1	1	100						
Pleural effusion	1	6	16	37.5						
Subpleural opacities	2	2	2	100						
Bronchiectasis	1	1	3	33.33						
<b>Treatment</b>										
Antiviral treatment	6	11	14	78.57	78.57	50.57	92.93	0	0	3.09
Antibiotics	6	14	14	100						
Corticosteroids	4	3	12	25	100	6.32	100	25.452.2	100	13.5
High flow nasal cannula	2	10	10	100						
Alpha-interferon aerosol inhalation	2	2	2	100						

	Lopinavir	2	2	2	100						
	Ritonavir	2	2	2	100						
	Oxygen therapy	4	3	12	25	100	6.32	100	25'452.2	100	13.5
	Moxifloxacin hydrochloride	2	2	2	100						
	Piperacillin tazobactam	1	1	1	100						
	Sulbactam	1	1	1	100						
<b>Outcome</b>											
	Death	7	0	9	0						
	Survived	7	9	9	100						
	Discharged	4	4	6	66.67	66.67	26.81	91.61	0	0	3.82
	Recovery	3	3	3	100						
	Remained hospitalized	3	1	3	33.33	33.33	1.98	92.51	0	0	3.82
<b>Complications</b>											
	Admission to intensive care unit	1	1	1	100						

\*Model does not convert if there is only one study or 100% of the patients have events.

**Supplementary Table 6:** Results of meta-analysis of prevalence in pediatric and neonatal COVID-19 patients

Variable	Numbers of studies	Patients	Total patients	Crude Prevalence [%]	Random effect model (REM) Prevalence*	REM (lower CI)	REM (upper CI)	Tau	I <sup>2</sup>	Q
<b>Comorbidities</b>										
Any comorbidity	1	2	25	8.00						
<b>Signs and symptoms</b>										
Any sign or symptom	27	903	1'055	85.59	84.95	72.02	92.52	1.642	81.0	62.59
Asymptomatic	26	149	1'054	14.14	16.17	8.51	28.59	1.389	79.1	60.84
Fever	22	170	320	53.13	67.51	51.63	80.17	0.935	66.5	67.56
Cough	14	149	311	47.91	51.29	35.07	67.25	0.833	73.9	37.03
Fatigue	6	20	235	8.51	8.51	5.56	12.82	0	0	6.64
Diarrhea	10	34	270	12.59	18.73	8.91	35.19	0.396	45.8	22.23
Sore throat	3	7	61	11.48	11.51	3.17	34.03	0.891	59.3	7.24
Sputum	6	14	51	27.45	36.45	11.22	72.24	0.714	44.5	12.11
Headache	2	4	39	10.26	10.26	3.9	24.33	0	0	0.05
Dyspnea	2	3	21	14.29	14.29	4.68	36.14	0	0	4.22
Nausea	5	10	55	18.18	29.10	6.56	70.6	1.781	67.6	14.12
Running nose (rhinorrhea)	10	26	259	10.04	10.67	3.36	29.12	0.034	5.2	17.14
Nasal congestion	4	13	198	6.57	6.57	3.85	10.98	0	0	5.29
Emesis or vomiting	2	15	177	8.47	23.67	2.69	77.7	2.566	85.8	13.46
<b>Imaging features</b>										
Pathologic findings	23	204	313	65.18	69.84	56.18	80.71	0.462	49.0	42.23
Pneumonia	22	194	298	65.10	67.76	56.08	77.58	0.233	31.7	36.24
Ground glass opacity (GGO)	13	108	278	38.85	48.36	34.88	62.08	0.329	50.0	27.87
Bilateral pneumonia	7	34	62	54.84	54.84	42.41	66.69	0	0	6.71
Unilateral pneumonia	5	15	55	27.27	27.27	17.16	40.43	0	0	6.00
Local patchy shadowing	8	52	223	23.32	34.11	16.01	58.44	0.535	57.2	21.97
Bilateral patchy shadowing	3	30	185	16.22	38.73	11.13	76.13	1.542	80.5	19.29
<b>Treatment</b>										
Antiviral treatment	8	40	83	48.19	98.41	17.38	99.99	19.812	94.0	71.5
Antibiotics	7	31	43	72.09	95.52	19.14	99.95	11.153	87.4	33.83
Corticosteroids	3	11	39	28.21	38.22	9.56	78.35	1.844	75.4	14.24
All mechanical ventilation	3	5	34	14.71	14.71	6.26	30.82	0	0	5.46
Invasive mechanical ventilation	2	4	33	12.12	12.12	4.62	28.18	0	0	1.44
High flow nasal cannula	2	5	9	55.56	55.56	25.13	82.32	0	0	1.27
Intravenous immunoglobulin	3	7	39	17.95	19.41	6	47.62	0.721	51.5	6.27
Alpha-interferon aerosol inhalation	6	31	52	59.62	99.34	4.17	100	13.796	90.8	31.17
<b>Outcome</b>										

Death	24	1	311	0.32	0.32	0.05	2.25	0	0	1.20
Survived	24	310	311	99.68	99.68	97.75	99.95	0	0	1.20
Discharged	16	216	290	74.48	76.94	48.64	92.16	2.765	83.5	100.48
Recovery	15	236	270	87.41	87.41	82.89	90.86	0	0	16.45
Remained hospitalized	12	90	293	30.72	72.69	29.11	94.52	6.921	93.2	155.50
<b>Complications</b>										
Admission to intensive care unit	5	6	41		14.91	6.08	32.18	0.157	13	4.21
All secondary infections	2	12	21	57.14	57.14	35.97	75.99	0	0	1.16
Respiratory failure	2	10	33	30.30	30.3	17.15	47.74	0	0	0.14
Pneumonia	2	16	26	61.54	61.54	42.07	77.9	0	0	1
Secondary infections (bacteria)	2	12	21	57.14	57.14	35.97	75.99	0	0	1.16


\*Model does not convert if there is only one study or 100% of the patients have events.

**Supplementary Table 7.** Laboratory findings reported for pediatric/neonatal COVID-19 patients

Laboratory parameters	Pediatric/neonatal
<b>Hematology</b>	
Hemoglobin, g/L	126 [118-135]
Leucocyte count, $\times 10^9/L$	6.8 [5.5-8.2]
Lymphocyte count, $\times 10^9/L$	2.9 [2.2-4.4]
Neutrophils count, $\times 10^9/L$	2.5 [1.8-3.7]
Platelet count, $\times 10^9/L$	200.5 [196-301]
<b>Coagulation</b>	
Prothrombin time, s	10.9 [10.6-11.3]
Activated partial thromboplastin time, s	42.1 [41.0-44.5]
Fibrinogen, g/L	2.1 [1.8-2.7]
<b>Electrolytes</b>	
Sodium, mmol/L	139.1 [138-140.5]
Potassium, mmol/L	4.8 [4.3-5.2]
<b>Clinical chemistry labs</b>	
Albumin, g/L	45.1 [44.5-47.9]
Alanine aminotransferase (ALT), U/L	15 [11-27]
Aspartate transaminase (AST), U/L	30 [24-42]
Total bilirubin, mmol/L	4 [3.6-4.2]
Lactate dehydrogenase, U/L	246 [207-305]
Creatine kinase, U/L	72 [29.9-74]
D-dimer, $\mu g/mL$	0.2 [0.2-0.4]
Creatinine, $\mu mol/L$	33.9 [26.1-42.7]
Blood urea nitrogen, mmol/L	4.1 [3.3-4.8]
<b>Inflammation</b>	
C-reactive protein, mg/L	4.0 [1.3-8.0]
Interleukin 6, pg/mL	11.59 [8.58-14.95]
Procalcitonin, ng/mL	0.05 [0.04-0.08]
Erythrocyte sedimentation rate, mm/h	6 [2-20]


*Normative values for children are highly age-dependent. Often a wide range of ages were reported as one value.*

## SUPPLEMENTARY FIGURES




**Supplementary Figure 1. All comorbidities in adult COVID-19 patients.** The key to the unique study identifier can be found in Table 1.







**Supplementary Figure 2. All imaging features in adult COVID-19 patients.** The key to the unique study identifier can be found in Table 1.




**Supplementary Figure 3. Symptoms and signs in pregnant COVID-19 patients.** The key to the unique study identifier can be found in Table 2.




**Supplementary Figure 4.** Imaging features in pregnant COVID-19 patients. The key to the unique study identifier can be found in Table 2.




**Supplementary Figure 5. Treatments administered to pregnant COVID-19 patients.** The key to the unique study identifier can be found in Table 2.




**Supplementary Figure 6. Symptoms and signs in pediatric and neonatal COVID-19 patients.** The key to the unique study identifier can be found in Table 3.



**Supplementary Figure 7. Imaging features in pediatric and neonatal COVID-19 patients.** The key to the unique study identifier can be found in Table 3.




**Supplementary Figure 8. Treatments administered to pediatric and neonatal COVID-19 patients.**  
The key to the unique study identifier can be found in Table 3.



**Supplementary Figure 9. Outcomes of pediatric and neonatal COVID-19 patients.** The key to the unique study identifier can be found in Table 3.





**Supplementary Figure 10.** Proportion of male and female patients in the non-severe, severe, survivors, and non-survivor patient groups. With the exception of the non-survivor with a larger proportion of male patients, the ration male:female patients was 1:1 in all the groups. The key to the study identifier can be found in Table 1 (adults).

Authors	Title	PMID	Unique study ID	Country	Language	Study type	Study population	Sample size	Age *	Male (%)	Female (%)
<b>Ai et al, 2020 [32]</b>	Correlation of Chest CT and RT-PCR Testing in Coronavirus Disease 2019 (COVID-19) in China: A Report of 1014 Cases	32101510	S1	China	English	Cohort Study	Adult	1014	51 (15)	467 (46)	547 (54)
<b>Albarell o et al, 2020 [159]</b>	2019-novel Coronavirus severe adult respiratory distress syndrome in two cases in Italy: An uncommon radiological presentation	32112966	S2	Italy	English	Case series	Adult	2	66.5 [66.25-66.75]	1 (50)	1 (50)
<b>An et al, 2020 [94]</b>	CT Manifestations of Novel Coronavirus Pneumonia: A Case Report	32157862	S3	China	English	Case Study	Adult	1	50	0 (0)	1 (100)
<b>Arentz et al, 2020 [59]</b>	Characteristics and Outcomes of 21 Critically Ill Patients With COVID-19 in Washington State	32191259	S4	USA	English	Cohort Study	Adult	21	70 [43-92]	11 (52)	10 (48)
<b>Bai et al, 2020 [128]</b>	Analysis of the first cluster of cases in a family of novel coronavirus pneumonia in Gansu Province	32064855	S5	China	Chinese	Case series	Adult	7	53.4	3 (43)	4 (57)
<b>Chan et al, 2020 [4]</b>	A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster	31986261	S6_adult	China	English	Case series	Adult	5	50 [36.25-64.50]	2 (40)	3 (60)
<b>Chang et al, 2020 [131]</b>	Epidemiologic and Clinical Characteristics of Novel Coronavirus Infections Involving 13 Patients Outside Wuhan, China	32031568	S7	China	English	Cohort Study	Adult	13	34 [34-48]	10 (77)	3 (33)
<b>Chen et al, 2020 [95]</b>	Analysis of clinical features of 29 patients with 2019 novel coronavirus pneumonia	32164089	S8	China	Chinese	Cohort Study	Adult	29	56 [range 26-79]	21 (72)	8 (28)
<b>Chen et al, 2020 [67]</b>	Analysis of myocardial injury in patients with COVID-19 and association between concomitant cardiovascular diseases and severity of COVID-19	32141280	S9_severe	China	Chinese	Cohort Study	Adult (severe)	24	68.5 (13.6)	18 (75)	6 (25)
<b>Chen et al, 2020 [67]</b>	Analysis of myocardial injury in patients with COVID-19 and association between concomitant cardiovascular diseases and severity of COVID-19	32141280	S9_non severe	China	Chinese	Cohort Study	Adult (non-severe)	126	57.1 (15.6)	66 (52)	60 (48)
<b>Chen et al, 2020 [39]</b>	Clinical progression of patients with COVID-19 in Shanghai, China	32171869	S10	China	English	Cohort Study	Adult	249	51 [36-64]	126 (51)	123 (49)
<b>Chen et al, 2020</b>	Epidemiological and clinical characteristics of 99 cases of 2019	32007143	S11	China	English	Cohort Study	Adult	99	55.5 (13.1)	67 (68)	32 (32)

[140]	novel coronavirus pneumonia in Wuhan, China: a descriptive study										
Chen et al, 2020 [161]	Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study	32217556	S12	China	English	Cohort Study	Adult	274	62 [44-70]	171 (62)	103 (38)
Cheng et al, 2020 [149]	Epidemiological characteristics of novel coronavirus pneumonia in Henan	32118390	S13	China	Chinese	Cohort Study	Adult	1079	46 [IQR: 24]	573 (53)	506 (47)
Cheng et al, 2020 [62]	Clinical Features and Chest CT Manifestations of Coronavirus Disease 2019 (COVID-19) in a Single-Center Study in Shanghai, China	32174128	S14	China	English	Cohort Study	Adult	11	50.36 (15.5)	8 (73)	3 (27)
Cheng et al, 2020 [157]	First case of Coronavirus Disease 2019 (COVID-19) pneumonia in Taiwan	32113824	S15	Taiwan	English	Case Study	Adult	1	55	0 (0)	1 (100)
** [69]	Early Epidemiological and Clinical Characteristics of 28 Cases of Coronavirus Disease in South Korea	32149037	S16	Korea	English	Cohort Study	Adult	28	42.6 [range 20-73]	15 (54)	13 (46)
Dai et al, 2020 [105]	CT Imaging and Differential Diagnosis of COVID-19	32129670	S17	China	English	Case Series	Adult	4	50 [47.75-55.125]	4 (100)	0 (0)
Deng et al, 2020 [68]	Clinical characteristics of fatal and recovered cases of coronavirus disease 2019 (COVID-19) in Wuhan, China: a retrospective study	32209890	S18_death	China	English	Cohort Study	Adult and pediatric	109	69[62-74]	73 (67)	36 (33)
Deng et al, 2020 [68]	Clinical characteristics of fatal and recovered cases of coronavirus disease 2019 (COVID-19) in Wuhan, China: a retrospective study	32209890	S18_survival	China	English	Cohort Study	Adult and pediatric	116	40[33-57]	51 (44)	65 (56)
Ding et al, 2020 [72]	The clinical characteristics of pneumonia patients coinfectd with 2019 novel coronavirus and influenza virus in Wuhan, China	32196707	S19	China	English	Case Series	Adult	5	49 [47-50]	2 (40)	3 (60)
Ding et al, 2020 [28]	A cured patient with 2019-nCoV pneumonia	32205073	S20	China	English	Case Study	Adult	1	57	0 (0)	1 (100)
Dong et al, 2020 [106]	Epidemiological characteristics of confirmed COVID-19 cases in Tianjin	32164400	S21	China	English	Cohort Study	Adult	135	48.62 (16.83)	72 (53)	63 (47)
Duan and Qin 2020 [144]	Pre- and Posttreatment Chest CT Findings - 2019 Novel Coronavirus (2019-nCoV) Pneumonia	32049602	S22	China	English	Case Study	Adult	1	46	0 (0)	1 (100)

<b>Fan et al, 2020 [49]</b>	Perinatal Transmission of COVID-19 Associated SARS-CoV-2: Should We Worry?	32182347	S23	China	English	Case Series	Adult	2	31.5 [30.25-32.75]	0 (0)	2 (100)
<b>Fang et al, 2020 [93]</b>	Changes of CT findings in a 2019 novel coronavirus (2019-nCoV) pneumonia patient	32073631	S24	China	English	Case Study	Adult	1	47	1 (100)	0 (0)
<b>Fang et al, 2020 [88]</b>	Comparisons of nucleic acid conversion time of SARS-CoV-2 of different samples in ICU and non-ICU patients	32209381	S25	China	English	Cohort Study	Adult	32	41	16 (50)	16 (50)
<b>Fang et al, 2020 [115]</b>	CT Manifestations of Two Cases of 2019 Novel Coronavirus (2019-nCoV) Pneumonia	32031481	S26	China	English	Case Series	Adult	2	38.5 [35.25-41.75]	1 (50)	1 (50)
<b>Gautret et al, 2020 [127]</b>	Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial	32205204	S27	France	English	Case Series	Adult	36	47 [24.5-61.5]	15 (42)	21 (58)
<b>Gross et al, 2020 [112]</b>	CT appearance of severe, laboratory-proven coronavirus disease 2019 (COVID-19) in a Caucasian patient in Berlin, Germany	32193883	S28	Germany	English	Case Study	Adult	1	61	1 (100)	0 (0)
<b>Guan et al, 2020 [25]</b>	Epidemiological investigation of a family clustering of COVID-19	32149484	S29	China	Chinese	Case Series	Adult	7	53.43	3 (43)	4 (57)
<b>Guan et al, 2020 [138]</b>	Clinical Characteristics of Coronavirus Disease 2019 in China	32109013	S30	China	English	Cohort Study	Adult	1099	47 [35-58]	639 (58)	460 (42)
<b>Guan et al, 2020 [142]</b>	CT Findings of Coronavirus Disease (COVID-19) Severe Pneumonia	32208010	S31	China	English	Case Study	Adult	1	59	0 (0)	1 (100)
<b>Guan et al, 2020 [120]</b>	Imaging Features of Coronavirusdisease 2019 (COVID-19): Evaluationon Thin-Section CT	32204990	S32	China	English	Cohort Study	Adult	53	42 [range 1-86]	25 (47)	28 (53)
<b>Han et al, 2020 [137]</b>	Early Clinical and CT Manifestations of Coronavirus Disease 2019 (COVID-19) Pneumonia	32181672	S33	China	English	Cohort Study	Adult	108	45	38 (35)	70 (65)
<b>Han et al, 2020 [104]</b>	The course of clinical diagnosis and treatment of a case infected with coronavirus disease 2019	32073161	S34	China	English	Case Study	Adult	1	47	1 (100)	0 (0)
<b>Hao, 2020 [30]</b>	Clinical features of atypical 2019 novel coronavirus pneumonia with an initially negative RT-PCR assay	32092387	S35	China	English	Case study	Adult	1	58	1 (100)	0 (0)
<b>He et al, 2020 [52]</b>	Impact of complicated myocardial injury on the clinical outcome of severe or critically ill COVID-19 patients	32171190	S36	China	Chinese	Cohort Study	Adult	54	68 [59.8-74.3]	34 (63)	20 (37)
<b>Hill et</b>	The index case of SARS-CoV-2 in	32205138	S37	Scotland	English	Case	Adult	1	51	1	0 (0)

<b>al, 2020 [118]</b>	Scotland: a case report					Study				(100)	
<b>Holshue et al, 2020 [113]</b>	First Case of 2019 Novel Coronavirus in the United States	32004227	S38	USA	English	Case Study	Adult	1	35	1 (100)	0 (0)
<b>Hosoda et al, 2020 [57]</b>	SARS-CoV-2 enterocolitis with persisting to excrete the virus for about two weeks after recovering from diarrhea: A case report	32188528	S39	Japan	English	Case Study	Adult	1	81	0 (0)	1 (100)
<b>Hu et al, 2020 [119]</b>	Clinical characteristics of 24 asymptomatic infections with COVID19 screened among close contacts in Nanjing, China	32146694	S40	China	English	Cohort Study	Adult	24	32.5 [19.0-57.0]	8 (33)	16 (64)
<b>Hu et al, 2020 [102]</b>	CT imaging of two cases of one family cluster 2019 novel coronavirus (2019-nCoV) pneumonia: inconsistency between clinical symptoms amelioration and imaging sign progression	32190575	S41	China	English	Case Series	Adult	2	42.5 [40.25-44.75]	1 (50)	1 (50)
<b>Huang et al, 2020 [76]</b>	Clinical characteristics of laboratory confirmed positive cases of SARS-CoV2 infection in Wuhan, China: A retrospective single center analysis	32114074	S42	China	English	Cohort Study	Adult	34	56.24 (17.14)	14 (41)	20 (59)
<b>Huang et al, 2020 [66]</b>	Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China	31986264	S43	China	English	Cohort Study	Adult	41	49 [41-58]	30 (73)	11 (27)
<b>Huang et al, 2020 [109]</b>	Use of Chest CT in Combination with Negative RT-PCR Assay for the 2019 Novel Coronavirus but High Clinical Suspicion	32049600	S44	China	English	Case Study	Adult	1	36	1 (100)	0 (0)
<b>Jin et al. 2020 [63]</b>	Epidemiological, clinical and virological characteristics of 74 cases of coronavirus-infected disease 2019 (COVID-19) with gastrointestinal symptoms	32213556	S45_wit hGI	China	English	Cohort Study	Adult - with GI Symptoms	74	46.14 (14.19)	37 (50)	37 (50)
<b>Jin et al. 2020 [63]</b>	Epidemiological, clinical and virological characteristics of 74 cases of coronavirus-infected disease 2019 (COVID-19) with gastrointestinal symptoms	32213556	S45_no GI	China	English	Cohort Study	Adult - No GI Symptoms	577	45.09 (14.45)	294 (51)	283 (49)
<b>Lee et al, 2020 [54]</b>	A case of COVID-19 and pneumonia returning from Macau in Taiwan: Clinical course and anti-SARS-CoV-2 IgG dynamic	32198005	S46	Vietnam	English	Case study	Adult	1	46	0 (0)	1 (100)

<b>Leung et al, 2020 [47]</b>	Clinical features of deaths in the novel coronavirus epidemic in China	32175637	S47	China	English	Cohort Study	Adult	46	70.6 (12.63)	31 (67)	15 (33)
<b>Li et al, 2020 [114]</b>	CT image visual quantitative evaluation and clinical classification of coronavirus disease (COVID-19)	32215691	S48	China	English	Cohort Study	Adult	78	44.6 (17.9)	38 (49)	40 (51)
<b>Li et al, 2020 [99]</b>	Characteristics of peripheral blood leukocyte differential counts in patients with COVID-19	32114745	S49	China	Chinese	Cohort Study	Adult	10	46.5 [36.5-64.3]	5 (50)	5 (50)
<b>Li et al, 2020 [96]</b>	Comparison of epidemic characteristics between SARS in 2003 and COVID-19 in 2020 in Guangzhou	32159317	S50	China	Chinese	Cohort Study	Adult	346	48 [range 3 months-90 yo]	167 (48)	179 (52)
<b>Li et al, 2020 [111]</b>	Comparison of the clinical characteristics between RNA positive and negative patients clinically diagnosed with 2019 novel coronavirus pneumonia	32087623	S51	China	Chinese	Cohort Study	Adult	31	54	15 (48)	16 (52)
<b>Lian et al, 2020 [77]</b>	Analysis of Epidemiological and Clinical features in older patients with Corona Virus Disease 2019 (COVID-19) out of Wuhan	32211844	S52_young	China	English	Cohort Study	Adult (young and middle-aged < 60 years)	652	41.15 (1.38)	349 (54)	303 (46)
<b>Lian et al, 2020 [77]</b>	Analysis of Epidemiological and Clinical features in older patients with Corona Virus Disease 2019 (COVID-19) out of Wuhan	32211844	S52_old	China	English	Cohort Study	Adult (elderly >= 60 years)	136	68.28 (7.31)	58 (43)	78 (57)
<b>Lin et al, 2020 [40]</b>	Novel coronavirus pneumonia outbreak in 2019: Computed tomographic findings in two cases	32056397	S53	China	English	Case Series	Adult	2	37 [36-38]	2 (100)	0 (0)
<b>Liu et al, 2020 [147]</b>	Clinical feature of COVID-19 in elderly patients: a comparison with young and middle-aged patients	32171866	S54_old	China	English	Cohort Study	Adult (elderly >= 60 years)	18	68.00 [65.25-69.75]	12 (67)	6 (33)
<b>Liu et al, 2020 [147]</b>	Clinical feature of COVID-19 in elderly patients: a comparison with young and middle-aged patients	32171866	S54_young	China	English	Cohort Study	Adult (young and middle-aged < 60 years)	38	47 [35.75-51.25]	19 (50)	19 (50)
<b>Liu et al, 2020 [153]</b>	Gross examination of report of a COVID-19 death autopsy	32198987	S55	China	Chinese	Case Study	Adult	1	85	1 (100)	0 (0)
<b>Liu et al, 2020 [90]</b>	Clinical characteristics of 30 medical workers infected with new coronavirus pneumonia	32062957	S56	China	Chinese	Cohort Study	Adult	30	35 [21-59]	10 (33)	20 (67)
<b>Liu et</b>	Analysis of factors associated with	32118640	S57	China	English	Cohort	Adult	78	38 [33-57]	39	39 (50)



<b>al, 2020 [81]</b>	disease outcomes in hospitalized patients with 2019 novel coronavirus disease					Study				(50)		
<b>Liu et al, 2020 [74]</b>	Clinical and biochemical indexes from 2019-nCoV infected patients linked to viral loads and lung injury	32048163	S58_adult	China	English	Case Series	Adult	12	63 [53.5-65]	8 (67)	4 (33)	
<b>Liu et al, 2020 [133]</b>	Clinical characteristics of novel coronavirus cases in tertiary hospitals in Hubei Province.	32044814	S59	China	English	Cohort Study	Adult	137	57 [range 20-83]	61 (45)	76 (55)	
<b>Liu et al, 2020 [146]</b>	Clinical and CT imaging features of the COVID-19 pneumonia: Focus on pregnant women and children	32171865	S60_adult	China	English	Cohort Study	Adult	14		5 (36)	9 (64)	
<b>Mo et al, 2020 [98]</b>	Clinical characteristics of refractory COVID-19 pneumonia in Wuhan, China	32173725	S61	China	English	Cohort Study	Adult	155	54 [42-66]	86 (55)	69 (45)	
<b>Pan et al, 2020 [132]</b>	Initial CT findings and temporal changes in patients with the novel coronavirus pneumonia (2019-nCoV): a study of 63 patients in Wuhan, China	32055945	S62	China	English	Cohort Study	Adult	63	44.9 (15.2)	33 (52)	30 (48)	
<b>Peng et al, 2020 [145]</b>	Clinical characteristics and outcomes of 112 cardiovascular disease patients infected by 2019-nCoV	32120458	S63_severe	China	Chinese	Cohort Study	Adult (severe)	16	57.5 [54-63]	9 (56)	7 (44)	
<b>Peng et al, 2020 [36]</b>	Clinical characteristics and outcomes of 112 cardiovascular disease patients infected by 2019-nCoV	32120458	S63_nonsevere	China	Chinese	Cohort Study	Adult (non-severe)	96	62 [55-67.5]	44 (46)	52 (54)	
<b>Qian et al, 2020 [80]</b>	A COVID-19 Transmission within a family cluster by presymptomatic infectors in China	32201889	S64_adult	China	English	Case series	Adult	7	57.5 [44.5-59]	3 (43)	4 (57)	
<b>Qian et al, 2020 [156]</b>	Epidemiologic and Clinical Characteristics of 91 Hospitalized Patients with COVID-19 in Zhejiang, China: A retrospective, multi-centre case series	32181807	S65	China	English	Cohort Study	Adult	91	50 [36.5-57]	37 (41)	54 (59)	
<b>Qu et al, 2020 [61]</b>	Platelet-to-lymphocyte ratio is associated with prognosis in patients with coronavirus disease-19	32181903	S66	China	English	Cohort Study	Adult	30	50.5 [36-65]	16 (53)	14 (47)	
<b>Ren et al, 2020 [165]</b>	Identification of a novel coronavirus causing severe pneumonia in human - a descriptive study	32004165	S67	China	English	Case Series	Adult	5	52 [49-61]	3 (60)	2 (40)	
<b>Ruan et al, 2020 [130]</b>	A case of 2019 novel coronavirus infected pneumonia with twice negative 2019-nCoV nucleic acid testing within 8 days	32149771	S68	China	English	Case study	Adult	1	47	0 (0)	1 (100)	
<b>Shi et al, 2020</b>	Association of Cardiac Injury With Mortality in Hospitalized Patients With	32211816	S69	China	English	Cohort Study	Adult	416	64 [range 21-90]	205 (49)	211 (51)	

<b>[44]</b>	COVID-19 in Wuhan, China										
<b>Shi et al, 2020 [51]</b>	Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study	32105637	S70	China	English	Cohort Study	Adult	81	49.5 (11)	42 (52)	39 (48)
<b>Shi et al. 2020 [103]</b>	Evolution of CT Manifestations in a Patient Recovered from 2019 Novel Coronavirus (2019-nCoV) Pneumonia in Wuhan, China	32032497	S71	China	English	Case Study	Adult	1	42	1 (100)	0 (0)
<b>Silverstein et al, 2020 [91]</b>	First imported case of 2019 novel coronavirus in Canada, presenting as mild pneumonia	32061312	S72	Canada	English	Case Study	Adult	1	56	1 (100)	0 (0)
<b>Song et al, 2020 [60]</b>	SARS-CoV-2 induced diarrhoea as onset symptom in patient with COVID-19	32139552	S73	China	English	Case Study	Adult	1	22	1 (100)	0 (0)
<b>Song et al, 2020 [164]</b>	Emerging 2019 Novel Coronavirus (2019-nCoV) Pneumonia	32027573	S74	China	English	Cohort Study	Adult	51	49 (16)	25 (49)	26 (51)
<b>Spiteri et al, 2020 [31]</b>	First cases of coronavirus disease 2019 (COVID-19) in the WHO European Region, 24 January to 21 February 2020	32156327	S75	Europe	English	Cohort Study	Adult	38	42 [range 2-81]	25 (66)	13 (34)
<b>Stoecklin et al, 2020 [160]</b>	First cases of coronavirus disease 2019 (COVID-19) in France: surveillance, investigations and control measures, January 2020	32070465	S76	France	English	Case Series	Adult	3	31 [30.5-39.5]	2 (67)	1 (33)
<b>Sun et al, 2020 [134]</b>	Epidemiological and Clinical Predictors of COVID-19	32211755	S77	Singapore	English	Cohort Study	Adult	54	42 [34-54]	29 (54)	25 (46)
<b>Sun et al, 2020 [89]</b>	Evolution of Computed Tomography Manifestations in Five Patients Who Recovered from Coronavirus Disease 2019 (COVID-19) Pneumonia.	32174054	S78	China	English	Case Series	Adult	5	45 [range 20-55]	2 (40)	3 (60)
<b>Tang et al, 2020 [108]</b>	Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia	32073213	S79	China	English	Cohort Study	Adult	183	54.1 (16.2)	98 (54)	85 (46)
<b>Tian et al, 2020 [148]</b>	Characteristics of COVID-19 infection in Beijing	32112886	S80	China	English	Cohort Study	Adult	262	47.5 [range 1-94]	127 (48)	135 (52)
<b>Tian et al, 2020 [84]</b>	Pulmonary Pathology of Early-Phase 2019 Novel Coronavirus (COVID-19) Pneumonia in Two Patients With Lung Cancer	32114094	S81_case1	China	English	Case Study	Adult	1	73	1 (100)	0 (0)



<b>Tian et al, 2020 [84]</b>	Pulmonary Pathology of Early-Phase 2019 Novel Coronavirus (COVID-19) Pneumonia in Two Patients With Lung Cancer	32114094	S81_ca se2	China	English	Case Study	Adult	1	84	0 (0)	1 (100)
<b>Tong et al, 2020 [38]</b>	Potential Presymptomatic Transmission of SARS-CoV-2, Zhejiang Province, China, 2020	32091386	S82	China	English	Case Series	Adult	6	23.00 [15.00-41.75]	3 (50)	3 (50)
<b>Van Cuong et al, 2020 [154]</b>	The first Vietnamese case of COVID-19 acquired from China	32085849	S83	Vietnam	English	Case Study	Adult	1	25	0 (0)	1 (100)
<b>Wan et al, 2020 [70]</b>	Clinical Features and Treatment of COVID-19 Patients in Northeast Chongqing	32198776	S84	China	English	Cohort Study	Adult	135	47 [36-55]	72 (53)	63 (47)
<b>Wang et al, 2020 [56]</b>	Clinical characteristics and therapeutic procedure for four cases with 2019 novel coronavirus pneumonia receiving combined Chinese and Western medicine treatment	32037389	S85	China	English	Case series	Adult	4	47.5 [28.75-63]	3 (75)	1 (25)
<b>Wang et al, 2020 [139]</b>	Clinical Characteristics of 138 Hospitalized Patients with 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China	32031570	S86	China	English	Cohort Study	Adult	138	56 [42 - 68]	75 (54)	63 (46)
<b>Wang et al, 2020 [100]</b>	Clinical Features of 69 Cases with Coronavirus Disease 2019 in Wuhan, China	32176772	S87	China	English	Cohort Study	Adult	69	42 [35-62]	32 (46)	37 (54)
<b>Wang et al, 2020 [45]</b>	Clinical Outcomes in 55 Patients With Severe Acute Respiratory Syndrome Coronavirus 2 Who Were Asymptomatic at Hospital Admission in Shenzhen, China	32179910	S88	China	English	Cohort Study	Adult	55	49 [range 2-69]	22 (40)	33 (60)
<b>Wang et al, 2020 [26]</b>	The clinical dynamics of 18 cases of COVID-19 outside of Wuhan, China	32139464	S89	China	English	Cohort Study	Adult	18	39 [29-55]	10 (56)	8 (44)
<b>Wu et al, 2020 [125]</b>	Clinical Characteristics of Imported Cases of COVID-19 in Jiangsu Province: A Multicenter Descriptive Study	32109279	S90	China	English	Cohort Study	Adult	80	46.1(15.42)	39 (49)	41 (51)
<b>Wu et al, 2020 [85]</b>	Biological characters analysis of COVID-19 patient accompanied with aplastic anemia	32145715	S91	China	Chinese	Case Study	Adult	1	48	1 (100)	0 (0)
<b>Xie et al, 2020 [82]</b>	Comparison of different samples for 2019 novel coronavirus detection by nucleic acid amplification tests	32114193	S92	China	English	Case Series	Adult	9	34 [26-45]	4 (44)	5 (56)

<b>Xiong et al, 2020 [50]</b>	Clinical and High-Resolution CT Features of the COVID-19 Infection: Comparison of the Initial and Follow-up Changes	32134800	S93	China	English	Cohort Study	Adult	42	49.5 (14.1)	25 (60)	17 (40)
<b>Xu et al, 2020 [73]</b>	Clinical and computed tomographic imaging features of novel coronavirus pneumonia caused by SARS-CoV-2	32109443	S94	China	English	Cohort Study	Adult	50	43.9 (16.8)	29 (58)	21 (42)
<b>Xu et al, 2020 [22]</b>	Imaging and clinical features of patients with 2019 novel coronavirus SARS-CoV-2	32107577	S95	China	English	Cohort Study	Adult	90	50 [range 18-86]	39 (43)	51 (57)
<b>Xu et al, 2020 [64]</b>	Pathological findings of COVID-19 associated with acute respiratory distress syndrome	32085846	S96	China	English	Case Study	Adult	1	50	1 (100)	0 (0)
<b>Xu et al, 2020 [8]</b>	Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARS-Cov-2) outside of Wuhan, China: retrospective case series	32075786	S97	China	English	Cohort Study	Adult	62	41 [32-52]	35 (56)	27 (44)
<b>Xu et al, 2020 [41]</b>	Clinical features and dynamics of viral load in imported and non-imported patients with COVID-19	32179140	S98_imported	China	English	Cohort Study	Adult	15	35	10 (67)	5 (33)
<b>Xu et al, 2020 [41]</b>	Clinical features and dynamics of viral load in imported and non-imported patients with COVID-19	32179140	S98_secondary	China	English	Cohort Study	Adult	17	37	7 (41)	10 (59)
<b>Xu et al, 2020 [41]</b>	Clinical features and dynamics of viral load in imported and non-imported patients with COVID-19	32179140	S98_tertiary	China	English	Cohort Study	Adult	19	53	8 (42)	11 (58)
<b>Yang et al, 2020 [126]</b>	Clinical characteristics and imaging manifestations of the 2019 novel coronavirus disease (COVID-19):A multi-center study in Wenzhou city, Zhejiang, China	32112884	S99	China	English	Cohort Study	Adult	149	45.11 (13.35)	81 (54)	68 (46)
<b>Yang et al, 2020 [9]</b>	Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study	32105632	S100	China	English	Cohort Study	Adult	52	59.7 (13.3)	35 (67)	17 (33)
<b>Yao et al, 2020 [42]</b>	Clinical characteristics and influencing factors of patients with novel coronavirus pneumonia combined with liver injury in Shaanxi region	32153170	S101	China	Chinese	Cohort Study	Adult	40	53.87 (15.84)	25 (63)	15 (37)
<b>Yao et al, 2020</b>	Epidemiological characteristics of 2019-nCoV infections in Shaanxi, China by	32139462	S102	China	English	Cohort Study	Adult	195	44.13 (15.8)	129 (66)	66 (34)

[143]	February 8, 2020										
Ye et al, 2020 [23]	Clinical characteristics of severe acute respiratory syndrome coronavirus 2 reactivation	32171867	S103	China	English	Case series	Adult	5	31 [30-32]	2 (40)	3 (60)
Yoon et al, 2020 [141]	Chest Radiographic and CT Findings of the 2019 Novel Coronavirus Disease (COVID-19): Analysis of Nine Patients Treated in Korea	32100485	S104	South Korea	English	Cohort Study	Adult	9	54	4 (44)	5 (56)
Young et al, 2020 [79]	Epidemiologic Features and Clinical Course of Patients Infected With SARS-CoV-2 in Singapore	32125362	S105	Singapore	English	Cohort Study	Adult	18	47 [31-71]	9 (50)	9 (50)
Yu et al, 2020 [150]	A Familial Cluster of Infection Associated With the 2019 Novel Coronavirus Indicating Possible Person-to-Person Transmission During the Incubation Period	32067043	S106	China	English	Case series	Adult	4	72 [68-78.25]	2 (50)	2 (50)
Yuan et al, 2020 [55]	Association of radiologic findings with mortality of patients infected with 2019 novel coronavirus in Wuhan, China	32191754	S107	China	English	Cohort Study	Adult	27	60 [47-69]	12 (44)	15 (56)
Zhang et al, 2020 [71]	CT image of novel coronavirus pneumonia: a case report	32189175	S108	China	English	Case Study	Adult	1	64	1 (100)	0 (0)
Zhang et al, 2020 [27]	Clinical features of 2019 novel coronavirus pneumonia in the early stage from a fever clinic in Beijing	32164091	S109	China	Chinese	Cohort Study	Adult	9	36 [15-49]	5 (56)	4 (44)
Zhang et al, 2020 [34]	Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China	32077115	S110	China	English	Cohort Study	Adult	140	57 [range 25-87]	71 (51)	69 (49)
Zhang et al, 2020 [155]	Epidemiological, clinical characteristics of cases of SARS-CoV-2 infection with abnormal imaging findings.	32205284	S111	China	English	Cohort Study	Adult	573	46.65 (13.83)	295 (51)	278 (49)
Zhang et al, 2020 [117]	High-resolution CT features of 17 cases of Corona Virus Disease 2019 in Sichuan province, China	32139463	S112	China	English	Cohort Study	Adult	17	48.6 [range 23-74]	8 (47)	9 (53)
Zhao et al, 2020 [162]	The characteristics and clinical value of chest CT images of novel coronavirus pneumonia	32199619	S113	China	English	Cohort Study	Adult	80	44 (1.77)	43 (54)	37 (46)
Zhao et al, 2020	A comparative study on the clinical features of COVID-19 pneumonia to	32161968	S114	China	English	Cohort Study	Adult	19	48 [27-56]	11 (58)	8 (42)

<b>[78]</b>	other pneumonias										
<b>Zhou et al, 2020 [48]</b>	Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study	32171076	S115	China	English	Cohort Study	Adult	191	56 [46.0-67.0]	119 (62)	72 (38)
<b>Zhu et al, 2020 [110]</b>	Comparison of heart failure and 2019 novel coronavirus pneumonia in chest CT features and clinical characteristics	32129583	S116	China	Chinese	Cohort Study	Adult	12	52 [32-73]	8 (67)	4 (33)
<b>Zhu et al. 2020 [58]</b>	Clinical and CT imaging features of 2019 novel coronavirus disease (COVID-19)	32142928	S117	China	English	Case Series	Adult	6	43 [32-56]	0 (0)	6 (100)

\* mean(sd) or median[Q1-Q3]

\*\* COVID-19 National Emergency Response Center, Epidemiology and Case Management Team, Korea Centers for Disease Control and Prevention, Cheongju, Korea et al. 2020

Authors	Title	PMID	Unique study ID	Country	Language	Study type	Study population	Sample size	Age *	Male (%)	Female (%)
<b>Chen et al, 2020 [29]</b>	Pregnant women with new coronavirus infection: a clinical characteristics and placental pathological analysis of three cases	32114744	S118	China	Chinese	Case Series	Pregnant	3	29.6	0 (0)	3 (100)
<b>Chen et al, 2020 [75]</b>	Chest computed tomography images of early coronavirus disease (COVID-19)	32162211	S119	China	English	Case Study	Pregnant	1	27	0 (0)	1 (100)
<b>Chen et al, 2020 [11]</b>	Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records	32151335	S120	China	English	Case series	Pregnant	9	28 [26-33]	0 (0)	9 (100)
<b>Dong et al. 2020 [46]</b>	Possible Vertical Transmission of SARS-CoV-2 From an Infected Mother to Her Newborn	32215581	S121_pregnant	China	English	Case Study	Pregnant	1	29	0 (0)	1 (100)
<b>Liao et al, 2020 [122]</b>	Chest CT Findings in a Pregnant Patient with 2019 Novel Coronavirus Disease	32212578	S122	China	English	Case Study	Pregnant	1	25	0 (0)	1 (100)
<b>Liu et al, 2020 [146]</b>	Clinical and CT imaging features of the COVID-19 pneumonia: Focus on pregnant women and children	32171865	S60_pregnant	China	English	Cohort Study	Pregnant	16	30 [26-35]	0 (0)	16 (100)
<b>Wang et al, 2020 [123]</b>	A case of 2019 Novel Coronavirus in a pregnant woman with preterm delivery	32119083	S123	China	English	Case study	Pregnant	1	28	0 (0)	1 (100)
<b>Wang et al, 2020 [152]</b>	A case report of neonatal COVID-19 infection in China	32161941	S124_pregnant	China	English	Case study	Pregnant	1	34	0 (0)	1 (100)
<b>Wen et al, 2020 [124]</b>	A patient with SARS-CoV-2 infection during pregnancy in Qingdao, China	32198004	S125	China	English	Case study	Pregnant	1	31	0 (0)	1 (100)
<b>Xia et al, 2020 [136]</b>	Emergency Caesarean delivery in a patient with confirmed coronavirus disease 2019 under spinal anaesthesia	32192711	S126	China	English	Case Study	Adult	1	27	0 (0)	1 (100)

\* mean(sd) or median[Q1-Q3]

Authors	Title	PMID	Unique study ID	Country	Language	Study type	Study population	Sample size	Age *	Male (%)	Female (%)
<b>Cai et al, 2020 [65]</b>	First case of 2019 novel coronavirus infection in children in Shanghai	32102141	S127	China	Chinese	Case Study	Pediatric	1	7	1 (100)	0 (0)
<b>Chan et al, 2020 [4]</b>	A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster	31986261	S6_pediatric	China	English	Case series	Pediatric	1	10	1 (100)	0 (0)
<b>Chen et al, 2020 [101]</b>	First case of severe childhood novel coronavirus pneumonia in China	32135586	S128	China	Chinese	Case Study	Pediatric	1	1.1	1 (100)	0 (0)
<b>Cui et al, 2020 [33]</b>	A 55-Day-Old Female Infant Infected With 2019 Novel Coronavirus Disease: Presenting With Pneumonia, Liver Injury, and Heart Damage	32179908	S129	China	English	Case study	Neonatal	1	55 (days)	0 (0)	1 (100)
<b>Dong et al. 2020 [46]</b>	Possible Vertical Transmission of SARS-CoV-2 From an Infected Mother to Her Newborn	32215581	S121_pediatric	China	English	Case Study	Neonatal	1	0	0 (0)	1 (100)
<b>Dong et al. 2020 [92]</b>	Epidemiological Characteristics of 2143 Pediatric Patients With 2019 Coronavirus Disease in China	DOI: 10.1542/peds.2020-0702	S130	China	English	Cohort Study	Pediatric	731	10 [2-13]	420 (57)	311 (43)
<b>Fan et al, 2020 [158]</b>	Anal swab findings in an infant with COVID-19	DOI : 10.1002/ped4.12186	S131	China	English	Case Study	Neonatal	1	0.25	0 (0)	1 (100)
<b>Feng et al, 2020 [87]</b>	Analysis of CT features of 15 children with 2019 novel coronavirus infection	32061200	S132	China	Chinese	Case series	Pediatric	15	7 [range 4-14]	5 (33)	10 (67)
<b>Ji et al, 2020 [135]</b>	Clinical features of pediatric patients with COVID-19: a report of two family cluster cases	32180140	S133	China	English	Case Series	Pediatric	2	12.0 [10.5-13.5]	2 (100)	0 (0)
<b>Le et al, 2020 [24]</b>	The first infant case of COVID-19 acquired from a secondary transmission in Vietnam	32213326	S134	Vietnam	English	Case Study	Neonatal	1	0.25	0 (0)	1 (100)
<b>Liu et al, 2020 [74]</b>	Clinical and biochemical indexes from 2019-nCoV infected patients linked to viral loads and lung injury	32048163	S58_pediatric	China	English	Case Study	Pediatric	1	10	1 (100)	0 (0)
<b>Liu et al, 2020 [86]</b>	Detection of Covid-19 in Children in Early January 2020 in Wuhan, China	32163697	S135	China	English	Case Series	Pediatric	6	3 [3-3.75]	2 (33)	4 (67)
<b>Liu et</b>	Clinical and CT imaging features of the	32171865	S60_pe	China	English	Cohort	Pediatric	4	3.0 [0.7- 6.0]	2 (50)	2 (50)

al, 2020 [146]	COVID-19 pneumonia: Focus on pregnant women and children		diatric			Study						
Lu et al, 2020 [37]	SARS-CoV-2 Infection in Children	32187458	S136	China	English	Cohort Study	Pediatric	171	6.7 [2-9.8]	104 (61)	67 (39)	
Park et al, 2020 [163]	First Pediatric Case of Coronavirus Disease 2019 in Korea	32193905	S137	South Korea	English	Case Study	Pediatric	1	10	0 (0)	1 (100)	
Qian et al, 2020 [80]	A COVID-19 Transmission within a family cluster by presymptomatic infectors in China	32201889	S64_pediatic	China	English	Case study	Pediatric (asymptomatic)	1	1.1	0 (0)	1 (100)	
Sun et al, 2020 [116]	Clinical features of severe pediatric patients with coronavirus disease 2019 in Wuhan: a single center's observational study	32193831	S138	China	English	Case series	Pediatric	8	10.2 [5.04-13.54]	6 (75)	2 (25)	
Tang et al, 2020 [43]	Detection of Novel Coronavirus by RT-PCR in Stool Specimen from Asymptomatic Child, China	32150527	S139	China	English	Case Study	Pediatric	1	10	1 (100)	0 (0)	
Wang et al, 2020 [83]	SARS-CoV-2 infection with gastrointestinal symptoms as the first manifestation in a neonate	32204755	S140	China	Chinese	Case Study	Neonatal	1	19 (days)	1 (100)	0 (0)	
Wang et al, 2020 [152]	A case report of neonatal COVID-19 infection in China	32161941	S124_pediatic	China	English	Case study	Neonatal	1	0	1 (100)	0 (0)	
Wang et al, 2020 [107]	Clinical analysis of 31 cases of 2019 novel coronavirus infection in children from six provinces (autonomous region) of northern China	32118389	S141	China	Chinese	Cohort Study	Pediatric	31	7.1 [0.6-17]	15 (48)	16 (52)	
Wei et al, 2020 [53]	Novel Coronavirus Infection in Hospitalized Infants Under 1 Year of Age in China	32058570	S142	China	English	Case Series	Neonatal	9	0.583 [0.33-0.75]	2 (22)	7 (78)	
Xia et al, 2020 [151]	Clinical and CT features in pediatric patients with COVID-19 infection: Different points from adults	32134205	S143	China	English	Cohort Study	Pediatric	20	2 [range 1 day-14 years 7 months]	13 (65)	7 (35)	
Xu et al, 2020 [35]	Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding	PMCID: PMC7095102	S144	China	English	Case Series	Pediatric	10	6.63 [2.17-13.4]	6 (60)	4 (40)	
Zeng et al, 2020 [97]	First case of neonate infected with novel coronavirus pneumonia in China	32065520	S145	China	Chinese	Case Study	Neonatal	1	17 (days)	1 (100)	0 (0)	
Zhang et al, 2020 [145]	2019-novel coronavirus infection in a three-month-old baby	32043842	S146	China	Chinese	Case study	Neonatal	1	0.25	0 (0)	1 (100)	

<b>Zheng et al, 2020 [129]</b>	Clinical Characteristics of Children with Coronavirus Disease 2019 in Hubei, China	32207032	S147	China	English	Cohort Study	Pediatric	25	3 [2-9]	14 (56)	11 (44)
<b>Zhou et al, 2020 [121]</b>	Clinical features and chest CT findings of coronavirus disease 2019 in infants and young children	32204756	S148	China	Chinese	Case Series	Pediatric	9	1 [range 7 months-3 years]	4 (44)	5 (56)

\* mean(sd) or median[Q1-Q3]



**Table 4:** Summary for random effects model for prevalence of comorbidities, clinical signs and symptoms, imaging features, treatments, outcome and complications in adult COVID-19 patients.

Variable	Numb of studies	Patients	Total patients	Crude prevalence [%]	Random effect model (REM) Prevalence	REM (lower CI)	REM (upper CI)	Tau $T^2$	$I^2$	Q
<b>Comorbidities</b>										
Any comorbidity	85	2'329	7'608	30.61	29.57	24.08	35.71	1.271	95.4	902.92
Hypertension	58	1'352	6'460	20.93	23.24	19.23	27.8	0.585	90.4	517.13
Diabetes mellitus	53	678	6'535	10.37	11.81	10.12	13.72	0.218	70.0	187.38
Carcinoma	36	111	6'033	1.84	2.15	1.56	2.95	0.447	56.7	95.51
Chronic obstructive pulmonary disease	29	86	5'232	1.64	1.70	0.92	3.1	1.976	84.5	147.80
Cardiovascular disease	28	180	3'747	4.80	6.09	4.04	9.10	1.014	85.0	208.10
Chronic kidney disease	20	56	3'521	1.59	1.85	0.93	3.63	1.536	79.4	84.55
Coronary heart disease	17	194	2'388	8.12	9.32	4.53	18.21	2.167	94.3	294.63
Any liver disease	15	51	580	8.79	3.85	1.44	9.89	2.215	83.5	95.23
Cerebrovascular disease	13	112	2'568	4.36	3.95	2.12	7.23	1.025	87.5	143.74
Current smoker	13	266	3'400	7.82	5.79	4.32	7.72	0.156	68.8	62.4
Hepatitis B	12	54	2'333		2.72	1.41	5.16	0.724	71.5	34.60
Chronic liver disease	11	95	2'576	3.69	3.69	3.03	4.49	0	0	15.53
Any respiratory system disease	10	49	1'020	4.80	2.95	1.28	6.67	1.045	78.8	45.62
Heart failure	5	37	354	10.45	20.12	2.25	73.36	5.885	95.6	43.48
Immunodeficiency	5	6	418	1.44	1.62	0.18	12.8	3.889	81.3	19.40
<b>Clinical signs and symptoms</b>										
Asymptomatic	69	148	2'749	5.38	0.4	0.07	2.21	11.535	93.6	664.40
Patients reported with any sign or symptom	65	1'936	2'597	74.55	98.03	92.48	99.51	9.123	96.2	864.92
Fever	110	6'955	8'859	78.51	82.96	79.13	86.21	0.968	91.6	1'096.03
Cough	102	4'778	8'885	53.78	58.38	53.92	62.70	0.528	90.1	1'671.74
Fatigue	69	1'996	7'980	25.01	29.25	24.03	35.07	0.918	94.2	1'140.98
Diarrhea	58	465	6'475	7.18	8.32	6.63	10.4	0.497	76.9	343.2
Sore throat	49	726	6'538	11.10	13.04	10.0	16.84	0.683	88.7	357.51
Sputum	48	1'437	6'118	23.49	25.06	19.68	31.35	0.850	94.3	904.12
Headache	48	710	7'564	9.39	10.4	8.29	12.97	0.511	86.0	326.48
Chest tightness	46	885	4'596	19.26	24.21	17.02	33.21	1.737	95.3	882.92
Myalgia	46	808	5'284	15.29	18.99	14.69	24.19	0.779	90.7	411.98

Dyspnea	39	705	5'730	12.30	15.20	10.54	21.43	1.446	94.8	881.01
Nausea	31	329	5'361	6.14	7.06	4.87	10.11	0.837	88.0	211.11
Running nose (rhinorrhea)	25	113	2'513	4.50	7.30	4.57	11.46	0.676	71.3	115.97
Nasal congestion	20	219	4'487	4.88	9.32	4.7	17.65	2.089	94.7	166.83
Dizziness or confusion	18	97	1'054	9.20	13.6	6.92	24.97	1.376	84.8	85.04
Hemoptysis	13	65	3'298	1.97	2.37	1.62	3.44	0.170	44.2	23.26
Anorexia	10	205	1'202	17.05	14.21	7.3	25.84	1.132	93.9	131.95
Emesis or vomiting	6	38	857	4.43	4.43	3.24	6.04	0	0	4.42
Chest pain	6	64	832	7.69	7.78	2.97	18.86	1.389	90.8	90.01
Abdominal pain	7	38	740	5.14	5.11	2.93	8.77	0.223	46.2	22.04
<b>Imaging features</b>										
Pathologic findings	93	6'969	7'780	89.58	97.83	95.38	99.00	5.934	97.4	952.20
Pneumonia	93	6'620	7'917	83.62	96.87	93.71	98.47	5.885	98.1	1'610.32
Ground glass opacity (GGO)	62	2'446	5'591	43.75	69.13	56.74	79.27	2.900	97.9	1'126.68
Bilateral pneumonia	48	2'745	4'247	64.63	77.29	70.08	83.17	1.173	94.6	931.56
Unilateral pneumonia	32	799	3'745	21.34	19.27	16.46	22.43	0.154	73.0	86.28
Consolidation	30	771	2'022	38.13	38.33	26.94	51.16	1.265	92.1	271.44
GGO with consolidation	15	153	323	47.37	49.53	40.35	58.73	0.174	43.1	26.58
Local patchy shadowing	8	424	1'161	36.52	35.79	15.64	62.63	1.426	75.4	28.40
Bilateral patchy shadowing	12	577	1'341	43.03	56.15	23.58	84.16	1.659	92.6	58.37
Nodular lesions	13	70	1'345	5.20	15.39	7.31	29.55	1.339	83.3	93.73
Air bronchogram	10	264	523	50.48	49.43	41.59	57.29	0.129	59.5	23.29
Pleural effusion	10	52	666	7.81	7.88	5.04	12.11	0.292	55.6	24.46
Reticulation / interlobular septal thickening	7	81	1'244	6.51	21.88	5.10	59.34	4.467	95.8	296.72
Interstitial abnormalities	5	163	1158	14.08	21.39	10.88	37.75	0.419	70.4	20.65
Crazy paving pattern	5	59	210	28.10	30.75	13.89	55.00	0.690	75.2	26.42
<b>Treatments</b>										
Antiviral treatment	57	4'475	6'068	73.75	92.74	85.65	96.47	5.031	98.4	2'064.73
Antibiotics	47	2'518	4'825	52.19	74.94	54.38	88.24	7.244	99.0	2'226.02
Corticosteroids	34	1'715	5'828	29.43	39.08	27.24	52.37	2.185	98.1	1'647.19
All mechanical ventilation	32	807	5'228	15.44	29.24	16.42	46.51	3.963	98.3	1'248.53
Invasive mechanical ventilation	25	238	3'506	6.79	8.84	4.39	16.97	2.969	95.6	356.53
High flow nasal cannula	20	1'298	2'745	47.29	47.39	27.93	67.67	2.654	98.3	499.24
Non-invasive mechanical ventilation	23	502	3'838	13.08	14.23	8.60	22.65	1.650	96.1	590.79
Intravenous immunoglobulin	20	781	3'162	24.70	21.67	15.47	29.50	0.070	94.0	486.21
Alpha interferon aerosol inhalation	15	367	745	49.26	89.41	55.01	98.31	6.313	97.4	331.79
Lopinavir	19	510	1'284	39.72	87.54	56.52	97.44	8.618	98.4	428.93

Ritonavir	19	510	1'284	39.72	87.54	56.52	97.44	8.618	98.4	428.93
Oxygen therapy	20	1300	1'872	69.44	83.83	72.76	90.96	1.519	95.2	406.24
Extracorporeal membrane oxygenation	22	31	4'651	0.67	0.51	0.16	1.63	4.517	86.9	93.79
Oseltamivir	13	443	1'159	38.22	96.39	41.42	99.9	9.269	91.8	89.74
Renal replacement therapy	18	62	4'572	1.36	1.35	0.48	3.78	4.010	92.7	154.71
Immune enhancing treatment	5	103	254	40.55	86.21	25.17	99.15	7.827	96.1	1'96.76
Antifungal treatment	5	70	1'516	4.62	6.81	3.68	12.28	0.401	81.7	32.66
<b>Outcome</b>										
Death	99	616	7'727	7.97	1.28	0.54	2.99	8.559	97.0	1'806.30
Survived	99	7'111	7'727	92.03	98.72	97.01	99.46	8.559	97.0	1'806.30
Discharged	56	1'751	5'401	32.42	52.15	35.25	68.58	5.257	98.5	2'161.24
Remained hospitalized	48	3'025	4'405	68.67	66.99	53.27	78.32	3.008	97.6	1'440.97
Recovery	34	1'012	3'741	27.05	53.76	32.35	73.87	5.495	98.5	1'685.14
<b>Complications</b>										
Admission to intensive care unit	23	195	2'877	6.78	9.68	5.41	16.73	1.685	91.4	314.57
Acute respiratory distress syndrome	27	759	5'122	14.82	22.97	12.69	37.94	3.121	98.0	1321.27
Shock	18	140	4'291	3.26	2.41	1.10	5.22	2.267	93.0	301.37
Acute kidney injury	18	241	4'113	5.86	7.17	3.75	13.28	1.889	95/0	335.95
Acute cardiac injury	13	242	1'250	19.36	13.54	8.58	20.72	0.631	88.2	109.82
All secondary infections	11	62	630	9.84	9.73	6.11	15.15	0.358	58.9	30.69
Respiratory failure	8	141	413	34.14	29.94	11.28	58.95	2.224	92.5	108.36
Pneumonia	7	1'031	1'489	69.24	33	8.36	72.68	4.67	98.2	476.86
Secondary infections (bacteria)	5	5	202	2.48	2.48	1.03	5.81	0	0	9.06
Heart failure	6	91	589	15.45	10.34	2.74	32.07	2.254	94.8	43.99

Table 5: Results of meta-analyses for patients with severe and non-severe disease outcome as well as survivors and non-survivors.

	Number of Studies	Number of events/ Number of severe	Number of events/ Number of non-severe	RR [95% CI]	p-value	Tau <sup>2</sup>	I <sup>2</sup>	Cochranes Q	Egger's test (p-value)
<b>Severe (cases) vs non-severe CoVID-19 disease (controls)</b>									
<b>Demographics</b>									
Sex: male	10	278/488	1059/1987	1.11 [1.01-1.22]	0.039	0.004	0%	7.67	0.763
Sex: female	10	210/488	925/1987	0.95 [0.82-1.10]	0.450	0.006	18.6%	11.05	0.395
Age	11	487	2059	SMD: 0.68 [0.40-0.97]	<0.001	0.154	81.8%	55.05	0.012
<b>Comorbidities</b>									
Any comorbidity	4	167/307	291/1205	2.11 [1.02-4.35]	0.046	0.160	79.8%	14.86	0.122
Hypertension	8	158/429	292/1734	2.15 [1.64-2.81]	<0.001	0.018	35.8%	10.91	0.664
Diabetes mellitus	7	84/427	127/1720	2.56 [1.50-4.39]	0.005	0.038	49.7%	11.92	0.279
Any heart condition	7	64/427	58/1720	4.09 [2.45-6.84]	<0.001	0.032	22.7%	7.76	0.548
COPD	6	23/403	15/1594	5.10 [3.08-8.45]	<0.001	0	0%	1.59	0.034
Carcinoma	5	15/345	19/1512	3.13 [0.63-15.64]	0.120	0.696	42.9%	7.00	0.339
<b>Symptoms and signs</b>									
Fever	8	399/462	1588/1847	1.02 [0.99-1.06]	0.187	<0.0001	41.3%	11.92	0.644
Fatigue	8	199/462	611/1847	1.21 [0.99-1.48]	0.059	0.004	46.0%	12.95	0.011
Myalgia	5	53/318	237/1454	1.01 [0.66-1.56]	0.929	<0.0001	20.7%	5.04	0.702
Headache	7	47/404	187/1765	1.14 [0.94-1.39]	0.146	<0.0001	0.0%	1.65	0.625
Cough	8	290/462	1051/1847	1.14 [1.02-1.27]	0.026	0.006	15.1%	8.25	0.633
Sputum	6	85/385	384/1549	1.05 [0.79-1.39]	0.460	<0.0001	14.8%	5.87	0.8731
Dyspnea	6	91/207	56/587	4.67 [0.99-21.91]	0.050	1.156	76.2%	21.03	0.148
Sore throat / Pharyngalgia	6	41/358	182/1549	1.40 [0.62-3.17]	0.337	0.218	50.9%	10.19	0.831
Diarrhea	6	41/403	77/1594	1.76 [0.72-4.32]	0.164	0.296	53.7%	10.80	0.384
<b>Treatment</b>									
Antibiotics	4	254/309	743/1410	1.63 [0.67-3.96]	0.177	0.285	93.5%	45.93	0.807
Antiviral treatment	6	249/347	888/1526	1.05 [0.90-1.22]	0.490	0.011	77.7%	22.45	0.604
Corticosteroids	5	200/345	416/1512	2.26 [1.32-3.87]	0.014	0.174	93.7%	63.66	0.211
<b>Imaging features (CT)</b>									
Pathological findings	7	400/416	1372/1631	1.06 [0.96-1.18]	0.192	0.009	90.1%	60.32	0.085
Pneumonia	5	373/389	1290/1539	1.05 [0.94-1.18]	0.299	0.008	92.1%	50.58	0.176

<b>Complications</b>									
Acute respiratory distress symptom (ARDS)	4	117/331	65/1457	10.59 [2.44-46.01]	0.014	0.606	84.1%	18.90	0.067
Acute kidney injury	4	16/331	8/1457	6.60 [0.37-116.33]	0.128	2.075	65.0%	8.56	0.909
Laboratory parameter	Number of studies	Number of severe	Number of non-severe	SMD [95% CI]	p-value	Tau <sup>2</sup>	I <sup>2</sup>	Cochranes Q	Egger's test (p-value)
Albumin	3	131	511	-1.60 [-2.97 - (-0.24)]	0.022	1.385	96%	50.01	0.790
Alanine aminotransferase (ALT)	6	184	695	0.27 [0.06-0.47]	0.011	0.014	22.1%	6.42	0.545
Aspartate transaminase (AST)	6	184	695	0.85 [0.61-1.09]	<0.001	0.031	36.5%	7.88	0.942
Creatinine	6	205	794	0.59 [0.12-1.07]	0.015	0.298	87.3%	39.30	0.501
C-reactive protein (CRP)	6	227	774	1.47 [0.88-2.07]	<0.001	0.487	91.2%	56.50	0.296
D-Dimer	4	143	361	0.55 [0.22-0.89]	0.001	0.066	59.4%	7.39	0.632
Hemoglobin	6	342	1618	-0.23 [-0.41- (-0.06)]	0.001	0.016	37.2%	7.96	0.927
Lactate dehydrogenase (LDH)	4	93	279	1.71 [1.08-2.34]	<0.001	0.294	77.3%	13.20	0.599
Leucocytes	7	412	1676	0.49 [-0.24-1.21]	0.187	0.905	97.0%	202.83	0.175
Lymphocytes	8	415	1703	-0.59 [-0.88 - (-0.30)]	<0.001	0.118	79.1%	33.54	0.986
Monocytes	3	59	239	-0.10 [-0.39- 0.19]	0.519	0	0%	0.58	0.180
Neutrophils	4	99	334	0.94 [0.27-1.61]	0.006	0.384	85.6%	20.8	0.409
Potassium	4	304	1437	-0.21 [-0.40 - (-0.02)]	0.034	0.015	41.2%	5.1	0.502
Procalcitonin	4	194	566	0.72 [0.06-1.38]	0.032	0.410	92.0%	37.55	0.848
Sodium	4	304	1437	-0.26 [-0.67-0.15]	0.201	0.137	86.3%	21.97	0.533
Thrombocytes	7	357	1621	-0.57 [-0.68-(-0.45)]	<0.001	0	0.0%	3.47	0.127
<b>Others</b>									
Time since onset of symptoms to admission	5	236	789	SMD: 0.14 [-0.12- 0.41]	0.291	0.056	64.8%	11.36	0.465


	Number of Studies	Number of events/ Number of non-survivors	Number of events/ Number of survivors	RR [95% CI]	p-value	Tau <sup>2</sup>	I <sup>2</sup>	Cochranes Q	Egger's test (p-value)
<b>Non-survivors (cases) vs survivors (controls)</b>									
<b>Demographics</b>									
Sex: male	7	236/340	326/617	1.32 [1.13-1.54]	0.005	0.002	21.8%	7.67	0.70
Sex: female	7	104/340	291/617	0.65 [0.53-0.83]	0.005	0	1.6%	6.10	0.54
Age	7	340	617	SMD: 1.25 [0.78-1.72]	<0.001	0.294	85.7%	41.97	0.012
<b>Comorbidities</b>									
Any comorbidity	6	207/308	234/597	1.69 [1.48-1.94]	<0.001	0	1	2.91	0.115
Hypertension	5	125/287	90/435	2.09 [1.65-2.64]	0.001	<0.001	0%	2.08	0.545
Diabetes mellitus	5	71/318	53/451	1.88 [1.26-2.81]	0.012	<0.001	0%	2.88	0.141
Any heart condition	5	48/318	15/451	3.95 [1.03-15.20]	0.047	.477	45.5%	7.35	0.666
Cerebrovascular disease	3	12/155	0/198	36.88 [8.50-160.04]	0.009	0	0%	0.07	0.305
Any lung disease	4	39/308	14/434	3.03 [0.61-15.04]	0.115	.429	49.8%	5.97	0.811
Carcinoma	5	12/318	8/451	2.26 [0.67-7.61]	0.136	0	0%	2.84	0.02
Current smoker	4	13/200	13/322	2.02 [0.61-6.72]	0.160	<0.001	0%	2.65	0.136
<b>Symptoms and signs</b>									
Fever	6	288/319	407/455	1.00 [0.95-1.05]	0.974	0	0%	4.9	0.022
Fatigue	3	109/276	129/414	1.24 [1.14-1.36]	0.009	0	0%	0.09	0.991
Myalgia	5	35/210	66/339	0.97 [0.61-1.55]	0.895	0.026	0%	3.14	0.385
Headache	4	19/255	29/301	0.83 [0.64-1.09]	0.120	0	0%	0.26	0.930
Cough	6	196/319	196/455	1.37 [0.58-3.24]	0.385	0.605	92.3%	64.86	0.389
Sputum	4	84/277	93/418	1.43 [0.65-3.15]	0.245	0.182	62.4%	7.99	0.886
Dyspnea	4	178/264	85/314	2.60 [0.58-11.65]	0.137	0.561	86%	21.49	0.611
Diarrhea	3	48/277	71/418	0.96 [0.38-2.43]	0.860	0.077	27.6%	2.76	0.838
<b>Treatment</b>									
Antibiotics	5	280/309	395/438	1.03 [0.99-1.07]	0.114	0	0%	2.09	0.293
Antiviral treatment	5	222/329	446/596	0.94 [0.79-1.13]	0.426	0.006	67.7%	12.38	0.260
Corticosteroids	4	229/308	227/434	1.29 [0.66-2.54]	0.321	0.136	80.6%	15.44	0.873
Immunoglobulin	4	143/308	122/434	1.88 [0.36-9.69]	0.309	0.979	92.5%	40.23	0.213
Oxygen nasal (high flow)	4	154/308	139/434	2.16 [0.09-50.50]	0.493	3.843	98.1%	158.98	0.03
All mechanical ventilation	5	298/319	115/455	6.05 [1.41-26.05]	0.026	1.126	84.5%	25.75	0.686
Non-invasive mech.	5	181/309	45/438	5.33 [1.52-18.71]	0.021	0.565	66.7%	12.02	0.765


ventilation									
Invasive mech. ventilation	5	89/309	5/438	14.14[138-145.09]	0.034	2.080	59.7%	9.92	0.181
Renal replacement therapy	4	22/200	1/322	10.36 [0.98-110.07]	0.051	0.194	0%	1.92	0.057
Extracorporeal membrane oxygenation (ECMO)	5	12/309	2/438	4.39 [1.64-11.78]	0.014	0	0%	1.35	0.033
<b>Imaging features (CT)</b>									
Pathological findings	6	562/577	325/335	0.97 [0.87-1.09]	0.588	0.006	75.9%	20.71	0.675
Pneumonia	3	159/168	254/302	1.07 [0.97-1.17]	0.089	<0.001	0%	1.34	0.680
<b>Complications</b>									
Acute respiratory distress symptom (ARDS)	6	298/319	115/455	4.24 [1.30-13.83]	0.026	1.115	92.8%	69.92	0.197
Shock	4	98/277	0/418	242.79 [23.70-2487.07]	0.005	0	0%	0.64	0.300
Acute cardiac injury	4	178/308	23/434	13.21 [0.70-248.38]	0.068	2.7831	81.8%	16.48	0.435
Acute kidney injury	5	88/309	5/435	20.77 [2.43-177.44]	0.017	2.301	67.7%	12.37	0.229
Laboratory parameter	Number of Studies	Number of cases	Number of controls	SMD [95% CI]	p-value	Tau <sup>2</sup>	I <sup>2</sup>	Cochranes Q	Egger's test (p-value)*
Albumin	2	110	120	-1.14 [-1.41 - (-0.85)]	<0.001	0	0%	0	n.a.
Alanine aminotransferase (ALT)	3	223	281	0.45 [0.08 - 0.82]	0.016	0.056	62.7%	6.37	0.984
Aspartate transaminase (AST)	2	114	165	0.17 [-0.07 - 0.41]	0.168	0	0%	0.76	n.a.
Creatinine	4	200	322	2.24 [-0.56 - 5.03]	0.117	7.719	98.8%	244.97	0.460
C-reactive protein (CRP)	2	114	165	0 [-0.24 - 0.24]	1.0	0	0%	0	n.a.
D-Dimer	4	174	274	1.54 [-0.17 - 3.25]	0.077	2.370	96.8%	94.99	0.672
Hemoglobin	3	142	140	-0.08 [-0.32 - 0.16]	0.504	0	0%	0.61	0.610
Lactate dehydrogenase (LDH)	2	110	120	1.61 [1.31 - 1.91]	<0.001	0	0%	0.3	n.a.
Leucocytes	4	277	418	2.21 [0.61 - 3.64]	0.006	1.989	97.9%	144.57	0.421
Lymphocytes	4	255	301	-0.92 [-1.3 - (-0.55)]	<0.001	0.079	64.6%	8.47	
Neutrophils	2	55	141	3.6 [3.12 - 4.08]	<0.001	0	0%	0.17	n.a.
Potassium	2	55	141	0.41 [0.1 - 0.77]	0.01	0	0%	0.01	n.a.
Thrombocytes	4	196	277	0.9 [-2.09 - 3.88]	0.556	8.916	99%	309.32	0.487
Partial thromboplastin time (PTT)	5	206	294	7.99 [4.64 - 11.34]	<0.001	13.245	98.9%	370.17	0.194
Activated partial thromboplastin time (APTT)	3	65	158	21.73 [4.34 - 39.13]	0.014	231.933	99.5%	363.82	0.386


Interleukin 6 (IL-6)	2	110	120	1.21 [0.93 – 1.5]	<0.001	0	0%	0.44	n.a.
<b>Others</b>									
Time since onset of symptoms to admission	3	195	273	0.47 [-0.09 – 1.02]	0.098	0.201	85.8%	14.05	0.797





\*Egger's test cannot be performed with less than three studies. Abbreviation: SMD: Standardize mean difference (negative number indicate lower values in cases, positive number indicate higher number in cases)



















**A****C****Survivors****D****Non-survivors**





A





B




C




D




A




B



C



D



# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.



# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.



# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.